



Ternary (σ, τ, ξ) -derivations on Banach ternary algebras

Madjid Eshaghi Gordji^{a,*}, Raziye Farrokhzad^b, Seyed Alireza Hosseinioun^c

^aDepartment of Mathematics, Semnan University, P. O. Box 35195-363, Semnan, Iran.

^bDepartment of Mathematics, Shahid Beheshti University, Tehran, Iran.

^c Department of Mathematical Sciences, University of Arkansas, Fayetteville, Arkansas 72701, USA.

Dedicated to the Memory of Charalambos J. Papaioannou

(Communicated by Th. M. Rassias)

Abstract

Let A be a Banach ternary algebra over a scalar field \mathbb{R} or \mathbb{C} and X be a Banach ternary A -module. Let σ, τ and ξ be linear mappings on A , a linear mapping $D : (A, []_A) \rightarrow (X, []_X)$ is called a ternary (σ, τ, ξ) -derivation, if

$$D([xyz]_A) = [D(x)\tau(y)\xi(z)]_X + [\sigma(x)D(y)\xi(z)]_X + [\sigma(x)\tau(y)D(z)]_X$$

for all $x, y, z \in A$.

In this paper, we investigate ternary (σ, τ, ξ) -derivation on Banach ternary algebras, associated with the following functional equation

$$f\left(\frac{x+y+z}{4}\right) + f\left(\frac{3x-y-4z}{4}\right) + f\left(\frac{4x+3z}{4}\right) = 2f(x).$$

Moreover, we prove the generalized Ulam–Hyers stability of ternary (σ, τ, ξ) -derivations on Banach ternary algebras.

Keywords: Banach ternary algebra; Banach ternary A -module; Ternary (σ, τ, ξ) -derivation.
2010 MSC: Mathematics Subject Classification. Primary 39B52; Secondary 39B82; 46B99; 17A40.

*Corresponding author

Email addresses: madjid.eshaghi@gmail.com (Madjid Eshaghi Gordji), raziyeh.farrokhzad@yahoo.com (Raziye Farrokhzad), shosseini@uark.edu (Seyed Alireza Hosseinioun)

1. Introduction

Ternary algebraic operations were considered in the 19 th century by several mathematicians such as A. Cayley [12] who introduced the notion of cubic matrix which in turn was generalized by Kapranov, Gelfand and Zelevinskii in 1990 ([30]). The simplest example of such non-trivial ternary operation is given by the following composition rule:

$$\{a, b, c\}_{ijk} = \sum_{l,m,n} a_{nil} b_{ljm} c_{mkn}, \quad i, j, k, \dots = 1, 2, \dots, N.$$

Ternary structures and their generalization, the so-called n-ary structures, raise certain hopes in view of their possible applications in physics. Some significant physical applications are as follows (see [3], [4], [8], [15]- [22], [24], [31], [32], [37], [54] and [56]). A ternary (associative) algebra $(A, [\])$ is a linear space A over a scalar field $\mathbb{F} = (\mathbb{R} \text{ or } \mathbb{C})$ equipped with a linear mapping, the so-called ternary product, $[\]: A \times A \times A \rightarrow A$ such that $[[abc]de] = [a[bcd]e]$ for all $a, b, c, d, e \in A$. This notion is a natural generalization of the binary case. Indeed if (A, \odot) is a usual (binary) algebra then $[abc] := (a \odot b) \odot c$ induced a ternary product making A into a ternary algebra which will be called trivial. It is known that unital ternary algebras are trivial and finitely generated ternary algebras are ternary subalgebras of trivial ternary algebras [9]. There are other types of ternary algebras in which one may consider other versions of associativity. Some examples of ternary algebras are (i) "cubic matrices" introduced by Cayley [12] which were in turn generalized by Kapranov, Gelfand and Zelevinskii [30]; (ii) the ternary algebra of polynomials of odd degrees in one variable equipped with the ternary operation $[p_1 p_2 p_3] = p_1 \odot p_2 \odot p_3$, where \odot denotes the usual multiplication of polynomials.

By a Banach ternary algebra we mean a ternary algebra equipped with a complete norm $\| \cdot \|$ such that $\| [abc] \| \leq \| a \| \| b \| \| c \|$. If a ternary algebra $(A, [\])$ has an identity, i.e. an element e such that $a = [aee] = [eae] = [eea]$ for all $a \in A$, then $a \odot b := [aeb]$ is a binary product for which we have

$$(a \odot b) \odot c = [[aeb]ec] = [ae[bec]] = a \odot (b \odot c)$$

and

$$a \odot e = [aee] = a = [eea] = e \odot a,$$

for all $a, b, c \in A$ and so $(A, [\])$ may be considered as a (binary) algebra. Conversely, if (A, \odot) is any (binary) algebra, then $[abc] := a \odot b \odot c$ makes A into a ternary algebra with the unit e such that $a \odot b = [aeb]$.

Let A be a Banach ternary algebra and X be a Banach space. Then X is called a ternary Banach A -module, if module operations $A \times A \times X \rightarrow X$, $A \times X \times A \rightarrow X$, and $X \times A \times A \rightarrow X$ are \mathbb{C} -linear in every variable. Moreover satisfy:

$$[[abc]_A dx]_X = [a[bcd]_A x]_X = [ab[cdx]_X]_X,$$

$$[abc]_A xd]_X = [a[bcx]_X d]_X = [ab[cxd]_X]_X,$$

$$[[xab]_X cd]_X = [x[abc]_A d]_X = [xa[bcd]_A]_X,$$

$$[[axb]_X cd]_X = [a[xbc]_X d]_X = [ax[bcd]_A]_X,$$

$$[[abx]_X cd]_X = [a[bxc]_X d]_X = [ab[xcd]_X]_X,$$

for all $x \in X$ and all $a, b, c, d \in A$, and

$$\max\{\|[xab]_X\|, \|[axb]_X\|, \|[abx]_X\|\} \leq \|a\|\|b\|\|x\|$$

for all $x \in X$ and all $a, b \in A$.

Let $(A, []_A)$ be a Banach ternary algebra over a scalar field \mathbb{R} or \mathbb{C} and $(X, []_X)$ be a ternary Banach A -module. A linear mapping $D : (A, []_A) \rightarrow (X, []_X)$ is called a ternary derivation, if

$$D([abc]_A) = [D(a)bc]_X + [aD(b)c]_X + [abD(c)]_X \quad (1.1)$$

for all $a, b, c \in A$.

Let σ, τ and ξ be linear mappings on A . A linear mapping $D : (A, []_A) \rightarrow (X, []_X)$ is called a ternary (σ, τ, ξ) -derivation, if

$$D([abc]_A) = [D(a)\tau(b)\xi(c)]_X + [\sigma(a)D(b)\xi(c)]_X + [\sigma(a)\tau(b)D(c)]_X \quad (1.2)$$

for all $a, b, c \in A$.

The stability of functional equations was first introduced by S. M. Ulam [55] in 1940. More precisely, let G_1 be a group, (G_2, d) be a metric group and ϵ be a positive number, S. M. Ulam asked, does there exist a $\delta > 0$ such that if a function $f : G_1 \rightarrow G_2$ satisfies the inequality $d(f(xy), f(x)f(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $T : G_1 \rightarrow G_2$ such that $d(f(x), T(x)) < \epsilon$ for all $x \in G_1$?. When this problem has a solution, we say that the homomorphism from G_1 to G_2 is stable.

In 1941, D. H. Hyers [27] gave a partial solution of *Ulam's* problem for the case of approximate additive mappings under the assumption that G_1 and G_2 are Banach spaces. In 1950, T. Aoki [1] was the second author to treat this problem for additive mappings (see also [11]). In 1978, Th. M. Rassias [48] generalized the theorem of Hyers by considering the stability problem with unbounded Cauchy differences. This phenomenon of stability that was introduced by Th. M. Rassias [48] is called the Hyers-Ulam-Rassias stability, according to J. M. Rassias Theorem, as follows:

Theorem 1.1. *Let $f : V \rightarrow W$ be a mapping from a norm vector space V into a Banach space W subject to the inequality*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p) \quad (1.3)$$

for all $x, y \in V$, where ϵ and p are constants with $\epsilon > 0$ and $p < 1$. Then there exists a unique additive mapping $T : V \rightarrow W$ such that

$$\|f(x) - T(x)\| \leq \frac{2\epsilon}{2-2^p} \|x\|^p \quad (1.4)$$

for all $x \in V$. If $p < 0$ then inequality (1.3) holds for all $x, y \neq 0$, and (1.4) for $x \neq 0$. Also, if the function $t \mapsto f(tx)$ from \mathbb{R} into W is continuous for each fixed $x \in V$, then T is linear.

On the other hand J. M. Rassias ([44]- [46]) generalized the Hyers stability result by presenting a weaker condition controlled by a product of different powers of norms. According to J. M. Rassias Theorem [51]:

Theorem 1.2. *If it is assumed that there exist constants $\Theta \geq 0$ and $p_1, p_2 \in \mathbb{R}$ such that $p = p_1 + p_2 \neq 1$, and $f : V \rightarrow W$ is a mapping from a norm space V into a Banach space W such that the inequality*

$$\|f(x + y) - f(x) - f(y)\| \leq \Theta \|x\|^{p_1} \|y\|^{p_2}$$

for all $x, y \in V$ holds, then there exists a unique additive mapping $T : V \rightarrow W$ such that

$$\|f(x) - T(x)\| \leq \frac{\Theta}{2 - 2^p} \|x\|^p,$$

for all $x \in V$. If in addition for every $x \in V$, $f(tx)$ is continuous in real t for each fixed x , then T is linear (see [38]-[46]).

During the last decades several stability problems of functional equations have been investigated by many mathematicians. A large list of references concerning the stability of functional equations can be found in [5], [6], [11], [13]-[24], [25], [28], [33], [47] and [49]-[53]. Recently, R. Badora , [7] and T. Miura et al. [34] proved the Ulam–Hyers stability, the Isac and Rassias–type stability [29], the Hyers–Ulam–Rassias stability and the Bourgin–type superstability of ring derivations on Banach algebras. On the other hand, C. Park [36] has contributed works to the stability problem of ternary homomorphisms and ternary derivations(see also [26]).

The main purpose of the present paper is to offer the Ulam–Hyers stability of ternary (σ, τ, ξ) -derivations on Banach ternary algebras subjected with the following functional equation

$$f\left(\frac{x + y + z}{4}\right) + f\left(\frac{3x - y - 4z}{4}\right) + f\left(\frac{4x + 3z}{4}\right) = 2f(x) . \tag{1.5}$$

2. Ternary (σ, τ, ξ) -derivations on Banach ternary algebras

In this section, we investigate ternary (σ, τ, ξ) -derivations on Banach ternary algebras. Throughout this section, assume that $(A, []_A)$ is a Banach ternary algebra and $(X, []_X)$ is a ternary Banach A -module.

Lemma 2.1. *Let V and W be linear spaces and let $f : V \rightarrow W$ be an additive mapping such that $f(\mu x) = \mu f(x)$ for all $x \in V$ and all $\mu \in \mathbb{T}^1(= \{\lambda \in \mathbb{C} ; |\lambda| = 1\})$. Then the mapping f is \mathbb{C} -linear. [35]*

Lemma 2.2. *Let $f : A \rightarrow X$ be a mapping such that*

$$f\left(\frac{\mu x + y + z}{4}\right) + f\left(\frac{3\mu x - y - 4z}{4}\right) + f\left(\frac{4\mu x + 3z}{4}\right) = 2\mu f(x), \tag{2.1}$$

for all $x, y, z \in A$ and $\mu \in \mathbb{T}^1$. Then f is \mathbb{C} -linear. [10]

The first result is as follows:

Theorem 2.3. *Let $p \neq 1$ and θ be nonnegative real numbers, and let $f : A \rightarrow X$ be a mapping and σ, τ , and ξ be linear mappings on A such that*

$$f\left(\frac{\mu x + y + z}{4}\right) + f\left(\frac{3\mu x - y - 4z}{4}\right) + f\left(\frac{4\mu x + 3z}{4}\right) = 2\mu f(x), \tag{2.2}$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$,

$$\|f([xyz]_A) - [f(x)\tau(y)\xi(z)]_X - [\sigma(x)f(y)\xi(z)]_X - [\sigma(x)\tau(y)f(z)]_X\| \leq \theta \|x\|^p \|y\|^p \|z\|^p \tag{2.3}$$

for all $x, y, z \in A$. Then the mapping $f : A \rightarrow X$ is a ternary (σ, τ, ξ) -derivation.

Proof . Assume $p < 1$. By Lemma 2.2, the mapping $f : A \rightarrow X$ is \mathbb{C} -linear. It follows from (2.3) that

$$\begin{aligned} & \|f([xyz]_A) - [f(x)\tau(y)\xi(z)]_X - [\sigma(x)f(y)\xi(z)]_X - [\sigma(x)\tau(y)f(z)]_X\| \\ &= \frac{1}{n^3} \|f([(nx)(ny)(nz)]_A) - [f(nx)\tau(ny)\xi(nz)]_X - [\sigma(nx)f(ny)\xi(nz)]_X - [\sigma(nx)\tau(ny)f(nz)]_X\| \\ &\leq \frac{\theta}{n^3} n^{3p} \|x\|^p \|y\|^p \|z\|^p \end{aligned}$$

for all $x, y, z \in A$. Thus, since $p < 1$, by letting n tend to ∞ in last inequality, we obtain

$$f([xyz]_A) = [f(x)\tau(y)\xi(z)]_X + [\sigma(x)f(y)\xi(z)]_X + [\sigma(x)\tau(y)f(z)]_X$$

for all $x, y, z \in A$. Hence, the mapping $f : A \rightarrow X$ is a ternary (σ, τ, ξ) -derivation. Similarly, one obtains the result for the case $p > 1$. \square

We prove the following Ulam stability problem for functional equation (1.5) controlled by the mixed type product-sum function

$$(x, y) \rightarrow \theta(\|x\|^{p_1} \|y\|^{p_2} \|z\|^{p_3} + \|x\|^p + \|y\|^p + \|z\|^p)$$

introduced by J. M. Rassias (see [47],[52]).

Theorem 2.4. *Let p, p_1, p_2, p_3 be real numbers such that $p < 1$, $p_1 + p_2 + p_3 < 1$, and $\theta > 0$. Suppose $f : A \rightarrow X$ is a mapping for which there exist mappings $g, h, k : A \rightarrow A$ whit $g(0) = h(0) = k(0) = 0$ such that*

$$\begin{aligned} & \|f\left(\frac{\mu x + y + z}{4}\right) + f\left(\frac{3\mu x - y - 4z}{4}\right) + f\left(\frac{4\mu x + 3z}{4}\right) - 2\mu f(x)\| \\ &\leq \theta(\|x\|^{p_1} \|y\|^{p_2} \|z\|^{p_3} + \|x\|^p + \|y\|^p + \|z\|^p), \end{aligned} \quad (2.4)$$

$$\|g(\lambda x + \lambda y) - \lambda g(x) - \lambda g(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.5)$$

$$\|h(\lambda x + \lambda y) - \lambda h(x) - \lambda h(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.6)$$

$$\|k(\lambda x + \lambda y) - \lambda k(x) - \lambda k(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.7)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$,

$$\|f([xyz]_A) - [f(x)h(y)k(z)]_X - [g(x)f(y)k(z)]_X - [g(x)h(y)f(z)]_X\| \leq \theta\|x\|^p \|y\|^p \|z\|^p \quad (2.8)$$

for all $x, y, z \in A$. Then there exist unique linear mappings σ, τ , and ξ from A to A and a unique ternary (σ, τ, ξ) -derivation $D : A \rightarrow X$ satisfying

$$\|g(x) - \sigma(x)\| \leq \theta \frac{2}{2 - 2^p} \|x\|^p \quad (2.9)$$

$$\|h(x) - \tau(x)\| \leq \theta \frac{2}{2 - 2^p} \|x\|^p \quad (2.10)$$

$$\|k(x) - \xi(x)\| \leq \theta \frac{2}{2 - 2^p} \|x\|^p \quad (2.11)$$

$$\|f(x) - D(x)\| \leq 2\theta \frac{2^p}{2 - 2^p} \|x\|^p \quad (2.12)$$

for all $x \in A$.

Proof . Setting $\mu = 1$ and $x = y = z = 0$ in (2.4), yields $f(0) = 0$. Let us take $\mu = 1$, $z = 0$ and $y = x$ in (2.4). Then we obtain

$$\|2f(\frac{x}{2}) - f(x)\| \leq 2\theta\|x\|^p, \quad (2.13)$$

for all $x \in A$. In (2.13), replacing $\frac{x}{2}$ by x and then dividing by 2, we get

$$\|f(x) - \frac{1}{2}f(2x)\| \leq 2^p\theta\|x\|^p,$$

for all $x \in A$. We easily prove that by induction that

$$\|f(x) - \frac{1}{2^n}f(2^n x)\| \leq 2\theta\|x\|^p \sum_{i=1}^n 2^{i(p-1)}. \quad (2.14)$$

In order to show that the functions $D_n(x) = \frac{1}{2^n}f(2^n x)$ form a convergent sequence, we use the Cauchy convergence criterion. Indeed, replace x by $2^m x$ and divide by 2^m in (2.14), where m is an arbitrary positive integer. We find that

$$\|\frac{1}{2^m}f(2^m x) - \frac{1}{2^{m+n}}f(2^{m+n} x)\| \leq 2\theta\|x\|^p \sum_{i=m+1}^{m+n} 2^{i(p-1)}$$

for all positive integers. Hence, by the Cauchy criterion the limit $D(x) = \lim_{n \rightarrow \infty} D_n(x)$ exists for each $x \in A$. By taking the limit as $n \rightarrow \infty$ in (2.12),

$$\|f(x) - D(x)\| \leq 2\theta\|x\|^p \sum_{i=1}^{\infty} 2^{i(p-1)}$$

and (2.12) holds for all $x \in A$. Now, we have

$$\begin{aligned} & \|D(\frac{\mu x + y + z}{4}) + D(\frac{3\mu x - y - 4z}{4}) + D(\frac{4\mu x + 3z}{4}) - 2\mu D(x)\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{2^n} \|f(\frac{2^n \mu x + 2^n y + 2^n z}{4}) + f(\frac{3 \cdot 2^n \mu x - 2^n y - 4 \cdot 2^n z}{4}) \\ &+ f(\frac{4 \cdot 2^n \mu x + 3 \cdot 2^n z}{4}) - 2\mu f(2^n x)\|_A \leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \theta(\|2^n x\|^{p_1} \|2^n y\|^{p_2} \|2^n z\|^{p_3} \\ &+ \|2^n x\|^p + \|2^n y\|^p + \|2^n z\|^p) = \lim_{n \rightarrow \infty} 2^{n(p_1+p_2+p_3-1)} \theta(\|x\|^{p_1} \|y\|^{p_2} \|z\|^{p_3}) \\ &+ \lim_{n \rightarrow \infty} 2^{n(p-1)} \theta(\|x\|^p + \|y\|^p + \|z\|^p) = 0 \end{aligned}$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$. Hence

$$D(\frac{\mu x + y + z}{4}) + D(\frac{3\mu x - y - 4z}{4}) + D(\frac{4\mu x + 3z}{4}) = 2\mu D(x)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$. So by Lemma 2.2, D is \mathbb{C} -linear.

Also put $\lambda = 1$ in (2.5) to obtain

$$\|g(x + y) - g(x) - g(y)\| \leq \theta(\|x\|^p + \|y\|^p) \tag{2.15}$$

fix $x \in A$. Replacing y by x and then dividing by 2 in (2.15), we get

$$\|\frac{1}{2}g(2x) - g(x)\| \leq \theta\|x\|^p$$

one can use the induction to show that

$$\|\frac{1}{2^{m+n}}g(2^{m+n}x) - \frac{1}{2^m}g(2^m x)\| \leq \theta\|x\|^p \sum_{i=m}^{m+n-1} 2^{(i-1)(p-1)} \tag{2.16}$$

for all $x \in A$. It follows from the convergence of series (2.16) that the sequence $\{\frac{g(2^n x)}{2^n}\}$ is cauchy. Hence, the limit $\sigma(x) = \lim_{n \rightarrow \infty} \frac{g(2^n x)}{2^n}$ exists for all $x \in A$. we easily prove that by (2.5) that $\sigma(\lambda x + \lambda y) = \lambda\sigma(x) + \lambda\sigma(y)$ and by (2.16) that

$$\|g(x) - \sigma(x)\| \leq \theta\|x\|^p \sum_{i=1}^{\infty} 2^{(i-1)(p-1)}$$

and (2.9) holds for all $x \in A$. Similarly there exist linear mappings τ and ξ on A such that (2.10) and (2.11) hold for all $x \in A$. On the other hand

$$\begin{aligned} & \|D([xyz]_A) - [D(x)\tau(y)\xi(z)]_X - [\sigma(x)D(y)\xi(z)]_X - [\sigma(x)\tau(y)D(x)]_X\| \\ &= \lim_{n \rightarrow \infty} \frac{1}{8^n} \|f([(2^n x)(2^n y)(2^n z)]_A) - [f(2^n x)h(2^n y)k(2^n z)]_X \\ & \quad - [g(2^n x)f(2^n y)k(2^n z)]_X - [g(2^n x)h(2^n y)f(2^n z)]_X\| \\ &\leq \lim_{n \rightarrow \infty} \frac{\theta}{8^n} \|2^n x\|^p \|2^n y\|^p \|2^n z\|^p \\ &= \lim_{n \rightarrow \infty} \theta 8^{n(p-1)} \|x\|^p \|y\|^p \|z\|^p \\ &= 0 \end{aligned}$$

for all $x, y, z \in A$, which means that

$$D([xyz]_A) = [D(x)\tau(t)\xi(z)]_X + [\sigma(x)D(y)\xi(z)]_X + [\sigma(x)\tau(y)D(z)]_X.$$

Therefore, we conclude that D is a ternary (σ, τ, ξ) -derivation. Suppose that there exists another ternary (σ, τ, ξ) -derivation $D' : A \rightarrow X$ satisfying (2.12). Since $D'(x) = \frac{1}{2^n}D'(2^n x)$, we see that

$$\begin{aligned} \|D(x) - D'(x)\| &= \frac{1}{2^n} \|D(2^n x) - D'(2^n x)\| \\ &\leq \frac{1}{2^n} (\|f(2^n x) - D(2^n x)\| + \|f(2^n x) - D'(2^n x)\|) \\ &\leq 4\theta \frac{2^p}{2-2^p} 2^{n(p-1)} \|x\|^p, \end{aligned}$$

which tends to zero as $n \rightarrow \infty$ for all $x \in A$. Therefore $D' = D$ as claimed and similarly we can prove that σ, τ and ξ are unique on A and the proof of the theorem is complete. \square

Theorem 2.5. *Let p, p_1, p_2, p_3 be real numbers such that $p > 1$, $p_1 + p_2 + p_3 > 1$, and $\theta > 0$. Suppose $f : A \rightarrow X$ is a mapping for which there exist mappings g, h, k on A with $g(0) = h(0) = k(0) = 0$ such that*

$$\begin{aligned} & \left\| f\left(\frac{\mu x + y + z}{4}\right) + f\left(\frac{3\mu x - y - 4z}{4}\right) + f\left(\frac{4\mu x + 3z}{4}\right) - 2\mu f(x) \right\| \\ & \leq \theta(\|x\|^{p_1} \|y\|^{p_2} \|z\|^{p_3} + \|x\|^p + \|y\|^p + \|z\|^p), \end{aligned} \quad (2.17)$$

$$\|g(\lambda x + \lambda y) - \lambda g(x) - \lambda g(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.18)$$

$$\|h(\lambda x + \lambda y) - \lambda h(x) - \lambda h(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.19)$$

$$\|k(\lambda x + \lambda y) - \lambda k(x) - \lambda k(y)\| \leq \theta(\|x\|^p + \|y\|^p) \quad (2.20)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$,

$$\|f([xyz]_A) - [f(x)h(y)k(z)]_X - [g(x)f(y)k(z)]_X - [g(x)h(y)f(z)]_X\| \leq \theta\|x\|^p\|y\|^p\|z\|^p \quad (2.21)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$. Then there exist unique linear mappings σ, τ and k on A and a unique ternary (σ, τ, ξ) -derivation $D : A \rightarrow X$ satisfying

$$\|g(x) - \sigma(x)\| \leq \theta \frac{2}{2^p - 2} \|x\|^p \quad (2.22)$$

$$\|h(x) - \tau(x)\| \leq \theta \frac{2}{2^p - 2} \|x\|^p \quad (2.23)$$

$$\|k(x) - \xi(x)\| \leq \theta \frac{2}{2^p - 2} \|x\|^p \quad (2.24)$$

$$\|D(x) - f(x)\| \leq 2\theta \frac{2^p}{2^p - 2} \|x\|^p \quad (2.25)$$

for all $x \in A$.

Proof . Setting $\mu = 1$ and $x = y = z = 0$ in (2.17), yields $f(0) = 0$. Let us take $\mu = 1$, $z = 0$ and $y = x$ in (2.17). Then we obtain

$$\|2f\left(\frac{x}{2}\right) - f(x)\| \leq 2\theta\|x\|^p, \quad (2.26)$$

for all $x \in A$. By induction, we get

$$\|2^n f\left(\frac{x}{2^n}\right) - f(x)\| \leq 2\theta\|x\|^p \sum_{i=0}^{n-1} 2^{i(1-p)}. \quad (2.27)$$

In order to show that the functions $D_n(x) = 2^n f\left(\frac{x}{2^n}\right)$ form a convergent sequence, we use the Cauchy convergence criterion. Indeed, replace x by $\frac{x}{2^m}$ and multiply by 2^m in (2.27), where m is an arbitrary positive integer. We find that

$$\|2^{m+n} f\left(\frac{x}{2^{m+n}}\right) - 2^m f\left(\frac{x}{2^m}\right)\| \leq 2\theta\|x\|^p \sum_{i=m}^{m+n-1} 2^{i(1-p)}$$

for all positive integers. Hence, by the Cauchy criterion the limit $D(x) = \lim_{n \rightarrow \infty} D_n(x)$ exists for each $x \in A$. By taking the limit as $n \rightarrow \infty$ in (2.27) we see that

$$\|D(x) - f(x)\| \leq 2\theta \|x\|^p \sum_{i=0}^{\infty} 2^{i(1-p)}$$

and (2.25) holds for all $x \in A$. Thus, we have

$$\begin{aligned} & \|D(\frac{\mu x + y + z}{4}) + D(\frac{3\mu x - y - 4z}{4}) + D(\frac{4\mu x + 3z}{4}) - 2\mu D(x)\| \\ &= \lim_{n \rightarrow \infty} 2^n \|f(\frac{2^{-n}\mu x + 2^{-n}y + 2^{-n}z}{4}) + f(\frac{3 \cdot 2^{-n}\mu x - 2^{-n}y - 4 \cdot 2^{-n}z}{4}) \\ &+ f(\frac{4 \cdot 2^{-n}\mu x + 3 \cdot 2^{-n}z}{4}) - 2\mu f(2^{-n}x)\| \\ &\leq \lim_{n \rightarrow \infty} 2^n \theta (\|2^{-n}x\|^{p_1} \|2^{-n}y\|^{p_2} \|2^{-n}z\|^{p_3} + \|2^{-n}x\|^p + \|2^{-n}y\|^p + \|2^{-n}z\|^p) \\ &= \lim_{n \rightarrow \infty} 2^{n(1-p_1+p_2+p_3)} \theta (\|x\|^{p_1} \|y\|^{p_2} \|z\|^{p_3}) + \lim_{n \rightarrow \infty} 2^{n(1-p)} \theta (\|x\|^p + \|y\|^p + \|z\|^p) \\ &= 0 \end{aligned}$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$. Hence,

$$D(\frac{\mu x + y + z}{4}) + D(\frac{3\mu x - y - 4z}{4}) + D(\frac{4\mu x + 3z}{4}) = 2\mu D(x)$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$. So by Lemma 2.2, D is \mathbb{C} -linear.

Also put $\lambda = 1$ in (2.18) to obtain

$$\|g(x + y) - g(x) - g(y)\| \leq \theta (\|x\|^p + \|y\|^p) \quad (2.28)$$

fix $x \in A$. Replacing y by x and then replacing x by $\frac{x}{2}$ in (2.28), we get

$$\|2g(\frac{x}{2}) - g(x)\| \leq 2\theta \|\frac{x}{2}\|^p$$

one can use the induction to show that

$$\|2^{m+n}g(\frac{x}{2^{m+n}}) - 2^m g(\frac{x}{2^m})\| \leq \theta \|x\|^p \sum_{i=m+1}^{m+n} 2^{i(1-p)} \quad (2.29)$$

for all $x \in A$. It follows from the convergence of series (2.29) that the sequence $\{2^n g(\frac{x}{2^n})\}$ is cauchy. Hence the limit $\sigma(x) = \lim_{n \rightarrow \infty} 2^n g(\frac{x}{2^n})$ exists for all $x \in A$. we easily prove that by (2.18) that $\sigma(\lambda x + \lambda y) = \lambda \sigma(x) + \lambda \sigma(y)$ and by (2.29) that

$$\|g(x) - \sigma(x)\| \leq \theta \|x\|^p \sum_{i=1}^{\infty} 2^{i(1-p)}$$

and (2.22) holds for all $x \in A$. Similarly there exist linear mappings τ and ξ on A such that (2.23) and (2.24) hold for all $x \in A$. Thus, we have

$$\begin{aligned} & \|D([xyz]_A) - [D(x)\tau(y)\xi(z)]_X - [\sigma(x)D(y)\xi(z)]_X - [\sigma(x)\tau(y)D(z)]_X\| \\ &= \lim_{n \rightarrow \infty} 8^n \| (f[(2^{-n}x)(2^{-n}y)(2^{-n}z)]_A) - [f(2^{-n}x)h(2^{-n}y)k(2^{-n}z)]_X \\ & \quad - [g(2^{-n}x)f(2^{-n}y)k(2^{-n}z)]_X - [g(2^{-n}x)h(2^{-n}y)f(2^{-n}z)]_X \| \\ & \leq \lim_{n \rightarrow \infty} 8^n \theta \| \frac{x}{2^n} \|^p \| \frac{y}{2^n} \|^p \| \frac{z}{2^n} \|^p = \lim_{n \rightarrow \infty} \theta 8^{n(1-p)} \|x\|^p \|y\|^p \|z\|^p \\ &= 0 \end{aligned}$$

for all $x \in A$, which means that

$$D([xyz]_A) = [D(x)\tau(y)\xi(z)]_X + [\sigma(x)D(y)\xi(z)]_X + [\sigma(x)\tau(y)D(z)]_X.$$

Therefore, we conclude that D is a ternary (σ, τ, ξ) -derivation. Suppose that there exists another ternary (σ, τ, ξ) -derivation $D' : A \rightarrow X$ satisfying (2.25). Since $D'(x) = 2^n D'(\frac{x}{2^n})$, we see that

$$\begin{aligned} \|D(x) - D'(x)\| &= 2^n \|D(\frac{x}{2^n}) - D'(\frac{x}{2^n})\| \\ &\leq 2^n (\|f(\frac{x}{2^n}) - D(\frac{x}{2^n})\| + \|f(\frac{x}{2^n}) - D'(\frac{x}{2^n})\|) \\ &\leq 4\theta \frac{2^p}{2^{p-2}} 2^{n(1-p)} \|x\|^p, \end{aligned}$$

which tends to zero as $n \rightarrow \infty$ for all $x \in A$. Hence, $D' = D$ as claimed and similarly we can prove that σ, τ and ξ are unique on A and proof of theorem is complete. \square

We are going to investigate the Hyers–Ulam–Rassias stability problem for functional equation (1.5).

Corollary 2.6. *Let $P \in (-\infty, 1) \cup (1, \infty)$, $\theta > 0$. Suppose $f : A \rightarrow X$ is a mapping for which there exist mappings g, h, k on A with $g(0) = h(0) = k(0) = 0$ such that*

$$\|f(\frac{\mu x + y + z}{4}) + f(\frac{3\mu x - y - 4z}{4}) + f(\frac{4\mu x + 3z}{4}) - 2\mu f(x)\| \leq \theta(\|x\|^p + \|y\|^p + \|z\|^p),$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$,

$$\|g(\lambda x + \lambda y) - \lambda g(x) - \lambda g(y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

$$\|h(\lambda x + \lambda y) - \lambda h(x) - \lambda h(y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

$$\|k(\lambda x + \lambda y) - \lambda k(x) - \lambda k(y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

for all $\lambda \in \mathbb{T}^1$ and all $x, y \in A$,

$$\|f([xyz]_A) - [f(x)h(y)k(z)]_X - [g(x)f(y)k(z)]_X - [g(x)h(y)f(z)]_X\| \leq \theta\|x\|^p\|y\|^p\|z\|^p$$

for all $x, y, z \in A$. Then there exist unique linear mappings σ, τ and k on A and a unique ternary (σ, τ, ξ) -derivation $D : A \rightarrow X$ satisfying

$$\|g(x) - \sigma(x)\| \leq \theta \frac{2}{|2^p - 2|} \|x\|^p$$

$$\|h(x) - \tau(x)\| \leq \theta \frac{2}{|2^p - 2|} \|x\|^p$$

$$\|k(x) - \xi(x)\| \leq \theta \frac{2}{|2^p - 2|} \|x\|^p$$

$$\|D(x) - f(x)\| \leq 2\theta \frac{2^p}{|2^p - 2|} \|x\|^p$$

for all $x \in A$.

By Theorems 2.4 and 2.5 we solve the following Hyers–Ulam stability problem for functional equation (1.5).

Corollary 2.7. *Let θ be a positive real number. Suppose $f : A \rightarrow X$ is a mapping for which there exist mappings g, h, k on A with $g(0) = h(0) = k(0) = 0$ such that*

$$\|f\left(\frac{\mu x + y + z}{4}\right) + \mu f\left(\frac{3\mu x - y - 4z}{4}\right) + f\left(\frac{4\mu x + 3z}{4}\right) - 2\mu f(x)\| \leq \theta,$$

for all $\mu \in \mathbb{T}^1$ and all $x, y, z \in A$,

$$\|g(\lambda x + \lambda y) - \lambda g(x) - \lambda g(y)\| \leq \theta$$

$$\|h(\lambda x + \lambda y) - \lambda h(x) - \lambda h(y)\| \leq \theta$$

$$\|k(\lambda x + \lambda y) - \lambda k(x) - \lambda k(y)\| \leq \theta$$

for all $\lambda \in \mathbb{T}^1$ and all $x, y \in A$,

$$\|f([xyz]_A) - [f(x)h(y)k(z)]_X - [g(x)f(y)k(z)]_X - [g(x)h(y)f(z)]_X\| \leq \theta$$

for all $x, y, z \in A$. Then there exist unique linear mappings σ, τ and k on A and a unique ternary (σ, τ, ξ) -derivation $D : A \rightarrow X$ satisfying

$$\|g(x) - \sigma(x)\| \leq \frac{\theta}{2}$$

$$\|h(x) - \tau(x)\| \leq \frac{\theta}{2}$$

$$\|k(x) - \xi(x)\| \leq \frac{\theta}{2}$$

$$\|D(x) - f(x)\| \leq \theta$$

for all $x \in A$.

References

- [1] V. Abramov, R. Kerner and B. Le Roy, Hypersymmetry a Z_3 graded generalization of supersymmetry, *J. Math. Phys.* 38, 1650 (1997).
- [2] M. Amyari, Stability of Generalized Lie (σ, τ) -Derivations, *Tamsui Oxford J. Math.* 24 (4), (2008) 389-399.
- [3] H. Azadi Kenary, M. Ghirati, C. Park and M. Eshaghi Gordji, Homomorphisms and derivations in induced fuzzy C^* -algebras, *Journal of Inequalities and Applications*, 2013, 2013:88, 12 pp.
- [4] H. Azadi Kenary, A. R. Zohdi and M. Eshaghi Gordji, Fixed points and stability for quartic mappings in β -normed left Banach modules on Banach algebras, *Acta Mathematica Scientia* 2013, 33B(4), 1113-1118.
- [5] T. Aoki, On the stability of the linear transformation in Banach spaces, *J. Math. Soc. Japan.* 2(1950), 64-66.
- [6] C. Baak and M. S. Moslehian, Stability of J^* -homomorphisms, *Nonlinear Analysis (TAM)*, 63 (2005), 42-48.
- [7] R. Badora, On approximate derivations, *Math. Inequal. Appl.* 9 (2006), 167-173.
- [8] F. Bagarello and G. Morchio, Dynamics of mean-field spin models from basic results in abstract differential equations, *J. Stat. Phys.* 66 (1992) 849-866. MR1151983 (93c:82034)
- [9] N. Bazunova, A. Borowiec and R. Kerner, Universal differential calculus on ternary algebras, *Lett. Math. Phys.* 67 (2004), no. 3, 195-206.
- [10] M. Bavand Savadkouhi, M. Eshaghi Gordji, J. M. Rassias, N. Ghobadipour, Approximate ternary Jordan derivations on Banach ternary algebras, *J. Math. Phys.* 50 (2009), no. 4, 042303, 9 pp.
- [11] D. G. Bourgin, Approximately isometric and multiplicative transformations on continuous function rings, *Duke Math. J.* 16, (1949). 385-397.
- [12] A. Cayley, On the 34 concomitants of the ternary cubic, *Am. J. Math.* 4, 1 (1881).
- [13] P. W. Cholewa, Remarks on the stability of functional equations, *Aequationes Math.* 27 (1984) 76-86. MR0758860 (86d:39016)
- [14] S. Czerwik, Stability of functional equations of Ulam-Hyers-Rassias type, *Hadronic Press*, 2003.
- [15] A. Ebadian, S. Kaboli Gharetapeh and M. Eshaghi Gordji, Nearly Jordan $*$ -homomorphisms between unital C^* -algebras, *Abs. Appl. Anal.*, Art.ID: 513128, (2011), 11 pages.
- [16] M. Eshaghi Gordji, Nearly involutions on Banach algebras; A fixed point approach, *Fixed Point Theory*, 14, No.1, 2013, 117-124.
- [17] M. Eshaghi Gordji and Z. Alizadeh, Stability and superstability of ring homomorphisms on non-Archimedean Banach algebras, *Abs. Appl. Anal.*, Vol. 2011, Article ID:123656, (2011), 10 pages.
- [18] M. Eshaghi Gordji, M. B. Ghaemi, H. Majani, C. Park and A. Najati, Fixed points and approximately heptic mappings in non-Archimedean normed spaces, *Adv. Difference Equ.* 2013, 2013:209, 10 pp.
- [19] M. Eshaghi Gordji, H. Khodaei and R. Khodabakhsh, On approximate n -ary derivations, *Int. J. Geom. Methods Mod. Phys.*, Vol8, No 3, (2011), 485-500.
- [20] M. Eshaghi Gordji, H. Khodaei and Th. M. Rassias, Fixed points and stability for quadratic mappings in β -normed left Banach modules on Banach algebras, *Results Math.*, 2011, DOI 10.1007/s00025-011-0123-z.
- [21] M. E. Gordji and H. Khodaei, A fixed point technique for investigating the stability of (α, β, γ) -derivations on Lie C^* -algebras, *Nonlinear Analysis-TMA* 76 (2013), 52-57.
- [22] M. Eshaghi Gordji, M. B. Savadkouhi and M. Bidkham, Additive-cubic functional equations in non-Archimedean spaces, *Filomat*, 27, 2013, No.5, 731-738.
- [23] Z. Gajda, On stability of additive mappings, *Internat. J. Math. Math. Sci.* 14(1991) 431-434.
- [24] R. Haag and D. Kastler, An algebraic approach to quantum field theory, *J. Math. Phys.* 5 (1964) 848-861. MR0165864
- [25] F. Hassani, A. Ebadian, M. Eshaghi Gordji and H. Azadi Kenary, Nearly n -homomorphisms and n -derivations in fuzzy ternary Banach algebras, *Journal of Inequalities and Applications*, 2013, 2013:71, 11 pp.
- [26] O. Hatori and J. Wada, Ring derivations on semi-simple commutative Banach algebras, *Tokyo J. Math.* 15 (1992), 223-229.
- [27] D. H. Hyers, On the stability of the linear functional equation, *Proc. Natl. Acad. Sci.* 27 (1941) 222-224.
- [28] D. H. Hyers, G. Isac and Th. M. Rassias, Stability of functional Equations in Several Variables, *Birkhauser, Boston, Basel, Berlin*, 1998.
- [29] G. Isac and Th. M. Rassias, On the Hyers-Ulam stability of n -additive mappings, *J. Approx. Theorey* 72 (1993), 131-137.
- [30] M. Kapranov, I. M. Gelfand and A. Zelevinskii, Discriminants, *Resultants and Multidimensional Determinants*, *Birkhauser, Berlin*, 1994.
- [31] R. Kerner, Ternary algebraic structures and their applications in physics, *Univ. P. M. Curie preprint, Paris* (2000), <http://arxiv.org/list/math-ph/0011>.
- [32] R. Kerner, The cubic chessboard: *Geometry and physics, Class. Quantum Grav.* 14, A203 (1997).

- [33] H. Khodaei, M. Eshaghi Gordji and R. Khodabakhsh, Fixed points, Lie *-homomorphisms and Lie *-derivations on lie C*-algebras, *Fixed Point Theory*, 14, No.2, 2013, 387-400.
- [34] T. Miura, G. Hirasawa and S. E. Takahasi, A perturbation of ring derivations on Banach algebras, *J. Math. Anal. Appl.* 319 (2006), 522-530.
- [35] C. Park, Homomorphisms between Poisson JC*-algebras, *Bull. Braz. Math. Soc.* 36 (2005) 79-97. MR2132832 (2005m:39047)
- [36] C. Park, Isomorphisms between C*-ternary algebras, *J. Math. Anal. Appl.* 327 (2007), 101-115.
- [37] C. Park, J. L. Cui, and M. Eshaghi Gordji, Orthogonality and quintic functional equations, *Acta Math. Sin. (Engl. Ser.)* 29 (2013), no. 7, 1381-1390.
- [38] J. M. Rassias, Complete solution of the multi-dimensional problem of Ulam. *Discuss. Math.* 14 (1994), 101-107.
- [39] J. M. Rassias, On the stability of a multi-dimensional Cauchy type functional equation. *Geometry, analysis and mechanics*, 365-375, World Sci.Publ., River Edge, NJ, 1994.
- [40] J. M. Rassias, Solution of a stability problem of Ulam. *Functional analysis, approximation theory and numerical analysis*, 241-249, World Sci.Publ., River Edge, NJ, 1994.
- [41] J. M. Rassias, Solution of a stability problem of Ulam, *Discuss. Math.* 12 (1992), 95-103 (1993).
- [42] J. M. Rassias, Solution of a problem of Ulam, *J. Approx. Theory* 57 (1989), no. 3, 268-273.
- [43] J. M. Rassias, On a new approximation of approximately linear mappings by linear mappings, *Discuss. Math.* 7 (1985), 193-196.
- [44] J. M. Rassias, On approximation of approximately linear mappings by linear mappings, *Bull. Sci. Math. (2)* 108 (1984), no. 4, 445-446.
- [45] J. M. Rassias, On approximation of approximately linear mappings by linear mappings, *J.Funct. Anal.* 46 (1982), no. 1, 126-130.
- [46] J. M. Rassias, Solution of the Ulam stability problem for quartic mappings, *Glas. Mat. Ser. III* 34(54) (1999), no. 2, 243-252.
- [47] J. M. Rassias and H. M. Kim, Approximate homomorphisms and derivations between C*-ternary algebras. *J. Math. Phys.* 49 (2008), no. 6, 063507, 10 pp. 46Lxx (39B82)
- [48] Th. M. Rassias, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.* 72 (1978) 297-300.
- [49] Th. M. Rassias, On the stability of functional equations and a problem of Ulam, *Acta Math. Appl.* 62 (2000) 23-130. MR1778016 (2001j:39042)
- [50] Th. M. Rassias, On the stability of functional equations in Banach spaces, *J. Math. Anal. Appl.* 251 (2000) 264-284. MR1790409 (2003b:39036)
- [51] Th. M. Rassias, The problem of S.M.Ulam for approximately multiplicative mappings, *J. Math. Anal. Appl.* 246(2)(2000),352-378.
- [52] K. Ravi, M. Arunkumar and J. M. Rassias, Ulam stability for the orthogonally general Euler-Lagrange type functional equation, *Int. J. Math. Stat.* 3 (2008), A08, 36-46. 39B55 (39B82)
- [53] K. Ravi, J. M. Rassias and B.V. Senthil Kumar, Ulam stability of reciprocal difference and adjoint functional equations, *To appear in Australian Journal of Mathematical Analysis and applications*
- [54] G. L. Sewell, Quantum Mechanics and its Emergent Macrophysics, *Princeton Univ. Press, Princeton, NJ*, 2002. MR1919619 (2004b:82001)
- [55] S. M. Ulam, Problems in Modern Mathematics, *Chapter VI, science ed. Wiley, New York*, 1940.
- [56] H. Zettl, A characterization of ternary rings of operators, *Adv. Math.* 48 (1983) 117-143. MR0700979 (84h:46093)