



Multi-point boundary value problems of higher-order nonlinear fractional differential equations

İsmail Yaslan*

Department of Mathematics, Pamukkale University 20070 Denizli, Turkey

(Communicated by Madjid Eshaghi Gordji)

Abstract

We investigate the existence and uniqueness of solutions for multi-point nonlocal boundary value problems of higher-order nonlinear fractional differential equations by using some well known fixed point theorems.

Keywords: boundary value problems; fractional derivative; fixed point theorems.
2010 MSC: Primary 26A33; Secondary 34B10, 34B27.

1. Introduction

Fractional differential equations have been of great interest recently. This is due to the intensive development of the theory of fractional calculus itself as well as its applications. They arise in various fields of science and engineering such as the mathematical modeling of systems and processes in the fields of physics, chemistry, biology, aerodynamics, porous structures and polymer rheology [6, 15, 20]. Therefore, in recent years, the study of the boundary value problems for fractional differential equations has received considerable attention (see, for instance, [2, 4, 5, 8, 9, 10, 16, 19, 23, 25, 26, 27, 28, 30, 31, 32, 34] and references therein). However, few papers have considered the multi-point boundary value problems for higher-order fractional differential equations (see [7, 13, 17, 21, 29, 33]).

Ahmad and Nieto [1] studied some existence results in a Banach space for a nonlocal fractional boundary value problem given by

$$\begin{cases} {}^C D^q(x(t)) = f(t, x(t)), & t \in (0, 1), \quad q \in (m-1, m], \quad m \geq 2, \\ x(0) = x'(0) = \dots = x^{(m-2)}(0) = 0, & x(1) = \alpha x(\eta), \quad 0 < \eta < 1, \end{cases}$$

*Corresponding author

Email address: iyaslan@pau.edu.tr (İsmail Yaslan)

where ${}^C D^\alpha$ is the Caputo fractional derivative.

Salem [22] investigated the existence of Pseudo solutions for the nonlinear m -point fractional boundary value problem

$$\begin{cases} D_{0+}^\alpha(x(t)) + q(t)f(t, x(t)) = 0, & t \in (0, 1), \quad \alpha \in (n-1, n], \quad n \geq 2, \\ x(0) = x'(0) = \dots = x^{(n-2)}(0) = 0, & x(1) = \sum_{i=1}^{m-2} \zeta_i x(\eta_i), \end{cases}$$

where $0 < \eta_1 < \dots < \eta_{m-2} < 1$, $\zeta_i > 0$ with $\sum_{i=1}^{m-2} \zeta_i \eta_i^{\alpha-1} < 1$.

Jia and Zhang [12] consider the multi-point boundary value problem of nonlinear fractional differential equation

$$\begin{cases} D_{0+}^\alpha u(t) + \lambda f(t, u(t)) = 0, & 0 < t < 1, \quad \alpha \geq 2, \quad n-1 < \alpha \leq n, \\ u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0, & u^{(i)}(1) = \sum_{j=0}^{m-2} \eta_j u'(\xi_j) ds, \quad 0 \leq i \leq n-2. \end{cases}$$

Ahmad and Ntouyas [3] are concerned with the existence of solutions for a fractional boundary value problem

$$\begin{cases} {}^C D^q(x(t)) = f(t, x(t)), & t \in (0, T), \quad 1 \leq q \leq 2, \\ \alpha_1 x(0) + \beta_1 ({}^C D^q(x(0))) = \gamma_1, \\ \alpha_2 x(1) + \beta_2 ({}^C D^q(x(1))) = \gamma_2, & 0 < p < 1. \end{cases}$$

Liu and Jia [18] investigated the nonlinear boundary value problem of fractional differential equation

$$\begin{cases} {}^C D^q(x(t)) = f(t, x(t), x'(t)), & t \in (0, 1), \quad q \in (n-1, n], \quad n > 2, \\ g_0(x(0), x'(0)) = 0, \\ g_1(x(1), x'(1)) = 0, \\ x''(0) = x'''(0) = \dots = x^{(n)}(0) = 0. \end{cases}$$

In [24], authors developed sufficient conditions for multiplicity of positive solutions to the boundary value problem

$$\begin{cases} D_{0+}^q(u(t)) + f(t, u(t)) = 0, & t \in (0, 1), \\ u^{(i)}(0) = 0, & 0 \leq i \leq n-2, \\ u(1) = \sum_{i=1}^{m-2} \delta_i u(\eta_i), \end{cases}$$

where $n-1 < q \leq n$ and D_{0+}^q is the standard Riemann-Liouville fractional derivative of order q , $n \geq 3$, $\delta_i, \eta_i \in (0, 1)$ with $\sum_{i=1}^{m-2} \delta_i \eta_i^{q-1} < 1$ and $f : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ is continuous.

Jiang and Wang [14] studied the existence and uniqueness of solutions to the following boundary value problem for fractional differential equation

$$\begin{cases} D_{0+}^\alpha u(t) + f(t, u(t), I_{0+}^\beta u(t)) = 0, & 0 \leq t \leq 1, \\ u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0, & D_{0+}^p u(1) = \sum_{i=1}^m a_i D_{0+}^q u(\xi_i), \end{cases}$$

where D_{0+}^α is the Riemann-Liouville fractional derivative of order $n-1 < \alpha \leq n$, $n \geq 2$, $0 < \beta < 1$, $p \in [1, n-2]$, $q \in [0, p]$, $0 < \xi_1 < \xi_2 < \dots < \xi_m < 1$, $f : [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$ is a continuous function and $a_i > 0$ ($i = 1, 2, \dots, m$).

Motivated by the aforementioned works, this paper is concerned with the existence of solutions to multi-point boundary value problem for higher order fractional differential equations:

$$\left\{ \begin{array}{l} -{}^R D_{0+}^{\alpha}(u(t)) + f(t, u(t)) = 0, \quad t \in [0, 1], \\ u''(0) = u'''(0) = \dots = u^{(n-2)}(0) = 0, \quad u''(1) = \sum_{i=1}^{m-2} \gamma_i u''(\xi_i), \\ au(0) - bu'(0) = \sum_{i=1}^{m-2} a_i u'(\xi_i), \\ cu(1) + du'(1) = \sum_{i=1}^{m-2} b_i u'(\xi_i), \end{array} \right. \quad (1.1)$$

where ${}^R D_{0+}^{\alpha}$ is the Riemann-Liouville fractional derivative of order α . Throughout the paper we suppose that $m \geq 3$, $n \geq 4$, $n - 1 < \alpha \leq n$ where $n, m \in \mathbb{N}$, $a, b, c, d > 0$, $a_i, b_i \geq 0$, $0 < \xi_1 < \dots < \xi_{m-2} < 1$ and $0 \leq \sum_{i=1}^{m-2} \gamma_i \xi_i^{n-1} < 1$. We assume that $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous. Also, we consider the analogous problem using the Caputo fractional derivative:

$$\left\{ \begin{array}{l} -{}^C D_{0+}^{\alpha}(u(t)) + f(t, u(t)) = 0, \quad t \in [0, 1], \\ u''(0) = u'''(0) = \dots = u^{(n-2)}(0) = 0, \quad u''(1) = \sum_{i=1}^{m-2} \gamma_i u''(\xi_i), \\ au(0) - bu'(0) = \sum_{i=1}^{m-2} a_i u'(\xi_i), \\ cu(1) + du'(1) = \sum_{i=1}^{m-2} b_i u'(\xi_i). \end{array} \right. \quad (1.2)$$

We mean a function $u \in C[0, 1]$ of class $C^n[0, 1]$ which satisfies the nonlocal fractional boundary value problem (1.1) (or (1.2)) by a solution of (1.1) (or (1.2)).

We have organized the paper as follows. First, we provide some definitions and preliminary lemmas which are key tools for our main results. Second, we obtain some existence and uniqueness results of the Riemann-Liouville multi-point boundary value problem (RLMBVP) (1.1) and the Caputo multi-point boundary value problem (CMBVP) (1.2).

We assume that the following conditions are satisfied:

(H1) If $m \geq 3$, then $c \sum_{i=1}^{m-2} a_i \geq a \sum_{i=1}^{m-2} b_i$ and if $m > 3$, then $ad > c \sum_{i=1}^{j-1} a_i \geq a \sum_{i=1}^{j-1} b_i > bc$ where

$$2 \leq j \leq m - 2.$$

(H2) $ad > a \sum_{i=1}^{m-2} b_i + c \sum_{i=1}^{m-2} a_i$.

2. Preliminaries

To state the main results of this paper, we will need the following lemmas and we present some notation.

Definition 2.1. ([15]) The Riemann-Liouville fractional derivative of order $\alpha > 0$ for a function $u : (0, \infty) \rightarrow \mathbb{R}$ is defined by

$${}^R D_{0+}^{\alpha} u(t) = \frac{1}{\Gamma(n - \alpha)} \frac{d^n}{dt^n} \int_0^t (t - s)^{n-\alpha-1} u(s) ds$$

where $n = [\alpha] + 1$, $[\alpha]$ denotes the integer part of the number α , provided that the right side is pointwise defined on $(0, \infty)$.

Definition 2.2. ([15]) The Riemann-Liouville fractional integral of order $\alpha > 0$ of a function $u : (0, \infty) \rightarrow \mathbb{R}$ is given by

$$I_{0+}^{\alpha}u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1}u(s)ds$$

provided the integral exists.

Lemma 2.3. ([15]) Let $\alpha > 0$. Then the following equality holds for $u \in L(0, 1)$, ${}^R D_{0+}^{\alpha}u \in L(0, 1)$;

$$I_{0+}^{\alpha} {}^R D_{0+}^{\alpha}u(t) = u(t) + c_1 t^{\alpha-1} + c_2 t^{\alpha-2} + \dots + c_n t^{\alpha-n},$$

$c_i \in \mathbb{R}$, $i = 1, \dots, n$, where $n - 1 < \alpha \leq n$.

Definition 2.4. ([15]) The fractional derivative of a function u in the Caputo sense is defined as

$${}^C D_{0+}^{\alpha}u(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1}u^{(n)}(s)ds$$

where $n = [\alpha] + 1$, $[\alpha]$ denotes the integer part of the number α , provided that the right side is pointwise defined on $(0, \infty)$.

Lemma 2.5. ([15]) Let $n - 1 < \alpha \leq n$, $u \in C^n[0, 1]$. Then

$$I_{0+}^{\alpha} {}^C D_{0+}^{\alpha}u(t) = u(t) + c_0 + c_1 t + c_2 t^2 + \dots + c_{n-1} t^{n-1},$$

for $c_i \in \mathbb{R}$, $i = 1, \dots, n$.

In the following, the RLMBVP (1.1) will be reduced to an equivalent integral equation. We know that ${}^R D_{0+}^{\alpha}(u(t)) = {}^R D_{0+}^{\alpha-2}({}^R D_{0+}^2 u(t)) = {}^R D_{0+}^{\alpha-2}(u''(t))$. If $-u''(t) = y(t)$ and $\alpha - 2 = q$, then the problem

$$\begin{cases} -{}^R D_{0+}^{\alpha-2}(u''(t)) + f(t, u(t)) = 0, & t \in [0, 1] \\ u''(0) = u'''(0) = \dots = u^{(n-2)}(0) = 0, & u''(1) = \sum_{i=1}^{m-2} \gamma_i u''(\xi_i) \end{cases}$$

is turned into problem

$$\begin{cases} {}^R D_{0+}^q y(t) + f(t, u(t)) = 0, & t \in [0, 1] \\ y(0) = y'(0) = \dots = y^{(n-4)}(0) = 0, & y(1) = \sum_{i=1}^{m-2} \gamma_i y(\xi_i). \end{cases} \tag{2.1}$$

Lemma 2.6. The boundary value problem (2.1) has a unique solution

$$y(t) = \int_0^1 H_1(t, s)f(s, u(s))ds$$

where

$$H_1(t, s) = \frac{1}{K_1 \Gamma(q)} \begin{cases} (1-s)^{q-1}t^{q-1} - \sum_{i=k}^{m-2} \gamma_i t^{q-1}(\xi_i - s)^{q-1}, & t \leq s, s \in J_k, k = 1, 2, \dots, m-1; \\ (1-s)^{q-1}t^{q-1} - \sum_{i=k}^{m-2} \gamma_i t^{q-1}(\xi_i - s)^{q-1} - K_1(t-s)^{q-1}, & t \geq s, s \in J_k, k = 1, 2, \dots, m-1, \end{cases}$$

and $K_1 = 1 - \sum_{i=1}^{m-2} \gamma_i \xi_i^{q-1}$.

Proof . According to Lemma 2.3, we can obtain that

$$y(t) = -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s)) ds + c_1 t^{q-1} + c_2 t^{q-2} + \dots + c_{n-2} t^{q-n+2}.$$

By boundary conditions of (2.1) we get $c_2 = c_3 = \dots = c_{n-2} = 0$ and

$$c_1 = \frac{1}{K_1 \Gamma(q)} \left[\int_0^1 (1-s)^{q-1} f(s, u(s)) ds - \sum_{i=1}^{m-2} \gamma_i \int_0^{\xi_i} (\xi_i - s)^{q-1} f(s, u(s)) ds \right].$$

Thus, the unique solution of problem (2.1) is

$$\begin{aligned} y(t) &= -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s)) ds + \frac{1}{K_1 \Gamma(q)} \int_0^1 (1-s)^{q-1} t^{q-1} f(s, u(s)) ds \\ &\quad - \frac{1}{K_1 \Gamma(q)} \sum_{i=1}^{m-2} \gamma_i \int_0^{\xi_i} (\xi_i - s)^{q-1} t^{q-1} f(s, u(s)) ds \\ &= \int_0^1 H_1(t, s) f(s, u(s)) ds. \end{aligned}$$

The proof is complete.

□

Lemma 2.7. Let $D := ac + ad + bc + \sum_{i=1}^{m-2} (ca_i - ab_i)$ and $J_1 = [0, \xi_1]$, $J_2 = [\xi_1, \xi_2], \dots, J_{m-2} = [\xi_{m-3}, \xi_{m-2}]$, $J_{m-1} = [\xi_{m-2}, 1]$. For $y \in C[0, 1]$, the boundary value problem

$$\begin{cases} -u''(t) = y(t), & t \in [0, 1], \\ au(0) - bu'(0) = \sum_{i=1}^{m-2} a_i u'(\xi_i) \\ cu(1) + du'(1) = \sum_{i=1}^{m-2} b_i u'(\xi_i) \end{cases} \quad (2.2)$$

has a unique solution

$$u(t) = \int_0^1 G(t, s) y(s) ds \quad (2.3)$$

where

$$G(t, s) = \frac{1}{D} \begin{cases} (at + b + \sum_{i=1}^{k-1} a_i)(c(1-s) + d - \sum_{i=k}^{m-2} b_i) + \sum_{i=k}^{m-2} a_i [c(t-s) + \sum_{i=1}^{k-1} b_i], \\ t \leq s, s \in J_k, k = 1, 2, \dots, m-1; \\ (as + b + \sum_{i=1}^{k-1} a_i)(c(1-t) + d - \sum_{i=k}^{m-2} b_i) + \sum_{i=1}^{k-1} b_i [a(t-s) + \sum_{i=k}^{m-2} a_i], \\ t \geq s, s \in J_k, k = 1, 2, \dots, m-1. \end{cases} \quad (2.4)$$

Proof . A direct calculation gives that if $y \in C[0, 1]$, then the boundary value problem (2.2) has the unique solution

$$\begin{aligned}
 u(t) = & - \int_0^t (t-s)y(s)ds + \frac{t}{D} \left\{ a \int_0^1 (c(1-s) + d)y(s)ds + \sum_{i=1}^{m-2} (ca_i - ab_i) \int_0^{\xi_i} y(s)ds \right\} \\
 & + \frac{1}{D} \left\{ (b + \sum_{i=1}^{m-2} a_i) \int_0^1 (c(1-s) + d)y(s)ds - (b + \sum_{i=1}^{m-2} a_i) \sum_{i=1}^{m-2} b_i \int_0^{\xi_i} y(s)ds \right. \\
 & \left. + \frac{1}{a} \left(c(b + \sum_{i=1}^{m-2} a_i) - D \right) \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} y(s)ds \right\}.
 \end{aligned}$$

Hence, we obtain (2.3). \square

Lemma 2.8. *The Green’s function $G(t, s)$ in (2.4) satisfies*

$$0 < G(t, s) \leq G(s, s)$$

for $(t, s) \in [0, 1] \times [0, 1]$.

Proof . From (H1) and (H2), we have $G(t, s) > 0$.

Now, we will show that $G(t, s) \leq G(s, s)$.

- (i) Let $s \in J_k, 1 \leq k \leq m - 2$ and $t \leq s$. Since $G(t, s)$ is increasing in t , we get $G(t, s) \leq G(s, s)$.
- (ii) Take $s \in J_k, 1 \leq k \leq m - 2$ and $t \geq s$. From (H1), $G(t, s)$ is decreasing in t . So we obtain $G(t, s) \leq G(s, s)$.

\square

From Lemma 2.6 and Lemma 2.7, we know that $u(t)$ is a solution of the problem (1.1) if and only if

$$u(t) = \int_0^1 G(t, s) \int_0^1 H_1(s, \tau) f(\tau, u(\tau)) d\tau ds. \tag{2.5}$$

Now, let $E = C[0, 1]$, with supremum norm $\|y\| = \sup_{t \in [0,1]} |y(t)|$ for any $y \in E$. We can define the operator $A : E \rightarrow E$ by

$$Au(t) = \int_0^1 G(t, s) \int_0^1 H_1(s, \tau) f(\tau, u(\tau)) d\tau ds, \tag{2.6}$$

where $u \in E$. Therefore solving (2.5) in E is equivalent to finding fixed points of the operator A .

Now, the CMBVP (1.2) will be reduced to an equivalent integral equation. We know that ${}^C D_{0+}^\alpha (u(t)) = {}^C D_{0+}^{\alpha-2} ({}^C D_{0+}^2 u(t)) = {}^C D_{0+}^{\alpha-2} (u''(t))$. If $-u''(t) = y(t)$ and $\alpha - 2 = q$, then the problem

$$\begin{cases} -{}^C D_{0+}^{\alpha-2}(u''(t)) + f(t, u(t)) = 0, & t \in [0, 1] \\ u''(0) = u'''(0) = \dots = u^{(n-2)}(0) = 0, & u''(1) = \sum_{i=1}^{m-2} \gamma_i u''(\xi_i) \end{cases}$$

is turned into problem

$$\begin{cases} {}^C D_{0+}^q y(t) + f(t, u(t)) = 0, & t \in [0, 1] \\ y(0) = y'(0) = \dots = y^{(n-4)}(0) = 0, & y(1) = \sum_{i=1}^{m-2} \gamma_i y(\xi_i). \end{cases} \quad (2.7)$$

Lemma 2.9. *The boundary value problem (2.7) has a unique solution*

$$y(t) = \int_0^1 H_2(t, s) f(s, u(s)) ds$$

where

$$H_2(t, s) = \frac{1}{K_2 \Gamma(q)} \begin{cases} (1-s)^{q-1} t^{n-1} - \sum_{i=k}^{m-2} \gamma_i t^{n-1} (\xi_i - s)^{q-1}, \\ t \leq s, s \in J_k, k = 1, 2, \dots, m-1; \\ (1-s)^{q-1} t^{n-1} - \sum_{i=k}^{m-2} \gamma_i t^{n-1} (\xi_i - s)^{q-1} - K_2 (t-s)^{q-1}, \\ t \geq s, s \in J_k, k = 1, 2, \dots, m-1, \end{cases}$$

$$\text{and } K_2 = 1 - \sum_{i=1}^{m-2} \gamma_i \xi_i^{n-1}.$$

Proof . According to Lemma 2.5, we can obtain that

$$y(t) = -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s)) ds + c_0 + c_1 t + \dots + c_{n-1} t^{n-1}.$$

By boundary conditions of (2.7) we get $c_0 = c_1 = \dots = c_{n-2} = 0$ and

$$c_{n-1} = \frac{1}{K_2 \Gamma(q)} \left[\int_0^1 (1-s)^{q-1} f(s, u(s)) ds - \sum_{i=1}^{m-2} \gamma_i \int_0^{\xi_i} (\xi_i - s)^{q-1} f(s, u(s)) ds \right].$$

Thus, the unique solution of problem (2.7) is

$$\begin{aligned} y(t) &= -\frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} f(s, u(s)) ds + \frac{1}{K_2 \Gamma(q)} \int_0^1 (1-s)^{q-1} t^{n-1} f(s, u(s)) ds \\ &\quad - \frac{1}{K_2 \Gamma(q)} \sum_{i=1}^{m-2} \gamma_i \int_0^{\xi_i} (\xi_i - s)^{q-1} t^{n-1} f(s, u(s)) ds \\ &= \int_0^1 H_2(t, s) f(s, u(s)) ds. \end{aligned}$$

The proof is complete.

□

From Lemma 2.7 and Lemma 2.9, we know that $u(t)$ is a solution of the problem (1.2) if and only if

$$u(t) = \int_0^1 G(t, s) \int_0^1 H_2(s, \tau) f(\tau, u(\tau)) d\tau ds. \tag{2.8}$$

We can define the operator $F : C[0, 1] \rightarrow C[0, 1]$ by

$$Fu(t) = \int_0^1 G(t, s) \int_0^1 H_2(s, \tau) f(\tau, u(\tau)) d\tau ds,$$

where $u \in C[0, 1]$. Therefore solving (2.8) in $C[0, 1]$ is equivalent to finding fixed points of the operator F .

3. Existence and uniqueness of solutions

In this section, first, we will use the following well-known contraction mapping theorem named also as the Banach fixed point theorem: Let E be a Banach space and S a nonempty closed subset of E . Assume $A : S \rightarrow S$ is a contraction, i.e., there is a λ ($0 < \lambda < 1$) such that $\|Ax - Ay\| \leq \lambda\|x - y\|$ for all x, y in S . Then A has a unique fixed point in S .

Theorem 3.1. *We assume that the function $f(t, x)$ satisfies the following Lipschitz condition (H3) There is a constant $L > 0$ such that*

$$|f(t, x) - f(t, y)| \leq L|x - y| \text{ for all } t \in [0, 1] \text{ and } x, y \in C[0, 1].$$

If we have

$$\frac{L}{K_1\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} ds < 1,$$

then the RLMBVP (1.1) has a unique solution in $C[0, 1]$.

Proof . For $u_1, u_2 \in C[0, 1]$ and $t \in [0, 1]$, we have

$$\begin{aligned} |(Au_1)(t) - (Au_2)(t)| &\leq \int_0^1 G(t, s) \int_0^1 |H_1(s, \tau)| |f(\tau, u_1(\tau)) - f(\tau, u_2(\tau))| d\tau ds \\ &\leq \int_0^1 G(s, s) \int_0^1 \frac{(1 - s)^{q-1}}{K_1\Gamma(q)} L |u_1(\tau) - u_2(\tau)| d\tau ds \\ &\leq \left(\frac{L}{K_1\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} ds \right) \|u_1 - u_2\| \end{aligned}$$

by using Lemma 2.8, Lemma 2.6 and the condition (H3). Hence, A is a contraction mapping and the theorem is proved. \square

Theorem 3.2. *Suppose that (H3) holds. Moreover,*

$$\frac{L}{K_2\Gamma(q)} \int_0^1 G(s, s)(1-s)^{q-1} ds < 1.$$

Then, the CMBVP (1.2) has a unique solution in $C[0, 1]$.

Proof . The proof of Theorem 3.2 is very similar to that of Theorem 3.1 and therefore omitted.

□

In the next theorem, the function $f(t, x)$ satisfies a Lipschitz condition on a subset of $C[0, 1]$.

Theorem 3.3. *We assume that*

(H4) *Let there exists a number $r > 0$ such that*

$$|f(t, x) - f(t, y)| \leq L|x - y| \text{ for each } t \in [0, 1],$$

for all x and y in $S = \{x \in C[0, 1] : \|x\| \leq r\}$, where $L > 0$ is a constant which may depend on r .

Also,

$$\frac{L}{K_1\Gamma(q)} \int_0^1 G(s, s)(1-s)^{q-1} ds < 1.$$

Suppose that there exists nonnegative function $g \in C[0, 1]$ such that $|f(t, u(t))| \leq g(t) |u(t)|$ and

$$\frac{1}{K_1\Gamma(q)} \int_0^1 G(s, s)(1-s)^{q-1} \int_0^1 g(\tau) d\tau ds \leq 1.$$

Then, the RLMBVP (1.1) has a unique solution $u \in C[0, 1]$ with $|u(t)| \leq r, \forall t \in [0, 1]$.

Proof . Let us take $S = \{u \in C[0, 1] : \|u\| \leq r\}$. Obviously, S is a closed subset of $C[0, 1]$. Let $A : C[0, 1] \rightarrow C[0, 1]$ be the operator defined by (2.6). For u_1 and u_2 in S , taking into account (H4), in exactly the same way in the proof of Theorem 3.1 we can get $\|Au_1 - Au_2\| \leq \lambda \|u_1 - u_2\|$, where $0 < \lambda < 1$.

It remains to show that A maps S into itself. If $u \in S$, then we obtain

$$\begin{aligned} |(Au)(t)| &\leq \int_0^1 G(t, s) \int_0^1 |H_1(s, \tau)| |f(\tau, u(\tau))| d\tau ds \\ &\leq \int_0^1 G(s, s) \int_0^1 \frac{(1-s)^{q-1}}{K_1\Gamma(q)} g(\tau) |u(\tau)| d\tau ds \\ &\leq \|u\|. \end{aligned}$$

Since $\|Au\| \leq r$, we have $A : S \rightarrow S$.

From the contraction mapping theorem, the RLMBVP (1.1) has a unique solution $u \in C[0, 1]$ with $|u(t)| \leq r, \forall t \in [0, 1]$.

□

Theorem 3.4. *We assume that (H4) holds. Also,*

$$\frac{L}{K_2\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} ds < 1.$$

Suppose that there exists nonnegative function $g \in C[0, 1]$ such that $|f(t, u(t))| \leq g(t) |u(t)|$ and

$$\frac{1}{K_2\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} \int_0^1 g(\tau) d\tau ds \leq 1.$$

Then, the CMBVP (1.2) has a unique solution $u \in C[0, 1]$ with $|u(t)| \leq r, \forall t \in [0, 1]$.

Proof . The proof of Theorem 3.4 is very similar to that of Theorem 3.3 and therefore omitted. \square

To get an existence theorem without uniqueness of solution, we will apply the following Leray-Schauder nonlinear alternative theorem.

Theorem 3.5. [11] *Let E be a real Banach space and Ω be a bounded open subset of $E, 0 \in \Omega, F : \bar{\Omega} \rightarrow E$ be a completely continuous operator. Then either there exist $x \in \partial\Omega, \lambda > 1$ such that $F(x) = \lambda x$, or there exists a fixed point $x^* \in \bar{\Omega}$.*

For convenience, we introduce the following notation. Let

$$B_1 = \frac{1}{K_1\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} \int_0^1 g(\tau) d\tau ds,$$

$$B_2 = \frac{1}{K_2\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} \int_0^1 g(\tau) d\tau ds,$$

$$D_1 = \frac{1}{K_1\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} \int_0^1 h(\tau) d\tau ds,$$

$$D_2 = \frac{1}{K_2\Gamma(q)} \int_0^1 G(s, s)(1 - s)^{q-1} \int_0^1 h(\tau) d\tau ds.$$

Theorem 3.6. *Suppose that there exist nonnegative functions $g, h \in C[0, 1]$ such that $|f(t, u(t))| \leq g(t) |u(t)| + h(t)$ and $B_1 < 1$. Then the RLMBVP (1.1) has at least one solution $u \in C[0, 1]$.*

Proof . Since $B_1 < 1$ and $D_1 > 0, r := D_1(1 - B_1)^{-1} > 0$. Let us take $\Omega = \{u \in C[0, 1] : \|u\| < r\}$. By Arzela-Ascoli theorem, it is easy to check that $A : \bar{\Omega} \rightarrow C[0, 1]$ is completely continuous operator.

If we take $u \in \partial\Omega$, $\lambda > 1$ such that $Au = \lambda u$, then

$$\begin{aligned} \lambda D_1(1 - B_1)^{-1} &= \lambda r = \lambda \|u\| = \|Au\| = \max_{t \in [0,1]} |Au(t)| \\ &\leq \max_{t \in [0,1]} \int_0^1 G(t, s) \int_0^1 |H_1(s, \tau)| |f(\tau, u(\tau))| d\tau ds \\ &\leq \int_0^1 G(s, s) \int_0^1 \frac{(1-s)^{q-1}}{K_1 \Gamma(q)} (g(\tau) |u(\tau)| + h(\tau)) d\tau ds \\ &\leq B_1 \|u\| + D_1 = B_1 r + D_1 = D_1(1 - B_1)^{-1}. \end{aligned}$$

Hence we obtain $\lambda \leq 1$, this contradicts $\lambda > 1$. By Theorem 3.5, the operator A has a fixed point in $\bar{\Omega}$. Thus, the RLMBVP (1.1) has at least one solution $u \in \bar{\Omega}$.

□

Theorem 3.7. *Suppose that there exist nonnegative functions $g, h \in C[0, 1]$ such that $|f(t, u(t))| \leq g(t) |u(t)| + h(t)$ and $B_2 < 1$. Then the CMBVP (1.2) has at least one solution $u \in C[0, 1]$.*

Proof . The proof of Theorem 3.7 is very similar to that of Theorem 3.6 and therefore omitted.

□

Example 3.8. *Consider the following nonlinear RLMBVP*

$$\begin{cases} -{}^R D_{0+}^{\frac{11}{2}}(u(t)) + 2 + t^2 u(t) = 0, & t \in [0, 1], \\ u''(0) = u'''(0) = u^{(4)}(0) = 0, & u''(1) = u''(\frac{1}{2}), \\ u(0) - u'(0) = u'(\frac{1}{2}), \\ 2u(1) + 3u'(1) = u'(\frac{1}{2}). \end{cases} \quad (3.1)$$

Taking $a = b = a_1 = b_1 = \gamma_1 = 1$, $\xi_1 = \frac{1}{2}$, $c = 2$, $d = m = 3$, $n = 6$, $\alpha = \frac{11}{2}$ and $q = \frac{7}{2}$, we obtain $K_1 = 1 - \frac{1}{4\sqrt{2}}$ and $G(s, s) = \frac{1}{4}(-s^2 + s + 2)$. Since we have

$$|f(t, x) - f(t, y)| = |t^2 x(t) - t^2 y(t)| \leq |x(t) - y(t)|, \quad \forall t \in [0, 1],$$

we take $L = 1$. Also, we obtain

$$\frac{L}{K_1 \Gamma(q)} \int_0^1 G(s, s)(1-s)^{q-1} ds \approx 0,055875 < 1.$$

Hence, by Theorem 3.1, the RLMBVP (3.1) has a unique solution in $C[0, 1]$.

References

- [1] B. Ahmad and JJ. Nieto, *Existence of solutions for nonlocal boundary value problems of higher-order nonlinear fractional differential equations*, Abstr. Appl. Anal. Art. ID 494720 (2009) 1–9.
- [2] B. Ahmad, JJ. Nieto and A. Alsaedi, *Existence and uniqueness of solutions for nonlinear fractional differential equations with non-separated type integral boundary conditions*, Acta Math. Sci. 31 (2011) 2122–2130.

- [3] B. Ahmad and SK. Ntouyas, *Fractional differential inclusions with fractional separated boundary conditions*, *Fract. Calc. Appl. Anal.* 15 (2012) 362–382.
- [4] B. Ahmad and SK. Ntouyas, *Initial value problems of fractional order Hadamard-type functional differential equations*, *Electron. J. Differ. Equ.* 2015 (2015) No. 77.
- [5] AS. Berdyshev, A. Cabada and BK. Turmetov, *On solvability of a boundary value problem for a nonhomogeneous biharmonic equation with a boundary operator of a fractional order*, *Acta Math. Sci. Ser. B Engl. Ed.* 34 (2014) 1695–1706.
- [6] S. Das, *Functional fractional calculus for system identification and controls*, New York, Springer, 2008.
- [7] M. El-Shahed and JJ. Nieto, *Nontrivial solutions for a nonlinear multi-point boundary value problem of fractional order*, *Comput. Math. Appl.* 59 (2010) 3438–3443.
- [8] FT. Fen, *On the solutions of a multi-point BVP for fractional differential equations with impulses*, *Miskolc Math. Notes* 19 (2018) 847–863.
- [9] FT. Fen, IY. Karaca and OB. Ozen, *Positive solutions of boundary value problems for p -Laplacian fractional differential equations*, *Filomat* 31 (2017) 1265–1277.
- [10] M. Günendi and İ. Yaslan, *Positive solutions of higher-order nonlinear multi-point fractional equations with integral boundary conditions*, *Fract. Calc. Appl. Anal.* 19 (2016) 989–1009.
- [11] D. Guo and V. Lakshmikantham, *Nonlinear problems in abstract cones*, Orlando, Academic Press, 1988.
- [12] Y. Jia and X. Zhang, *Positive solutions for a class of fractional differential equation multi-point boundary value problems with changing sign nonlinearity*, *J. Appl. Math. Comput.* 47 (2015) 15–31.
- [13] M. Jia, X. Zhang and X. Gu, *Nontrivial solutions for a higher fractional differential equation with fractional multi-point boundary conditions*, *Bound. Value Probl.* 70 (2012) 1–16.
- [14] J. Jiang and H. Wang, *Existence and uniqueness of solutions for a fractional differential equation with multi-point boundary value problems*, *Journal of Applied Analysis and Computation* 9 (2019) 2156–2168.
- [15] AA. Kilbas, HM. Srivastava and J.J. Trujillo, *Theory and applications of fractional differential equations*, Amsterdam, Elsevier, 2006.
- [16] Y. Li and W. Jiang, *Existence and nonexistence of positive solutions for fractional three-point boundary value problems with a parameter*, *J. Funct. Space. Art. ID 9237856* (2019) 1–10.
- [17] Y. Li and A. Qi, *Positive solutions for multi-point boundary value problems of fractional differential equations with p -Laplacian*, *Math. Meth. Appl. Sci.* 39 (2016) 1425–1434.
- [18] X. Liu and M. Jia, *Multiple solutions for fractional differential equations with nonlinear boundary conditions*, *Comput. Math. Appl.* 59 (2010) 2880–2886.
- [19] X. Liu, M. Jia and W. Ge, *Multiple solutions of a p -Laplacian model involving a fractional derivative*, *Adv. Differ. Equ.* 2013 (2013) No. 126 1–12.
- [20] J. Podlubny, *Fractional differential equations*, New York, Academic Press, 1999.
- [21] H. Qu and X. Liu, *Existence of nonnegative solutions for a fractional m -point boundary value problem at resonance*, *Bound. Value Probl.* 127 (2013) 1–10.
- [22] HAH. Salem, *On the fractional order m -point boundary value problem in reflexive Banach spaces and weak topologies*, *J. Comput. Appl. Math.* 224 (2009) 565–572.
- [23] HAH. Salem and AMA. El-Sayed, *Weak solution for fractional order integral equations in reflexive Banach spaces*, *J. Comput. Appl. Math.* 224 (2009) 565–572.
- [24] K. Shah, S. Zeb and RA. Khan, *Multiplicity results of multi-point boundary value problem of nonlinear fractional differential equations*, *Appl. Math. Inf. Sci.* 12 (2018) 727–734.
- [25] GT. Wang, B. Ahmad and LH. Zhang, *Some existence results for impulsive nonlinear fractional differential equations with mixed boundary conditions*, *Comput. Math. Appl.* 62 (2011) 1389–1397.
- [26] GT. Wang, B. Ahmad and LH. Zhang, *On nonlocal integral boundary value problems for impulsive nonlinear differential equations of fractional order*, *Fixed Point Theory* 15 (2014) 265–284.
- [27] Y. Wang, *Existence and nonexistence of positive solutions for mixed fractional boundary value problem with parameter and p -Laplacian operator*, *J. Funct. Space. Art. ID 1462825* (2018) 1–6.
- [28] X. Xu, D. Jiang and C. Yuan, *Multiple positive solutions for the boundary value problem of a nonlinear fractional differential equation*, *Nonlinear Anal.* 71 (2009) 4676–4688.
- [29] X. Xu and H. Zhang, *Multiple positive solutions to singular positive and semipositone m -point boundary value problems of nonlinear fractional differential equations*, *Bound. Value Probl.* 34 (2018) 1–18.
- [30] A. Yang, J. Henderson and C. Nelms Jr., *Extremal points for a higher-order fractional boundary-value problem*, *Electron. J. Differential Equations* 2015 (2015) No. 161 1–12.
- [31] İ. Yaslan and M. Günendi, *Higher order multi-point fractional boundary value problems with integral boundary conditions*, *Int. J. Nonlinear Anal. Appl.* 9 (2018) 247–260.

- [32] X. Zhang and Q. Zhong, *Triple positive solutions for nonlocal fractional differential equations with singularities both on time and space variables*, Appl. Math. Lett. 80 (2018) 12–19.
- [33] C. Zhou, *Existence and uniqueness of positive solutions to higher-order nonlinear fractional differential equation with integral boundary conditions*, Electron. J. Differential Equations 2012 (2012) No. 234.
- [34] Y. Zou and G. He, *On the uniqueness of solutions for a class of fractional differential equations*, Appl. Math. Lett. 74 (2017) 68–73.