



A new approach for solution of Telegraph equation

Mohammad Zarebnia^{a,*}, Reza Parvaz^a

^aDepartment of Mathematics, University of Mohaghegh Ardabili, 56199-11367 Ardabil, Iran.

(Communicated by Madjid Eshaghi Gordji)

Abstract

In this paper, B-spline collocation method is developed for the solution of one-dimensional hyperbolic telegraph equation. The convergence of the method is proved. Also the method is applied on some test examples and the numerical results have been compared with the analytical solutions. The L_∞, L_2 and Root-Mean-Square errors (RMS) in the solutions show the efficiency of the method computationally.

Keywords: B-spline, Telegraph equation, Collocation method, Convergence
2010 MSC: Primary 65D07 ; Secondary 65N35 .

Telegraph equation is a kind of hyperbolic partial differential equation. This type of equations are used in the propagation of electromagnetic waves in the earth-ionosphere waveguide, ecological and cosmological phenomena modeling and mechanical wave [3]. Also, hyperbolic partial differential equations are commonly used in signal analysis for transmission and propagation of electrical signals [8] and also has applications in other fields [16, 14]. Telegraph equation can describe the voltage and current on an electrical transmission [1]. There are many types of telegraph equation as quantum and fractional telegraph equation [15, 2]. The quantum telegraph equation is used for the dynamics of probabilities and information in Quantum Mechanics [15]. The fractional telegraph equation is used for the anomalous diffusion processes observed in blood flow experiments [2]. In recent years, many different methods have been used to estimate the solution of the one-dimensional hyperbolic telegraph equation; see, for example, in [9], modified B-spline based on differential quadrature method is used, unconditionally stable difference schemes is used in [5], collocation method based on cubic b-spline is proposed in [18], and other method can be found in [6, 4, 10].

*Corresponding author

Email addresses: zarebnia@uma.ac.ir (Mohammad Zarebnia), rparvaz@uma.ac.ir (Reza Parvaz)

Consider the second-order linear hyperbolic partial differential equation in one-space dimension:

$$\frac{\partial^2 u}{\partial t^2} + 2\alpha \frac{\partial u}{\partial t} + \beta^2 u = \frac{\partial^2 u}{\partial x^2} + f(x, t), \quad a \leq x \leq b, \quad t \geq 0, \tag{0.1}$$

with the initial conditions

$$u(x, 0) = f_0(x), \tag{0.2}$$

$$\frac{\partial u}{\partial t}(x, 0) = f_1(x), \tag{0.3}$$

and boundary conditions

$$u(a, t) = g_0(t), \quad u(b, t) = g_1(t), \tag{0.4}$$

$$\frac{\partial u}{\partial x}(a, t) = g_2(t), \quad \frac{\partial u}{\partial x}(b, t) = g_3(t), \tag{0.5}$$

where α and β are constants.

The balance of this paper is organized as follows. In Section 2, the quintic B-spline collocation method for the numerical solution of the one-dimensional hyperbolic telegraph equation is described. In Section 3 we derive convergence of the B-spline collocation method. In Section 4, the results of numerical experiments are presented. A summary is given at the end of the paper in Section 5.

1. Quintic B-spline collocation method

The interval $[a, b]$ is partitioned into a mesh of uniform length $h := \frac{b-a}{N}$ by the knots $x_i, i = 0, 1, \dots, N$ such that $x_i := x_0 + ih$ and $a = x_0 < x_1 < x_2 < \dots < x_{N-1} < x_N = b$. To solve the equation (0.1) by collocation method with quintic B-splines as basis functions, we define the approximation $U^n(x)$ as following

$$U^n(x) = \sum_{i=-2}^{N+2} c_i^n B_i(x). \tag{1.1}$$

where $U^n(x)$ is a shape function that approximates $u(x, t_n)$ for the time level $t_n = nk$ where k is a time step size. For each time level t_n , the set $\{c_{-2}^n, c_{-1}^n, \dots, c_{N+1}^n, c_{N+2}^n\}$ are unknown real coefficients, which are to be found, and the $B_i(x)$ are the quintic B-spline functions defined by [17, 13]

$$B_i(x) = \frac{1}{h^5} \begin{cases} (x - x_{i-3})^5, & x \in [x_{i-3}, x_{i-2}), \\ (x - x_{i-3})^5 - 6(x - x_{i-2})^5, & x \in [x_{i-2}, x_{i-1}), \\ (x - x_{i-3})^5 - 6(x - x_{i-2})^5 + 15(x - x_{i-1})^5, & x \in [x_{i-1}, x_i), \\ (x_{i+3} - x)^5 - 6(x_{i+2} - x)^5 + 15(x_{i+1} - x)^5, & x \in [x_i, x_{i+1}), \\ (x_{i+3} - x)^5 - 6(x_{i+2} - x)^5, & x \in [x_{i+1}, x_{i+2}), \\ (x_{i+3} - x)^5, & x \in [x_{i+2}, x_{i+3}), \end{cases} \tag{1.2}$$

where $B_{-2}, B_{-1}, B_0, B_1, \dots, B_{N+1}, B_{N+2}$ form a basis over the region $a \leq x \leq b$. The values of $B_i(x)$ and its derivatives may be tabulated as in Table 1. Using approximate function (1.1) and Table 1, we have

$$u(x_i, t_n) \approx U_i^n = c_{i-2}^n + 26c_{i-1}^n + 66c_i^n + 26c_{i+1}^n + c_{i+2}^n, \tag{1.3}$$

$$\frac{\partial u}{\partial x}(x_i, t_n) \approx (U')_i^n = \frac{1}{h}(-5c_{i-2}^n - 50c_{i-1}^n + 50c_{i+1}^n + 5c_{i+2}^n), \tag{1.4}$$

$$\frac{\partial^2 u}{\partial x^2}(x_i, t_n) \approx (U'')_i^n = \frac{1}{h^2}(20c_{i-2}^n + 40c_{i-1}^n - 120c_i^n + 40c_{i+1}^n + 20c_{i+2}^n). \tag{1.5}$$

Table 1: B_i, B'_i and B''_i at the node points.

x	x_{i-3}	x_{i-2}	x_{i-1}	x_i	x_{i+1}	x_{i+2}	x_{i+3}
$B_i(x)$	0	1	26	66	26	1	0
$hB'_i(x)$	0	5	50	0	-50	-5	0
$h^2B''_i(x)$	0	20	40	-120	40	20	0

To apply the proposed method, discretizing the time derivative in the usual finite difference way, with using following finite difference formulae [7], we can write:

$$\left(\frac{\partial^2 u}{\partial t^2}\right)_i^n \approx \frac{u_i^{n+1} - 2u_i^n + u_i^{n-1}}{\Gamma(k)^2}, \tag{1.6}$$

$$\left(\frac{\partial u}{\partial t}\right)_i^n \approx \frac{u_i^{n+1} - u_i^{n-1}}{2\Gamma(k)}, \tag{1.7}$$

$$\left(\frac{\partial^2 u}{\partial x^2}\right)_i^n \approx \frac{\left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n-1} + \left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n+1}}{2}, \tag{1.8}$$

where k is a time step size, $\left(\frac{\partial^2 u}{\partial x^2}\right)_i^n := \frac{\partial^2 u}{\partial x^2}(x_i, t_n)$, $u_i^n := u(x_i, t_n)$ and $\Gamma(k)$ is a selected function of k satisfying the following equation

$$\Gamma(k)^2 = (k)^2(1 + \mathcal{O}(k)^j), \quad j = 0, 1, \dots \tag{1.9}$$

In the numerical computations, we applied the following two possible choices for $\Gamma(k)$ to improve the accuracy: k and $2 \sin(\frac{k}{2})$. Hence (0.1) can be written as:

$$\frac{u_i^{n+1} - 2u_i^n + u_i^{n-1}}{\Gamma(k)^2} + 2\alpha \frac{u_i^{n+1} - u_i^{n-1}}{2\Gamma(k)} + \beta^2 u_i^n = \frac{\left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n-1} + \left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n+1}}{2} + f(x_i, t_n). \tag{1.10}$$

Rearranging the terms and simplifying we get

$$vu_i^{n+1} + w\left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n+1} = \Phi^n(x_i), \tag{1.11}$$

where

$$\Phi^n(x_i) := \left(2 - (\beta\Gamma(k))^2\right)u_i^n + (\alpha\Gamma(k) - 1)u_i^{n-1} + \frac{\Gamma(k)^2}{2}\left(\frac{\partial^2 u}{\partial x^2}\right)_i^{n-1} + \Gamma(k)^2 f(x_i, t_n), \tag{1.12}$$

$$v := 1 + \alpha\Gamma(k), \tag{1.13}$$

$$w := -\frac{\Gamma(k)^2}{2}. \tag{1.14}$$

Substituting the approximate solution U for u and putting the values (1.3) and (1.5) in (1.11) yields the following difference equation with the variables $c_i, i = -2, \dots, N + 2$.

$$\begin{aligned} &vU_i^{n+1} + w(U'')_i^{n+1} = \\ &(v + 20\frac{w}{h^2})c_{i+2}^{n+1} + (26v + 40\frac{w}{h^2})c_{i+1}^{n+1} + (66v - 120\frac{w}{h^2})c_i^{n+1} \\ &+ (26v + 40\frac{w}{h^2})c_{i-1}^{n+1} + (v + 20\frac{w}{h^2})c_{i-2}^{n+1} = \Psi_i^n, \quad i = 0, \dots, N, \end{aligned} \tag{1.15}$$

where

$$\Psi_i^n := \left(2 - (\beta\Gamma(k))^2\right)U_i^n + (\alpha\Gamma(k) - 1)U_i^{n-1} + \frac{\Gamma(k)^2}{2}(U''_i)^{n-1} + \Gamma(k)^2 f(x_i, t_n). \tag{1.16}$$

The system (1.15) consists of $(N+1)$ linear equations in $(N+5)$ unknowns $\tilde{C} := \{c_{-2}, c_{-1}, \dots, c_{N+1}, c_{N+2}\}$. To obtain a unique solution for \tilde{C} we must use the boundary conditions. From the boundary conditions we can write

$$c_{-2}^{n+1} + 26c_{-1}^{n+1} + 66c_0^{n+1} + 26c_1^{n+1} + c_2^{n+1} = g_0(t_{n+1}), \tag{1.17}$$

$$c_{N-2}^{n+1} + 26c_{N-1}^{n+1} + 66c_N^{n+1} + 26c_{N+1}^{n+1} + c_{N+2}^{n+1} = g_1(t_{n+1}), \tag{1.18}$$

and

$$\frac{1}{h}(-5c_{-2}^{n+1} - 50c_{-1}^{n+1} + 50c_1^{n+1} + 5c_2^{n+1}) = g_2(t_{n+1}), \tag{1.19}$$

$$\frac{1}{h}(-5c_{N-2}^{n+1} - 50c_{N-1}^{n+1} + 50c_{N+1}^{n+1} + 5c_{N+2}^{n+1}) = g_3(t_{n+1}). \tag{1.20}$$

By using (1.17)-(1.20), we obtain

$$c_{-1}^{n+1} = -\frac{33}{8}c_0^{n+1} - \frac{9}{4}c_1^{n+1} - \frac{1}{8}c_2^{n+1} + \frac{hg_2(t_{n+1})}{80} + \frac{g_0(t_{n+1})}{16}, \tag{1.21}$$

$$c_{-2}^{n+1} = \frac{165}{4}c_0^{n+1} + \frac{65}{2}c_1^{n+1} + \frac{9}{4}c_2^{n+1} - \frac{13hg_2(t_{n+1})}{40} - \frac{5g_0(t_{n+1})}{8}, \tag{1.22}$$

$$c_{N+1}^{n+1} = -\frac{33}{8}c_N^{n+1} - \frac{9}{4}c_{N-1}^{n+1} - \frac{1}{8}c_{N-2}^{n+1} - \frac{hg_3(t_{n+1})}{80} + \frac{g_1(t_{n+1})}{16}, \tag{1.23}$$

$$c_{N+2}^{n+1} = \frac{165}{4}c_N^{n+1} + \frac{65}{2}c_{N-1}^{n+1} + \frac{9}{4}c_{N-2}^{n+1} + \frac{13hg_3(t_{n+1})}{40} - \frac{5g_1(t_{n+1})}{8}. \tag{1.24}$$

Hence we have the following system consists of $(N + 1)$ linear equations in $(N + 1)$ unknowns $\{c_0, c_1, \dots, c_{N-1}, c_N\}$. The B-spline method in matrix form can be written as follows :

$$AC = Q, \tag{1.25}$$

where

$$\mathbf{A} := \begin{pmatrix} \acute{c} - \frac{33\acute{b}}{8} + \frac{165\acute{a}}{4} & \frac{65\acute{a}}{2} - \frac{5\acute{b}}{4} & \frac{13\acute{a}}{4} - \frac{\acute{b}}{8} & 0 & \dots & 0 & \dots & 0 \\ \acute{b} - \frac{33\acute{a}}{8} & \acute{c} - \frac{9\acute{a}}{4} & \acute{b} - \frac{\acute{a}}{8} & a & 0 & \dots & \dots & 0 \\ \acute{a} & \acute{b} & \acute{c} & \acute{b} & \acute{a} & \dots & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & \dots & \acute{a} & \acute{b} & \acute{c} & \acute{b} & \acute{a} & \dots \\ 0 & \dots & 0 & \acute{a} & \acute{b} - \frac{33\acute{a}}{8} & \acute{c} - \frac{9\acute{a}}{4} & \acute{b} - \frac{\acute{a}}{8} & \dots \\ 0 & \dots & 0 & 0 & \acute{c} - \frac{33\acute{b}}{8} + \frac{165\acute{a}}{4} & \frac{65\acute{a}}{2} - \frac{5\acute{b}}{4} & \frac{13\acute{a}}{4} - \frac{\acute{b}}{8} & \dots \end{pmatrix}, \tag{1.26}$$

$$C := \left(c_0^{n+1}, c_1^{n+1}, \dots, c_{N-1}^{n+1}, c_N^{n+1}\right)^T, \tag{1.27}$$

and

$$\mathbf{Q} := \begin{pmatrix} \Psi^n(x_0) + (-\frac{bh}{80} + \frac{13ah}{40})g_2(t_{n+1}) + (-\frac{b}{16} + \frac{5a}{8})g_0(t_{n+1}) \\ \Psi^n(x_1) - \frac{ha}{80}g_2(t_{n+1}) - \frac{a}{16}g_0(t_{n+1}) \\ \Psi^n(x_2) \\ \vdots \\ \Psi^n(x_{N-2}) \\ \Psi^n(x_{N-1}) + \frac{ha}{80}g_3(t_{n+1}) - \frac{a}{16}g_1(t_{n+1}) \\ \Psi^n(x_N) + (\frac{bh}{80} - \frac{13ah}{40})g_3(t_{n+1}) + (-\frac{b}{16} + \frac{5a}{8})g_1(t_{n+1}) \end{pmatrix}, \tag{1.28}$$

with

$$\acute{a} := v + 20\frac{w}{h^2}, \tag{1.29}$$

$$\acute{b} := 26v + 40\frac{w}{h^2}, \tag{1.30}$$

$$\acute{c} := 66v - 120\frac{w}{h^2}. \tag{1.31}$$

The computer algebra system *Mathematica-9* is used for solving the system (1.25). To start any computation, it is necessary to know the value of u at the nodal points of first time level, that is, at $t = k$. A Taylor series expansion at $t = k$ may be written as

$$u(x, k) = f_0(x) + kf_1(x) + \frac{k^2}{2}(f(x, 0) - \beta^2 f_0(x) - 2\alpha f_1(x) + \frac{\partial f_0(x)}{\partial x}) + \mathcal{O}(k^3). \tag{1.32}$$

2. Convergence analysis

Theorem 2.1. *Suppose that $u(x, t)$ be the exact solution of (0.1) and $u(x, t) \in C^5[a, b]$ also $|\frac{\partial^5 u(x, t)}{\partial x^5}| \leq L$ and $U(x, t)$ be the numerical approximation by our methods, then we can write*

$$\| u(x, t) - U(x, t) \|_\infty = \mathcal{O}(h^2 + k). \tag{2.1}$$

Before we prove, we recall following theorem and lemma.

Theorem 2.2. *Suppose that $f(x) \in C^5[a, b]$. Then for the unique quintic spline $S(x)$ associated with f , we have*

$$\| f^{(j)} - S^{(j)} \|_\infty \leq K_j \omega_5(h) h^{4-j}, \quad j = 0, 1, 2, 3. \tag{2.2}$$

where $\omega_5(h)$ denotes the modulus of continuity of $f^{(5)}$ and the coefficients λ_j are independent of f and h .

Proof . For the proof see [12]. \square

Remark 2.3. *By using Theorem 2.2 and definition of the modulus of continuity, we can say that if $|f^{(5)}(x)| \leq L$, we can write (2.2) as*

$$\| f^{(j)} - S^{(j)} \|_\infty \leq \lambda_j L h^{4-j}, \quad j = 0, 1, 2, 3. \tag{2.3}$$

Lemma 2.4. *For the B-splines $\{B_{-2}, \dots, B_{N+2}\}$ we have the following inequality:*

$$\left| \sum_{i=-2}^{N+2} B_i(x) \right| \leq 186, \quad (a \leq x \leq b). \tag{2.4}$$

Proof . From the real analysis we have $|\sum_{i=-2}^{N+2} B_i(x)| \leq \sum_{i=-2}^{N+2} |B_i(x)|$. If $x = x_i, i = 1, \dots, N$, then, we have

$$|\sum_{i=-2}^{N+2} B_i(x)| = 120 \leq 186, \tag{2.5}$$

and if $x_{i-1} \leq x \leq x_i$, then, we can write

$$\begin{aligned} |\sum_{i=-2}^{N+2} B_i(x)| &\leq |B_{i-3}(x)| + |B_{i-2}(x)| + |B_{i-1}(x)| + |B_i(x)| \\ &+ |B_{i+1}(x)| + |B_{i+2}(x)| \leq 1 + 26 + 66 + 66 + 26 + 1 \leq 186 \end{aligned}$$

□ Now we prove theorem 2.1.

Proof . Suppose that $\varepsilon_i = u(t_i) - U^i$ be the local truncation error for (1.10) at the i th. By using the truncation error, we can write

$$|\varepsilon_i| \leq \varrho_i k^2, \quad i \geq 2. \tag{2.6}$$

In addition we have $|\varepsilon_1| \leq \varrho_1 k^3$. To continue we assume that e_{n+1} be the global error in time discretizing process and $\varrho = \max\{\varrho_1, \dots, \varrho_n\}$. We can write the following global error estimate at $n + 1$ level

$$e_{n+1} = \sum_{i=1}^n \varepsilon_i, \quad (k \leq T/n), \tag{2.7}$$

with the help of (2.6)-(2.7), we can write

$$|e_{n+1}| = |\sum_{i=1}^n \varepsilon_i| \leq n\varrho k^2 \leq n\varrho \frac{T}{n} k = \rho k, \tag{2.8}$$

where $\rho = \varrho T$.

Now at the $(n + 1)$ th time step we assume that $u(x)$ be the exact solution of (1.11) and $U(x) = \sum_{i=-2}^{N+2} c_i B_i(x)$ be the B-spline approximation to $u(x)$. Also we assume that $S^*(x) = \sum_{i=-2}^{N+2} c_i^* B_i(x)$ be the unique spline interpolate to the exact solution. In order to derive a bound for $\|u(x) - U(x)\|_\infty$, we need to estimate the $\|u(x) - S^*(x)\|_\infty$ and $\|S^*(x) - U(x)\|_\infty$. Now we substituting $S^*(x)$ in (1.11) the we get the following result

$$AC^* = Q^*, \tag{2.9}$$

With considering (1.25) and (2.9), we get

$$A(C^* - C) = (Q^* - Q). \tag{2.10}$$

From (1.15), we can write

$$|\Psi_i^* - \Psi_i| \leq v |S^*(x_i) - U(x_i)| + w |S^{*''}(x_i) - U''(x_i)|. \tag{2.11}$$

By using (2.11) and Theorem 2.2, we can write

$$\|Q^* - Q\|_\infty \leq M_1 h^2, \tag{2.12}$$

where $M_1 = v\lambda_0 L h^2 + w\lambda_2 L$. In this step from (2.10), we can write

$$(C^* - C) = A^{-1}(Q^* - Q). \tag{2.13}$$

By taking the infinity norm from (2.13) and applying (2.12), we get

$$\|C^* - C\|_\infty \leq \|A^{-1}\|_\infty \|Q^* - Q\|_\infty \leq M_1 h^2 \|A^{-1}\|_\infty. \quad (2.14)$$

By using the theory of matrices, we can write

$$\sum_{i=1}^{N+1} a_{ki}^{-1} \eta_i = 1, \quad (2.15)$$

where a_{ki}^{-1} are the elements of A^{-1} and η_i ($1 \leq i \leq N+1$) is the summation of the i th row of the matrix A . As a result we can write

$$\|A^{-1}\|_\infty = \sum_{i=1}^{N+1} |a_{ki}^{-1}| \leq \frac{1}{\min_{1 \leq i \leq N} \eta_i} \leq \frac{1}{\Lambda}, \quad (2.16)$$

where Λ is constant. Following result is obtained by substituting (2.16) into (2.14), we get

$$\|C^* - C\|_\infty \leq \frac{M_1 h^2}{\Lambda} \leq M_2 h^2, \quad (2.17)$$

where $M_2 = \frac{M_1}{\Lambda}$ is constant. Considering the B-spline collocation approximation and the computed spline approximation, we can write:

$$S^*(x) - U(x) = \sum_{i=-2}^{N+2} (c_i^* - c_i) B_i(x), \quad (2.18)$$

taking norm from (2.18) and by using (2.17) and lemma 2.4, we obtain

$$\|S^*(x) - U(x)\|_\infty = \left\| \sum_{i=-2}^{N+2} (c_i^* - c_i) B_i(x) \right\|_\infty \leq \left| \sum_{i=-2}^{N+2} B_i(x) \right| \|C^* - C\|_\infty \leq 186 M_2 h^2. \quad (2.19)$$

Also from Theorem 2.2 we can write

$$\|u - S^*(x)\|_\infty \leq \lambda_0 L h^4, \quad (2.20)$$

and therefore with helping (2.19) and (2.20) we get

$$\|u - U(x)\|_\infty \leq \varpi h^2, \quad (2.21)$$

where $\varpi = \lambda_0 L h^2 + 186 M_2$. \square

3. Numerical examples

In order to illustrate the performance of the quintic B-spline collocation method in solving the One-dimensional hyperbolic telegraph equation and justify the accuracy and efficiency of the present method, we consider the following examples. To show the efficiency of the present method

for our problem in comparison with the exact solution, we report the RMS error, L_∞ and L_2 using formulae

$$RMS = \frac{(\sum_{i=1}^N |u(x_i, t) - U_n(x_i)|^2)^{\frac{1}{2}}}{N^{\frac{1}{2}}},$$

$$L_\infty = \max_i |U_n(x_i) - u(x_i, t)|, L_2 = h \left| \sum_{i=1}^N (u(x_i, t) - U_n(x_i))^2 \right|,$$

where $U(x, t)$ denotes numerical solution and $u(x, t)$ denotes analytical solution.

Example 1. Consider the hyperbolic telegraph equation (0.1) with $\alpha = \pi, \beta = \pi$, in the interval $[0, 1]$. In this case we have $f(x, t) = \pi^2 \sin(\pi x)(\sin(\pi t) + 2 \cos(\pi t))$. The analytical solution given by $u(x, t) = \sin(\pi t) \sin(\pi x)$. The boundary conditions and the initial conditions are taken from the exact solution. Table 2 shows the absolute error between the analytical solution and the numerical solution at different points for $t = 0.5$. Table 3 shows the L_2 errors at different partitions. The graph of the solution is given in Figure 1 . Also, Figure 2 shows that the solution obtained by our method is close to the exact solution

Table 2: A comparison of absolute errors of Example 1 at different points with $h = 1/100, k = 1/200$.

Method	present method		method in [11]
x	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	$\eta = \frac{1}{60}, \gamma = \frac{1}{2}$ grid
0.2	3.23677×10^{-5}	3.26277×10^{-5}	$5.858898718658607 \times 10^{-1}$
0.4	5.32737×10^{-5}	5.32737×10^{-5}	$9.479897263432836 \times 10^{-1}$
0.6	5.32737×10^{-5}	5.32737×10^{-5}	$9.479897263432840 \times 10^{-1}$
0.8	5.32737×10^{-5}	3.26277×10^{-5}	$5.858898718658610 \times 10^{-1}$

Table 3: A comparison of L_2 errors of Example 1 at different partitions.

Partitions	N=100, k=0.01		N=400, k=0.001	
Time	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$
0.5	1.5799×10^{-5}	1.58168×10^{-5}	1.57942×10^{-7}	7.91938×10^{-8}
1	6.94858×10^{-6}	6.61076×10^{-6}	6.94312×10^{-8}	3.33013×10^{-8}
1.5	1.50368×10^{-5}	1.51490×10^{-5}	1.50334×10^{-7}	7.57557×10^{-8}
2	7.25141×10^{-6}	6.89868×10^{-6}	7.24547×10^{-8}	3.4743×10^{-8}

Example 2. We consider the hyperbolic telegraph equation (0.1) with $f(x, t) = (3 - 4\alpha + \beta^2) \exp(-2t) \sinh(x)$ and the analytical solution $u(x, t) = \exp(-2t) \sinh(x)$, in the interval $[0, 1]$. The boundary conditions and the initial conditions are taken from the exact solution. Tables 4 and 5 give a comparison between the L_∞ errors found by our method and the method in [6]. Also Table 5 shows RMS and L_2 errors. Figure 3, shows absolute error for different values of time with $N = 200, k = 0.001$.




Figure 1: The graph of the solution for Example 1 with $\Gamma(k) = k, N = 200$ and $k=0.01$.




Figure 2: Comparisons between numerical and analytical solutions with $\Gamma(k) = k$ (left) and $\Gamma(k) = 2\sin(k/2)$ (right) for Example 1 at different times with $N = 200, k = 0.01$.

Table 4: A comparison of L_∞ errors of Example 2 for $\alpha = 20, \beta = 10$ at different time and $N = 21, k = 0.01$.

Method	present method		method in [6]	
	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	Uniform Grid	Nonuniform grid
0.5	2.73131×10^{-6}	2.59359×10^{-6}	2.25640×10^{-4}	2.22066×10^{-4}
1	1.5714×10^{-6}	1.48998×10^{-6}	5.75205×10^{-3}	1.41294×10^{-4}
1.5	7.01914×10^{-7}	6.65254×10^{-7}	2.43143×10^{-2}	6.93111×10^{-5}
2	2.8573×10^{-7}	2.70754×10^{-7}	—————	2.27829×10^{-5}

Table 5: A comparison of L_∞ errors of Example 2 for $\alpha = 20, \beta = 10$ at different time and $N = 21, k = 0.0001$.

Method	present method		method in [6]	
	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	Uniform Grid	Nonuniform grid
0.5	1.9517×10^{-9}	1.92002×10^{-9}	2.23874×10^{-6}	2.21693×10^{-6}
1	1.08081×10^{-9}	1.06278×10^{-9}	1.72404×10^{-5}	1.22688×10^{-6}
1.5	4.75261×10^{-10}	4.67199×10^{-10}	4.14630×10^{-3}	6.73102×10^{-7}
2	1.91059×10^{-10}	1.87808×10^{-10}	—————	2.66660×10^{-7}

Table 6: L_2 and RMS errors of Example 2 for $\alpha = 20, \beta = 10$ at different time and $N = 21, k = 0.01$.

Γ	$2\sin(\frac{k}{2})$		k	
	L_2	RMS	L_2	RMS
0.2	2.36511×10^{-6}	1.08383×10^{-5}	2.25651×10^{-6}	1.03406×10^{-5}
0.4	2.82003×10^{-6}	1.2923×10^{-5}	2.67987×10^{-6}	1.22807×10^{-5}
0.8	2.0589×10^{-6}	9.43506×10^{-6}	1.95292×10^{-6}	8.94939×10^{-6}
1	1.5714×10^{-6}	7.20105×10^{-6}	1.48999×10^{-6}	6.82797×10^{-6}

Example 3. In this example we Consider the hyperbolic telegraph equation (0.1) with $\alpha = 10, \beta = 5$ in $x \in [0, 1]$ and $f(x, t) = -2\alpha \sin(t) \sin(x) + \beta^2 \cos(t) \cos(x)$. The exact solution for this case is $u(x, t) = \cos(t) \sin(x)$. The boundary conditions and the initial conditions are taken from the exact solution. In order to compare the solutions with [6], we have taken $k = 0.001$ and $N = 21$. Table 7 gives a comparison between the L_∞ error found by our method and by method in [6]. Table 8 shows L_2 in different partitions. Table 9 shows RMS and L_2 errors. Figure 4, shows absolute error for different values of time with $N = 50, k = 0.001$.




Figure 3: Absolute errors with $\Gamma(k) = k$ (left) and $\Gamma(k) = 2\sin(k/2)$ (right) for Example 2 with $k = 0.001$, $N = 200$ and $\alpha = 20, \beta = 10$.

Table 7: A comparison of L_∞ errors for Example 3 at different time and $N = 21, k = 0.001$.

Method	present method		method in [6]	
Time	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	Uniform Grid	Nonuniform grid
0.5	7.59175×10^{-9}	5.35357×10^{-9}	1.67216×10^{-5}	1.28551×10^{-5}
1	3.5984×10^{-9}	2.39252×10^{-8}	4.71302×10^{-4}	3.20793×10^{-5}
1.5	1.61922×10^{-8}	4.2016×10^{-8}	8.62796×10^{-4}	5.71454×10^{-5}
2	2.49884×10^{-8}	5.20784×10^{-8}	—————	6.82827×10^{-5}

Table 8: A comparison of L_2 errors of Example 3 at different partitions.

Partitions	N=200,k=0.01		N=400,k=0.001	
	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$	$\Gamma : 2\sin(\frac{k}{2})$	$\Gamma : k$
0.5	3.69769×10^{-8}	2.46374×10^{-8}	2.65293×10^{-10}	1.73258×10^{-10}
1	1.40829×10^{-8}	1.16929×10^{-7}	9.95706×10^{-11}	8.27160×10^{-10}
1.5	8.1959×10^{-8}	2.09527×10^{-7}	5.80334×10^{-10}	1.48296×10^{-10}
2	1.28423×10^{-7}	2.61864×10^{-7}	9.09458×10^{-10}	1.85373×10^{-9}

Table 9: L_2 and RMS errors of Example 3 at different time and $N = 80, k = 0.05$.

Time	$2\sin(\frac{k}{2})$		k	
	L_2	RMS	L_2	RMS
0.5	1.37362×10^{-6}	1.2286×10^{-5}	9.94623×10^{-7}	8.89618×10^{-6}
1	5.5496×10^{-7}	4.96372×10^{-6}	4.61873×10^{-6}	4.13112×10^{-5}
1.5	3.22627×10^{-6}	2.88566×10^{-5}	8.26127×10^{-6}	7.38911×10^{-5}
2	5.05609×10^{-6}	4.52231×10^{-5}	1.03189×10^{-5}	9.22947×10^{-5}

4. Conclusion

The quintic B-spline collocation method is used to solve the one-dimensional hyperbolic telegraph equation with initial and boundary conditions. The convergence analysis of the method is shown. The numerical solutions are compared with the exact solution by finding the RMS, L_2 and L_∞ errors. The numerical results given in the previous section demonstrate the good accuracy of the scheme proposed in this research.




Figure 4: Absolute errors with $\Gamma(k) = k$ (left) and $\Gamma(k) = 2\sin(k/2)$ (right) for Example 3 with $k = 0.001$, $N = 50$.

References

- [1] A. Atangana, *On the stability and convergence of the time-fractional variable order telegraph equation*, J. Comput. Phys. 293 (2015) 104–114.
- [2] R.C. Cascaval, E. Eckstein, C.L. Frota, J.A. Goldstein, *Fractional telegraph equations*, J. Math. Ana. Appl. 276(1) (2002) 145–159.
- [3] V. Devi, R.K. Maurya, S. Singh, V. K.Singh, *Lagrange's operational approach for the approximate solution of two-dimensional hyperbolic telegraph equation subject to Dirichlet boundary conditions*, Appl. Math. Comput. 367 (2020) 124717.
- [4] H.F. Ding, Y.X. Zhangb, J.X Caoa, J.H Tianb, *A class of difference scheme for solving telegraph equation by new non-polynomial spline methods*, Appl. Math. Comput. 278 (2012) 4671–4683.
- [5] F. Gao, C. Chi, *Unconditionally stable difference schemes for a one-space-dimensional linear hyperbolic equation*, Appl. Math. Comput. 187 (2007) 1272–1276.
- [6] R. Jiware, S. Pandit, R.C. Mittal, *A differential quadrature algorithm for solution of the second order one dimensional hyperbolic telegraph equation*, IJNS 13 (2012) 259–266.
- [7] S.A. Khuri, A. Sayfy, *A spline collocation approach for the numerical solution of a generalized nonlinear Klein Gordon equation*, Appl. Math. Comput. 216 (2010) 1047–1056.
- [8] A. C. Metaxas and R. J. Meredith, *Industrial microwave*, Heating, Peter Peregrinus, London, 1993.
- [9] R.C. Mittal, R. Bhatia, *A numerical study of two dimensional hyperbolic telegraph equation by modified B-spline differential quadrature method*, Appl. Math. Comput. 244 (2014) 976–997.
- [10] R.K. Mohanty, M.K. Jain, K. George, *On the use of high order difference methods for the system of one space second order nonlinear hyperbolic equations with variable coefficients*, J. Comput. Appl. Math. 72 (1996) 421–431.
- [11] R.K. Mohanty, *An unconditionally stable finite difference formula for a linear second order one space dimensional hyperbolic equation with variable coefficients*, Appl. Math. Comput. 165 (2005) 229–236.
- [12] G. Pittaluga, L. Sacripante, *Quintic spline interpolation on uniform meshes*, Acta. Math. Hung. 72 (1996) 4167–175.
- [13] P.M. Prenter, *Spline and Variational Methods*, Wiley, New York, 1975.
- [14] G. Roussy, J. A. Percy, *Foundations and Industrial Applications of Microwaves and Radio Frequency Fields*, John Wiley, New York, 1995.
- [15] P. Sancho, *Relativistic Extension of the Quantum Telegraph Equation*, Open. Syst. Inf. Dyn. 7(2) (2000) 157–164.
- [16] J. N. Sharma, K. Singh, J. N. Sharma, *Partial differential equations for engineers and scientists*, second edition, Alpha Science International, 2009.
- [17] J. Stoer, R. Bulirsch, *Introduction to numerical analysis*, third edition, Springer-Verlag, 2002.
- [18] M. Zarebnia., R. Parvaz, *Cubic B-spline collocation method for numerical solution of the one-dimensional hyperbolic telegraph equation*, JARSC 4(4) (2012) 46–60.