



Existence Theory for Higher-Order Nonlinear Ordinary Differential Equations with Nonlocal Stieltjes Boundary Conditions

Bashir Ahmad^{a,*}, Ahmed Alsaedi^a, Nada Al-Malki^a

^aNonlinear Analysis and Applied Mathematics (NAAM) Research Group, Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

(Communicated by Madjid Eshaghi Gordji)

Abstract

In this paper, we develop the existence theory for some boundary value problems of nonlinear n th-order ordinary differential equations supplemented with nonlocal Stieltjes boundary conditions. Our results are based on some standard theorems of fixed point theory and are well illustrated with the aid of examples.

Keywords: higher-order differential equations; Stieltjes; nonlocal boundary conditions; fixed point.

2010 MSC: Primary 34B10; Secondary 34B15, 34A60.

1. Introduction

The study of nonlinear boundary value problems of differential equations is of central importance in mathematics in view of their extensive applications in applied sciences such as fluid mechanics, geophysics, mathematical physics, etc. The recent development of the subject includes several kinds of nonlocal and integral boundary conditions. The nonlocal conditions take care of peculiarities of physical, chemical or other processes occurring at some intermediate positions of the given domain while the integral conditions provide an alternative for the assumption of ‘circular cross-section’ throughout the vessels in the study of fluid flow problems. For examples and details of nonlocal

*Corresponding author

Email addresses: bashirahmad_qau@yahoo.com (Bashir Ahmad), aalsaedi@hotmail.com (Ahmed Alsaedi), mass_nana@hotmail.com (Nada Al-Malki)

problems, we refer the reader to a series of papers ([1]-[12]), whereas the works involving integral boundary conditions can be found in ([13]-[21]) and the references cited therein.

In this paper, we consider nonlocal multi-point and strip type Riemann-Stieltjes integral boundary conditions and establish the existence theory for boundary value problems of n th-order ordinary differential equations supplemented with these conditions. The concept of Stieltjes conditions provides a unified approach for dealing with a variety of boundary conditions such as multipoint and integral boundary conditions. For details on Riemann-Stieltjes integral conditions, we refer the reviews by Whyburn [22] and Conti [23]. Recently, Webb [24, 25] and Webb and Infante [26] studied the higher order problems with Stieltjes integral boundary conditions via fixed point index. Some more works on Riemann-Stieltjes integral conditions can be found in a series of papers [27, 28, 29, 30] and the references cited therein.

The rest of the paper is organized as follows. In Section 2, we formulate and solve a model problem. The existence results for this problem are obtained via contraction mapping principle and Schauder's fixed point and are explained with the help of some examples. In Section 3, we discuss some more problems involving Stieltjes conditions.

2. Model Problem

We consider a nonlocal Stieltjes type boundary value problem involving an n th-order nonlinear ordinary differential equation given by

$$\begin{cases} u^{(n)}(t) = f(t, u(t)), & t \in [0, 1], \\ u(0) = \delta u(\xi), \quad u'(0) = 0, \quad u''(0) = 0, \dots, \quad u^{(n-2)}(0) = 0, \\ \alpha u(1) + \beta u'(1) = \int_0^1 u(s) d\mu(s), \quad 0 < \xi < 1, \end{cases} \quad (2.1)$$

where $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a given continuous function, μ is function of bounded variation and $\alpha, \beta, \delta, \xi$ are real constants satisfying the relation:

$$\Lambda = \left(\alpha - \int_0^1 d\mu(s) \right) (\delta \xi^{n-1}) + \left(\alpha + \beta(n-1) - \int_0^1 s^{n-1} d\mu(s) \right) (1 - \delta) \neq 0. \quad (2.2)$$

In the following lemma, we solve a linear variant of problem (2.1).

2.1. Basic result

Lemma 2.1. *For any $y \in C([0, 1], \mathbb{R})$, the linear differential equation*

$$u^{(n)}(t) = y(t), \quad t \in [0, 1], \quad (2.3)$$

supplemented with boundary conditions of problem (2.1) is equivalent to the integral equation:

$$\begin{aligned} u(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} y(s) ds + \sigma_1(t) \left[\int_0^1 \left(\int_0^s \frac{(s-p)^{n-1}}{(n-1)!} y(p) dp \right) d\mu(s) \right. \\ &\quad \left. - \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} y(s) ds \right] \\ &\quad + \sigma_2(t) \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} y(s) ds, \end{aligned} \quad (2.4)$$

where

$$\sigma_1(t) = \frac{1}{\Lambda} [\delta \xi^{n-1} + t^{n-1}(1 - \delta)], \tag{2.5}$$

$$\sigma_2(t) = \frac{\delta}{\Lambda} \left[\alpha + \beta(n - 1) - \int_0^1 s^{n-1} d\mu(s) - t^{n-1} \left(\alpha - \int_0^1 d\mu(s) \right) \right], \tag{2.6}$$

and Λ is given by (2.2).

Proof . It is well known that the solution of the differential equation (2.3) can be written as

$$u(t) = \int_0^t \frac{(t - s)^{n-1}}{(n - 1)!} y(s) ds + c_0 + c_1 t + c_2 t^2 + \dots + c_{n-2} t^{n-2} + c_{n-1} t^{n-1}, \tag{2.7}$$

where c_i ($i = 0, 1, \dots, n - 1$) are arbitrary real constants. Using the boundary conditions: $u'(0) = 0$, $u''(0) = 0, \dots, u^{(n-2)}(0) = 0$ in (2.7), we find that $c_1 = c_2 = \dots = c_{n-2} = 0$, and consequently, (2.7) takes the form:

$$u(t) = \int_0^t \frac{(t - s)^{n-1}}{(n - 1)!} y(s) ds + c_0 + c_{n-1} t^{n-1}. \tag{2.8}$$

Now using the remaining boundary conditions:

$$u(0) = \delta u(\xi), \quad \alpha u(1) + \beta u'(1) = \int_0^1 u(s) d\mu(s),$$

in (2.8), we get

$$A_1 c_0 - A_2 c_{n-1} = A_3 \quad \text{and} \quad E_1 c_0 + E_2 c_{n-1} = E_3, \tag{2.9}$$

where

$$\begin{aligned} A_1 &= 1 - \delta, \quad A_2 = \delta \xi^{n-1}, \quad A_3 = \delta \int_0^\xi \frac{(\xi - s)^{n-1}}{(n - 1)!} y(s) ds, \\ E_1 &= \alpha - \int_0^1 d\mu(s), \quad E_2 = \alpha + (n - 1)\beta - \int_0^1 s^{n-1} d\mu(s), \quad E_3 = B_3 - B_2 - B_1, \\ B_1 &= \alpha \int_0^1 \frac{(1 - s)^{n-1}}{(n - 1)!} y(s) ds, \quad B_2 = \beta \int_0^1 \frac{(1 - s)^{n-2}}{(n - 2)!} y(s) ds, \\ B_3 &= \int_0^1 \left(\int_0^s \frac{(s - p)^{n-1}}{(n - 1)!} y(p) dp \right) d\mu(s). \end{aligned}$$

Solving the system (2.9), we obtain

$$c_0 = \frac{E_3 A_2 + E_2 A_3}{E_1 A_2 + A_1 E_2}, \quad c_{n-1} = \frac{E_3 A_1 - A_3 E_1}{A_2 E_1 + A_1 E_2},$$

where $A_2 E_1 + A_1 E_2 \neq 0$. Substituting the values of c_0, c_{n-1} in (2.8), we get the solution (4.3). \square

3. Existence results

In view of Lemma 2.1, we define a fixed point problem related to problem (2.1) as $Fu = u$, where $F : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ is defined by

$$\begin{aligned}
 (Fu)(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds + \sigma_1(t) \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} f(g, u(g)) dg \right) d\mu(s) \right. \\
 &\quad \left. - \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} f(s, u(s)) ds \right] \\
 &\quad + \sigma_2(t) \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} f(s, u(s)) ds,
 \end{aligned}
 \tag{3.1}$$

For the subsequent analysis, we define $\|u\| = \sup_{t \in [0,1]} |u(t)|$, and

$$\vartheta = \left\{ \frac{1}{n!} + h_1 \left[\int_0^1 \frac{s^n}{n!} d\mu(s) + \frac{(\beta n + \alpha)}{n!} \right] + h_2 \frac{\xi^n}{n!} \right\},
 \tag{3.2}$$

where $\max_{t \in [0,1]} |\sigma_1(t)| = h_1$, $\max_{t \in [0,1]} |\sigma_2(t)| = h_2$.

Theorem 3.1. *Assume that $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function satisfying the Lipschitz condition: $|f(t, u) - f(t, v)| \leq \ell|u - v|$, $\ell > 0$, $\forall u, v \in \mathbb{R}, t \in [0, 1]$. Then the boundary value problem (2.1) has a unique solution if $\ell\vartheta < 1$, where ϑ is given by (3.2).*

Proof . In the first step, we show that $FB_a \subset B_a$, where F is the operator defined by (3.1) and $B_a = \{u \in C([0, 1], \mathbb{R}) : \|u\| \leq a\}$, with $a \geq \frac{b\vartheta}{1 - \ell\vartheta}$ and $b = \sup_{t \in [0,1]} |f(t, 0)|$. Using $|f(t, u(t))| = |f(t, u(t)) - f(t, 0) + f(t, 0)| \leq |f(t, u(t)) - f(t, 0)| + |f(t, 0)| \leq \ell\|u\| + b \leq \ell a + b$ for any $u \in B_a, t \in [0, 1]$, we obtain

$$\begin{aligned}
 |(Fu)(t)| &\leq \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds \right. \\
 &\quad + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} |f(g, u(g))| dg \right) d\mu(s) \right. \\
 &\quad \left. + \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} |f(s, u(s))| ds \right] \\
 &\quad \left. + |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds \right\}. \\
 &\leq (\ell a + b) \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} ds + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} dg \right) d\mu(s) \right. \right. \\
 &\quad \left. \left. + \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} ds \right] + |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} ds \right\} \\
 &\leq (\ell a + b)\vartheta \leq a,
 \end{aligned}$$

which implies that $\|Fu\| \leq a$. In consequence, it follows that $FB_a \subset B_a$. Next, for $u, v \in C([0, 1], \mathbb{R})$

and for each $t \in [0, 1]$, we have that

$$\begin{aligned}
 & |(Fu)(t) - (Fv)(t)| \\
 & \leq \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u(s)) - f(s, v(s))| ds \right. \\
 & + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} |f(g, u(g)) - f(g, v(g))| dg \right) d\mu(s) \right. \\
 & + \left. \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} |f(s, u(s)) - f(s, v(s))| ds \right] \\
 & + \left. |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} |f(s, u(s)) - f(s, v(s))| ds \right\} \\
 & \leq \ell \|u - v\| \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} ds \right. \\
 & + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} dg \right) d\mu(s) + \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} ds \right] \\
 & + \left. |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} ds \right\}.
 \end{aligned}$$

Taking maximum over the interval $[0, 1]$, we get $\|(Fu) - (Fv)\| \leq \ell\vartheta \|u - v\|$, where ϑ is given by (3.2). By the assumption: $\ell\vartheta < 1$, we deduce that F is a contraction. Hence, by the contraction mapping principle, problem (2.1) has a unique solution. \square

Example 3.2. Consider the following boundary value problem

$$\begin{cases} u'''(t) = \frac{3}{8} \left(\frac{|u|}{1+|u|} + u + \frac{1}{2} \right), & t \in [0, 1], \\ u(0) = \frac{1}{2}u(1/4), \quad u'(0) = 0, \quad u(1) + u'(1) = \int_0^1 u(s)d\mu(s). \end{cases} \tag{3.3}$$

Here $n = 3$, $\alpha = 1$, $\beta = 1$, $\delta = 1/2$, $\xi = 1/4$, and $f(t, u) = (3/8) \left(\frac{|u|}{1+|u|} + u + \frac{1}{2} \right)$ and $\mu(s) = s^2/2$. Using the given data, we find that $\ell = 3/4$ as $|f(t, u) - f(t, v)| \leq (3/4) \|u - v\|$, $\vartheta \simeq 0.436189$, $\Lambda \simeq 1.390625$. Obviously $\ell\vartheta \simeq 0.327141854 < 1$. Thus, all the conditions of Theorem 3.1 are satisfied. Hence, by the conclusion of Theorem (3.1), problem (3.3) has a unique solution on $[0, 1]$.

Our next existence result is based on Schauder’s fixed point theorem.

Theorem 3.3. Assume that

(A₁) $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous;

(A₂) there exists a positive constant M such that $|f(t, u)| \leq M$ for each $t \in [0, 1]$ and all $u \in \mathbb{R}$.

Then the problem (2.1) has at least one solution on $[0, 1]$.

Proof . We show that the operator F defined by (3.1) satisfies the hypotheses of Schauder’s fixed point theorem. This will be done in several steps.

Step 1: F is continuous.

Let $\{u_k\}$ be a sequence such that $u_k \rightarrow u$ in $C([0, 1], \mathbb{R})$. Then, for each $t \in [0, 1]$, we have

$$\begin{aligned}
 & |F(u_k)(t) - F(u)(t)| \\
 \leq & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u_k(s)) - f(s, u(s))| ds \\
 & + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} |f(g, u_k(g)) - f(g, u(g))| dg \right) d\mu(s) \right. \\
 & + \left. \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} |f(s, u_k(s)) - f(s, u(s))| ds \right] \\
 & + |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} |f(s, u_k(s)) - f(s, u(s))| ds \\
 \leq & \|f(\cdot, u_k(\cdot)) - f(\cdot, u(\cdot))\| \sup_{t \in [0,1]} \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} ds + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} dg \right) d\mu(s) \right. \right. \\
 & + \left. \left. \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} ds \right] + |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} ds \right\} \\
 \leq & \vartheta \|f(\cdot, u_k(\cdot)) - f(\cdot, u(\cdot))\| \rightarrow 0 \text{ as } k \rightarrow \infty,
 \end{aligned}$$

in view of continuity of f (ϑ is given by (3.2)).

Step 2: F maps bounded sets into bounded sets in $C([0, 1], \mathbb{R})$.

Indeed, it is enough to show that for any $\eta^* > 0$, there exists a positive constant L such that for each $u \in B_{\eta^*} = \{u \in C([0, 1], \mathbb{R}) : \|u\| \leq \eta^*\}$, we have $\|F(u)\| \leq L$. For each $t \in [0, 1]$, by the condition (A_2) , we have that

$$\begin{aligned}
 |F(u)(t)| \leq & \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds \\
 & + |\sigma_1(t)| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} |f(g, u(g))| dg \right) d\mu(s) \right. \\
 & + \left. \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} |f(s, u(s))| ds \right] \\
 & + |\sigma_2(t)| \int_0^\xi \frac{(\xi-s)^{n-1}}{(n-1)!} |f(s, u(s))| ds.
 \end{aligned}$$

Taking the norm for $t \in [0, 1]$, the above inequality yields $\|F(u)\| \leq \vartheta L$, where ϑ is given by (3.2).

Step 3: F maps bounded sets into equicontinuous sets of $C([0, 1], \mathbb{R})$.

Let $t_1, t_2 \in (0, 1)$, $t_1 < t_2$, and B_{η^*} be a bounded set in $C([0, 1], \mathbb{R})$ as in Step 2. Then, for $u \in B_{\eta^*}$,

we have

$$\begin{aligned}
 & |F(u)(t_2) - F(u)(t_1)| \\
 \leq & \left| \int_0^{t_1} \frac{[(t_2 - s)^{n-1} - (t_1 - s)^{n-1}]}{(n-1)!} f(s, u(s)) ds + \int_{t_1}^{t_2} \frac{(t_2 - s)^{n-1}}{(n-1)!} f(s, u(s)) ds \right| \\
 & + |\Lambda(1 - \delta)(t_2^{n-1} - t_1^{n-1})| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} f(g, u(g)) dg \right) d\mu(s) \right. \\
 & \left. + \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} f(s, u(s)) ds \right] \\
 & + \left| \Lambda\delta \left(\alpha - \int_0^1 d\mu(s) \right) (t_2^{n-1} - t_1^{n-1}) \right| \int_0^\xi \frac{(\xi - s)^{n-1}}{(n-1)!} f(s, u(s)) ds. \\
 \leq & M \left\{ \left| \int_0^{t_1} \frac{[(t_2 - s)^{n-1} - (t_1 - s)^{n-1}]}{(n-1)!} ds + \int_{t_1}^{t_2} \frac{(t_2 - s)^{n-1}}{(n-1)!} ds \right| \right. \\
 & + |\Lambda(1 - \delta)(t_2^{n-1} - t_1^{n-1})| \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} dg \right) d\mu(s) \right. \\
 & \left. + \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} ds \right] \\
 & \left. + \left| \Lambda\delta \left(\alpha - \int_0^1 d\mu(s) \right) (t_2^{n-1} - t_1^{n-1}) \right| \int_0^\xi \frac{(\xi - s)^{n-1}}{(n-1)!} ds \right\}. \\
 \leq & \frac{M}{n!} [2(t_2 - t_1)^n + |t_2^n - t_1^n|] \\
 & + M |\Lambda(1 - \delta)(t_2^{n-1} - t_1^{n-1})| \left[\int_0^1 \frac{s^n}{n!} d\mu(s) + \frac{(\beta n + \alpha)}{n!} \right] \\
 & + \frac{M\xi^n}{n!} \left| \Lambda\delta \left(\alpha - \int_0^1 d\mu(s) \right) (t_2^{n-1} - t_1^{n-1}) \right|.
 \end{aligned}$$

Clearly the right hand side of the above inequality tends to zero independent of u as $(t_2 - t_1) \rightarrow 0$. In view of the above three Steps, the Arzelá-Ascoli theorem applies and consequently the operator $F : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ is continuous and completely continuous.

Step 4: A priori bounds.

We show that the set $\varepsilon = \{u \in C([0, 1], \mathbb{R}) : u = \lambda F(u) \text{ for some } 0 < \lambda < 1\}$ is bounded. Let $u \in \varepsilon$. Then $u = \lambda F(u)$ for some $0 < \lambda < 1$. Thus, for each $t \in [0, 1]$, we have

$$\begin{aligned}
 u(t) &= \lambda \left\{ \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds + \sigma_1(t) \left[\int_0^1 \left(\int_0^s \frac{(s-g)^{n-1}}{(n-1)!} f(g, u(g)) dg \right) d\mu(s) \right. \right. \\
 & \left. \left. - \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} f(s, u(s)) ds \right] \right. \\
 & \left. + \sigma_2(t) \int_0^\xi \frac{(\xi - s)^{n-1}}{(n-1)!} f(s, u(s)) ds \right\}.
 \end{aligned}$$

Using the condition (A_2) , it is easy to show that $\|F(u)\| \leq M\vartheta$. This shows that set ε is bounded. Thus, it follows by Schauder’s fixed point theorem that the operator F has a fixed point, which is a solution of the problem (2.1). \square

Example 3.4. Consider the following nonlinear boundary value problem

$$\begin{cases} u^{(4)}(t) = \frac{e^{-u^2(t)} + 2 \sin(1 + 3u(t)) + \cos(3 + 5u^3(t)) + 3t^4}{1 + u^2(t)}, & t \in [0, 1], \\ u(0) = u(1/3), \quad u'(0) = 0, \quad u''(0) = 0, \quad \frac{2}{3}u(1) + \frac{3}{4}u'(1) = \int_0^1 u(s)d\mu(s). \end{cases} \quad (3.4)$$

Here $f(t, u(t)) = \frac{e^{-u^2(t)} + 2 \sin(1 + 3u(t)) + \cos(3 + 5u^3(t)) + 3t^4}{1 + u^2(t)}$. Clearly $f(t, u(t))$ is continuous and $|f(t, u(t))| \leq M$ with $M = 7$ for each $t \in [0, 1]$ and all $u \in \mathbb{R}$. Thus the conclusion of Theorem 3.3 applies and the problem (3.4) has a solution on $[0, 1]$.

4. Some related problems

In this section, we study two more n th-order boundary value problems involving Stieltjes type integral boundary conditions.

4.1. Problem I

We consider the following n th-order ordinary differential equation

$$u^{(n)}(t) = f(t, u(t)), \quad t \in [0, 1], \quad (4.1)$$

supplemented with the Stieltjes integral boundary conditions:

$$\begin{cases} u(0) = \delta \int_0^\xi u(s)d\tau(s), \quad u'(0) = 0, \quad u''(0) = 0, \dots, u^{(n-2)}(0) = 0, \\ \alpha u(1) + \beta u'(1) = \sum_{i=1}^m \gamma_i \int_0^{\beta_i} u(s)d\psi(s), \quad 0 < \xi < \beta_1 < 1, \end{cases} \quad (4.2)$$

where $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a given continuous function, $\alpha, \beta, \gamma_i, \delta, \xi, \beta_i$ ($i = 1, 2, \dots, m$) are real constants to be chosen appropriately, $\tau(s)$ and $\psi(s)$ are functions of bounded variation.

Observe that the problem (4.1)-(4.2) differs from the problem (2.1) in the sense that it considers the boundary conditions:

$$u(0) = \delta \int_0^\xi u(s)d\tau(s), \quad \alpha u(1) + \beta u'(1) = \sum_{i=1}^m \gamma_i \int_0^{\beta_i} u(s)d\psi(s)$$

instead of the following conditions assumed in the problem (2.1):

$$u(0) = \delta u(\xi), \quad \alpha u(1) + \beta u'(1) = \int_0^1 u(s)d\mu(s),$$

whereas the other conditions remain the same.

Lemma 4.1. *The unique solution of problem (4.1)-(4.2) is equivalent to the integral equation:*

$$\begin{aligned}
 u(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds \\
 &+ \lambda_1(t) \left[\sum_{i=1}^m \gamma_i \int_0^{\beta_i} \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\psi(s) \right. \\
 &\left. - \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} f(s, u(s)) ds \right] \\
 &+ \lambda_2(t) \int_0^\xi \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\tau(s),
 \end{aligned} \tag{4.3}$$

where

$$\lambda_1(t) = \frac{1}{\chi} \left[\delta \int_0^\xi s^{n-1} d(\tau(s)) + t^{n-1} \left(1 - \delta \int_0^\xi d\tau(s) \right) \right], \tag{4.4}$$

$$\begin{aligned}
 \lambda_2(t) &= \frac{1}{\chi} \left[\alpha + (n-1)\beta - \sum_{i=1}^m \gamma_i \int_0^{\beta_i} s^{n-1} d\psi(s) \right. \\
 &\left. - t^{n-1} \left(\alpha - \sum_{i=1}^m \gamma_i \int_0^{\beta_i} d\psi(s) \right) \right],
 \end{aligned} \tag{4.5}$$

$$\begin{aligned}
 \chi &= \left(\alpha - \sum_{i=1}^m \gamma_i \int_0^{\beta_i} d\psi(s) \right) \left(\delta \int_0^\xi s^{n-1} d\tau(s) \right) \\
 &- \left(\alpha + (n-1)\beta - \sum_{i=1}^m \gamma_i \int_0^{\beta_i} s^{n-1} d\psi(s) \right) \left(1 - \delta \int_0^\xi d\tau(s) \right) \neq 0.
 \end{aligned} \tag{4.6}$$

Proof . The proof is similar to that of Lemma 2.1. So we omit it. \square

By Lemma 4.1, we consider a fixed point problem associated with the problem (4.1)-(4.2) as $Gu = u$, where the operator $G : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ is defined by

$$\begin{aligned}
 (Gu)(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds \\
 &+ \lambda_1(t) \left[\sum_{i=1}^m \gamma_i \int_0^{\beta_i} \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\psi(s) \right. \\
 &\left. - \int_0^1 \frac{(1-s)^{n-2} [\beta(n-1) + \alpha(1-s)]}{(n-1)!} f(s, u(s)) ds \right] \\
 &+ \lambda_2(t) \int_0^\xi \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\tau(s).
 \end{aligned} \tag{4.7}$$

Moreover, we set

$$\vartheta_I = \left\{ \frac{1}{n!} + m_1 \left[\sum_{i=1}^m \gamma_i \int_0^{\beta_i} \frac{s^n}{n!} d\psi(s) + \frac{(\beta n + \alpha)}{n!} \right] + m_2 \int_0^\xi \frac{s^n}{n!} d\tau(s) \right\}, \tag{4.8}$$

where $\max_{t \in [0,1]} |\lambda_1(t)| = m_1$, $\max_{t \in [0,1]} |\lambda_2(t)| = m_2$ (λ_1 and λ_2 are respectively given by (4.4) and (4.5)).

Following the method of proof used for obtaining the two existence results for the problem (2.1) in the previous section, we can establish the similar results for the problem (4.1)-(4.2) with the aid of the operator G defined by (4.7) and the constant ϑ_I given by (4.8). The existence results for the problem (4.1)-(4.2) can be formulated as follows.

Theorem 4.2. *Let $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function satisfying the Lipschitz condition: $|f(t, u) - f(t, v)| \leq \ell_1|u - v|$, $\ell_1 > 0$, $\forall u, v \in \mathbb{R}$, $t \in [0, 1]$. Then there exists a unique solution for the problem (4.1)-(4.2) on $[0, 1]$ if $\ell_1\vartheta_I < 1$, where ϑ_I is given by (4.8).*

Theorem 4.3. *Assume that the function $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and there exists a positive constant M_1 such that $|f(t, u)| \leq M_1$ for each $t \in [0, 1]$ and for all $u \in \mathbb{R}$. Then the problem (4.1)-(4.2) has at least one solution on $[0, 1]$.*

Example 4.4. *Consider the third-order boundary value problem given by*

$$\begin{cases} u'''(t) = f(t, u(t)), & t \in [0, 1], \\ u(0) = \frac{1}{2} \int_0^\xi u(s) d\tau(s), \quad u'(0) = 0, \quad u(1) + u'(1) = \sum_{i=1}^4 \gamma_i \int_0^{\beta_i} u(s) d\psi(s). \end{cases} \quad (4.9)$$

Here $n = 3$, $\alpha = 1$, $\beta = 1$, $\delta = 1/2$, $\xi = 1/4$, $m = 4$, $\beta_1 = 1/3$, $\beta_2 = 1/2$, $\beta_3 = 2/3$, $\beta_4 = 3/4$, $\gamma_1 = 3/4$, $\gamma_2 = 2/3$, $\gamma_3 = 1/2$, $\gamma_4 = 1/3$, $f(t, u) = \frac{2|u|}{3(1+|u|)} + \frac{u}{3} + e^t$, $\tau(s) = s$, and $\psi(s) = s^2$. Using the given values, it is found that $\chi \simeq 1.083833$, $\vartheta_I \simeq 0.715664$ and $\ell_1 = 1$ as $|f(t, u) - f(t, v)| \leq \|u - v\|$. Obviously $\ell_1\vartheta_I \simeq 0.715664 < 1$. Thus, all the conditions of Theorem (4.2) are satisfied. In consequence, by the conclusion of Theorem (4.2), the problem (4.9) has a unique solution on $[0, 1]$.

4.2. Problem II

We replace the condition ‘ $\alpha u(1) + \beta u'(1) = \sum_{i=1}^m \gamma_i \int_0^{\beta_i} u(s) d\psi(s)$ ’ by ‘ $\alpha u(\eta) + \beta u'(\eta) = \int_\zeta^1 u(s) d\rho(s)$ ’ in (4.2) and consider the following problem:

$$\begin{cases} u^{(n)}(t) = f(t, u(t)), & t \in [0, 1], \\ u(0) = \delta \int_0^\xi u(s) d\mu(s), \quad u'(0) = 0, \quad u''(0) = 0, \dots, \quad u^{(n-2)}(0) = 0, \\ \alpha u(\eta) + \beta u'(\eta) = \int_\zeta^1 u(s) d\rho(s), \quad 0 < \xi < \eta < \zeta < 1, \end{cases} \quad (4.10)$$

where $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is a given continuous function, and α, β, δ are real constants to be chosen appropriately, $\mu(s)$ and $\rho(s)$ are functions of bounded variation.

As before, associated with the problem (4.10), we define an operator $H : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$ as

$$\begin{aligned} (Hu)(t) &= \int_0^t \frac{(t-s)^{n-1}}{(n-1)!} f(s, u(s)) ds + \nu_1(t) \int_0^\xi \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\mu(s) \\ &+ \nu_2(t) \left[\int_\zeta^1 \left(\int_0^s \frac{(s-x)^{n-1}}{(n-1)!} f(x, u(x)) dx \right) d\rho(s) \right. \\ &\left. - \alpha \int_0^\eta \frac{(\eta-s)^{n-1}}{(n-1)!} f(s, u(s)) ds - \beta \int_0^\eta \frac{(\eta-s)^{n-2}}{(n-2)!} f(s, u(s)) ds \right], \end{aligned} \quad (4.11)$$

where

$$\begin{aligned}
 \nu_1(t) &= \frac{\delta}{\gamma} \left[\alpha \eta^{n-1} + \beta(n-1)\eta^{n-2} - \int_{\zeta}^1 s^{n-1} d\rho(s) - t^{n-1} \left(\alpha - \int_{\zeta}^1 d(\rho(s)) \right) \right] \\
 \nu_2(t) &= \frac{1}{\gamma} \left[\delta \int_0^{\xi} s^{n-1} d\mu(s) + t^{n-1} \left(1 - \delta \int_0^{\xi} d\mu(s) \right) \right], \\
 \gamma &= \left(\alpha \eta^{n-1} + \beta(n-1)\eta^{n-2} - \int_{\zeta}^1 s^{n-1} d\rho(s) \right) \left(1 - \delta \int_0^{\xi} d\mu(s) \right) \\
 &+ \left(\alpha - \int_{\zeta}^1 d\rho(s) \right) \left(\delta \int_0^{\xi} s^{n-1} d\mu(s) \right) \neq 0.
 \end{aligned} \tag{4.12}$$

Notice that the problem (4.10) has solutions only if the operator equation $Hu = u$ has fixed points. In the sequel, we use the notation:

$$Q_{II} = \left\{ \frac{1}{n!} + n_1 \int_0^{\xi} \frac{s^n}{n!} d\mu(s) + n_2 \left[\int_{\zeta}^1 \frac{s^n}{n!} d(\rho(s)) + \frac{\alpha \eta^n}{n!} + \frac{\beta \eta^{n-1}}{(n-1)!} \right] \right\}. \tag{4.13}$$

where $\max_{t \in [0,1]} |\nu_i(t)| = n_i, i = 1, 2$.

Now we present the existence results for the problem (4.10). The method of proof for these results is similar to the one employed in Section 3, so we omit the proofs.

Theorem 4.5. *Assume that the function $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and satisfies the Lipschitz condition: $|f(t, u) - f(t, v)| \leq \ell_2 |u - v|, \ell_2 > 0, \forall u, v \in \mathbb{R}, \ell_2 > 0 t \in [0, 1]$. Then the problem (4.10) has a unique solution on $[0, 1]$ provided that $\ell_2 Q_{II} < 1$, where Q_{II} is given by (4.13).*

Theorem 4.6. *Assume that the following conditions hold:*

(B₁) *the function $f : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous;*

(B₂) *there exists a positive constant N such that $|f(t, u)| \leq N$ for each $t \in [0, 1]$ and for all $u \in \mathbb{R}$.*

Then there exists at least one solution for the problem (4.10) on $[0, 1]$.

Example 4.7. *Let us consider the problem (4.10) with $n = 3, \alpha = 1, \beta = 1, \delta = 1/2, \xi = 1/4, \zeta = 1/3, \eta = 3/4$, and $f(t, u) = \frac{2e^t |u|}{5(1 + |u|)} + 3, \mu(s) = \frac{s^2}{2}$ and $\rho(s) = \frac{s^3}{3}$.*

Using the given data, we find that $\gamma \simeq 1.834539, Q_{II} \simeq 0.370316$ and $\ell_2 = 2/5$ as $|f(t, u) - f(t, v)| \leq (2/5) \|u - v\|$. Clearly

$$\ell_2 Q_{II} \simeq 0.148126 < 1.$$

Since all the conditions of Theorem 4.5 are satisfied, therefore, the conclusion of Theorem 4.5 applies and the problem (4.10) with the chosen data has a unique solution on $[0, 1]$.

References

- [1] M. Greguš, F. Neumann and FM. Arscott, *Three-point boundary value problems in differential equations*, Proc. London Math. Soc. 3 (1964) 459–470.
- [2] AV. Bicadze and AA. Samarskii, *Some elementary generalizations of linear elliptic boundary value problems (Russian)*, Anal. Dokl. Akad. Nauk SSSR, 185 (1969) 739–740.
- [3] VA. Il'in and EI. Moiseev, *A nonlocal boundary value problem of the first kind for the Sturm-Liouville operator in differential and difference interpretations (Russian)*, Differential'nye Uravneniya 23 (1987) 1198–1207.
- [4] J. Andres, *A four-point boundary value problem for the second-order ordinary differential equations*, Arch. Math. (Basel) 53 (1989) 384–389.
- [5] CP. Gupta, *Solvability of a three-point nonlinear boundary value problem for a second order ordinary differential equations*, J. Math. Anal. Appl. 168 (1998), 540–551.
- [6] PW. Elloe and B. Ahmad, *Positive solutions of a nonlinear n th order boundary value problem with nonlocal conditions*, Appl. Math. Lett. 18 (2005) 521–527.
- [7] S. Clark and J. Henderson, *Uniqueness implies existence and uniqueness criterion for non local boundary value problems for third-order differential equations*, Proc. Amer. Math. Soc. 134 (2006), 3363–3372.
- [8] JRL. Webb and G. Infante, *Positive solutions of nonlocal boundary value problems: A unified approach*, J. London Math. Soc. 74 (2006) 673–693.
- [9] J.R. Graef, J.R.L. Webb, *Third order boundary value problems with nonlocal boundary conditions*, Nonlinear Anal. 71 (2009) 1542–1551.
- [10] Y. Sun, L. Liu, J. Zhang and RP. Agarwal, *Positive solutions of singular three-point boundary value problems for second-order differential equations*, J. Comput. Appl. Math. 230 (2009) 738–750.
- [11] FT. Akyildiz, H. Bellout, K. Vajravelu and RA. Van Gorder, *Existence results for third order nonlinear boundary value problems arising in nano boundary layer fluid flows over stretching surfaces*, Nonlinear Anal. Real World Appl. 12 (2011) 2919–2930.
- [12] JR. Graef, L. Kong, Q. Kong and M. Wang, *Uniqueness and parameter dependence of positive solutions to higher order boundary value problems with fractional q -derivatives*, J. Appl. Anal. Comput. 3 (2013) 21–35.
- [13] G. Infante, *Eigenvalues and positive solutions of ODEs involving integral boundary conditions*, Discrete Contin. Dyn. Syst. (2005), suppl., 436–442.
- [14] Z. Yang, *Positive solutions of a second order integral boundary value problem*, J. Math. Anal. Appl. 321 (2006), 751–765.
- [15] B. Ahmad and A. Alsaedi, *Existence of approximate solutions of the forced Duffing equation with discontinuous type integral boundary conditions*, Nonlinear Anal. Real World Appl. 10 (2009), 358–367.
- [16] A. Boucherif, *Second-order boundary value problems with integral boundary conditions*, Nonlinear Anal. 70 (2009) 364–371.
- [17] B. Ahmad, S.K. Ntouyas, H.H. Alsulami, *Existence results for n -th order multipoint integral boundary-value problems of differential inclusion*, Electron. J. Differential Equations (2013) No. 203, 13 pp.
- [18] J. Xu, D. O'Regan and Z. Yang, *Positive solutions for a n th-order impulsive differential equation with integral boundary conditions*, Differ. Equ. Dyn. Syst. 22 (2014) 427–439.
- [19] Y. Li and H. Zhang, *Positive solutions for a nonlinear higher order differential system with coupled integral boundary conditions*, J. Appl. Math. (2014) Art. ID 901094, 7 pp.
- [20] J. Henderson, *Smoothness of solutions with respect to multi-strip integral boundary conditions for n th order ordinary differential equations*, Nonlinear Anal. Model. Control 19 (2014) 396–412.
- [21] IY. Karaca and FT. Fen, *Positive solutions of n th-order boundary value problems with integral boundary conditions*, Math. Model. Anal. 20 (2015) 188–204.
- [22] WM. Whyburn, *Differential equations with general boundary conditions*, Bull. Amer. Math. Soc. 48 (1942) 692–704.
- [23] R. Conti, *Recent trends in the theory of boundary value problems for ordinary differential equations*, Boll. Un. Mat. Ital. 22 (1967) 135–1788.
- [24] JRL. Webb, *Nonlocal conjugate type boundary value problems of higher order*, Nonlinear Anal. 71 (2009) 1933–1940.
- [25] JRL. Webb, *Positive solutions of some higher order nonlocal boundary value problems*, Electron. J. Qual. Theory Differ. Equ. (2009), Special Edition I, No. 29, 15 pp.
- [26] JRL. Webb and G. Infante, *Non-local boundary value problems of arbitrary order*, J. Lond. Math. Soc. 79 (2009) 238–258.
- [27] JRL. Webb and G. Infante, *Positive solutions of nonlocal boundary value problems involving integral conditions*, NoDEA Nonlinear Differential Equations Appl. 15 (2008) 45–67.

- [28] Y. Wang, L. Liu, Y. Wu, *Positive solutions for a nonlocal fractional differential equation*, *Nonlinear Anal.* 74 (2011) 3599–3605.
- [29] X. Zhang and Y. Han, *Existence and uniqueness of positive solutions for higher order nonlocal fractional differential equations*, *Appl. Math. Lett.* 25 (2012) 555–560.
- [30] B. Ahmad and SK. Ntouyas, *Existence results for fractional differential inclusions arising from real estate asset securitization and HIV models*, *Adv. Difference Equ.* (2013), 2013:216, 15 pp.