

An Analog of Titchmarsh's Theorem for the Dunkl Transform in the Space $L^2_\alpha(\mathbb{R})$

R. Daher, M. El Hamma *

Department of Mathematics, Faculty of Science Ain Chock, University Hassan II, Casablanca, Morocco.

(Communicated by A. Ebadian)

Abstract

In this paper, using a generalized Dunkl translation operator, we obtain an analog of Titchmarsh's Theorem for the Dunkl transform for functions satisfying the Lipschitz-Dunkl condition in $L_{2,\alpha} = L^2_\alpha(\mathbb{R}) = L^2(\mathbb{R}, |x|^{2\alpha+1}dx)$, $\alpha > \frac{-1}{2}$.

Keywords: Dunkl Operator, Dunkl Transform, Generalized Dunkl Translation.
2010 MSC: 44A15, 46E30.

1. Introduction and Preliminaries

Dunkl operators are differential-difference operators introduced in 1989, by Dunkl [5]. On the real line, these operators, which are denoted by $D = D_\alpha$, depend on a real parameter $\alpha > \frac{-1}{2}$ and they associated with the reflection group \mathbb{Z}_2 on \mathbb{R} . For more details about these operators see [4, 5, 6, 7, 10] and the references therein.

Recently in mathematical papers a new class of generalized translations was described and put into use, namely, the generalized Dunkl translations. The generalized Dunkl translations are constructed on the base of certain differential-difference operators which are widely used in mathematical physics (e.g., [5, 9]).

Titchmarsh's [11, Thorem 85] characterized the set of functions in $L_{2,\alpha}$ satisfying the cauchy Lipschitz condition by means of an asymptotic estimate growth of the norm of their Fourier transform, we have

*Corresponding author

Email addresses: ra_daher@yahoo.fr (R. Daher), m_elhamma@yahoo.fr (M. El Hamma)

Theorem 1.1. [11] Let $\alpha \in (0, 1)$ and assume that $f \in L^2(\mathbb{R})$. Then the following are equivalents:

$$(1) \|f(t+h) - f(t)\|_{L^2(\mathbb{R})} = O(h^\alpha) \text{ as } h \rightarrow 0$$

$$(2) \int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 d\lambda = O(r^{-2\alpha}) \quad r \rightarrow \infty$$

where \widehat{f} stands for the Fourier transform of f .

In this paper, we obtain an analog of Theorem 1.1 in the Dunkl operator setting by means of generalized Dunkl translation. We point out that similar results have been established in the context of noncompact rank 1 Riemannian symmetric spaces [8].

The Dunkl operator is differential-difference operator D which satisfies the condition

$$Df(x) = \frac{df}{dx}(x) + \left(\alpha + \frac{1}{2}\right) \frac{f(x) - f(-x)}{x}, \quad f \in L_{2,\alpha}.$$

Let $j_\alpha(x)$ be a normalized Bessel function of the first kind, i.e.,

$$j_\alpha(x) = \frac{2^\alpha \Gamma(\alpha + 1) J_\alpha(x)}{x^\alpha},$$

where $J_\alpha(x)$ is a Bessel function of the first kind ([2], chap 7). The function $j_\alpha(x)$ is infinitely differentiable and even.

We understand a generalized exponential function as the function

$$e_\alpha(x) = j_\alpha(x) + ic_\alpha x j_{\alpha+1}(x), \tag{1.1}$$

where $c_\alpha = (2\alpha + 2)^{-1}$.

The function $y = e_\alpha(x)$ satisfies the equation $Dy = y$.

Using the correlation

$$j'_\alpha(x) = -\frac{x j_{\alpha+1}(x)}{2(\alpha + 1)}.$$

we conclude that the function $e_\alpha(x)$ admits the representation

$$e_\alpha(x) = j_\alpha(x) - i j'_\alpha(x) \tag{1.2}$$

Let $L_{2,\alpha}$ is the Hilbert space consists of measurable functions $f(x)$ defined on \mathbb{R} with the norm

$$\|f\| = \|f\|_{2,\alpha} = \left(\int_{-\infty}^{+\infty} |f(x)|^2 |x|^{2\alpha+1} dx \right)^{1/2}.$$

The Dunkl transform is defined by

$$\widehat{f}(\lambda) = \int_{-\infty}^{+\infty} f(x) e_\alpha(\lambda x) |x|^{2\alpha+1} dx, \quad \lambda \in \mathbb{R}.$$

The inverse Dunkl transform is defined by the formula

$$f(x) = (2^{\alpha+1}\Gamma(\alpha + 1))^{-2} \int_{-\infty}^{+\infty} \widehat{f}(\lambda)e_{\alpha}(-\lambda x)|\lambda|^{2\alpha+1}d\lambda .$$

The Dunkl transform satisfies the Parseval equality

$$\int_{-\infty}^{+\infty} |f(x)|^2|x|^{2\alpha+1}dx = (2^{\alpha+1}\Gamma(\alpha + 1))^{-2} \int_{-\infty}^{+\infty} |\widehat{f}(\lambda)||\lambda|^{2\alpha+1}d\lambda .$$

In $L_{2,\alpha}$, we define the operator of the generalized Dunkl translation (see [3])

$$T_h f(x) = C \left(\int_0^\pi f_e(G(x, h, \varphi))h^e(x, h, \varphi)\sin^{2\alpha}\varphi d\varphi + \int_0^\pi f_0(G(x, h, \varphi))h^0(x, h, \varphi)\sin^{2\alpha}\varphi d\varphi \right)$$

where

$$C = \frac{\Gamma(\alpha + 1)}{\Gamma(\frac{1}{2})\Gamma(\alpha + \frac{1}{2})}, \quad G(x, h, \varphi) = \sqrt{x^2 + h^2 - 2|xh|\cos\varphi}$$

$$h^e(x, h, \varphi) = 1 - \operatorname{sgn}(xh)\cos\varphi .$$

and

$$h^0(x, h, \varphi) = \frac{(x + h)h^e(x, h, \varphi)}{G(x, h, \varphi)} \text{ for } (x, h) \neq (0, 0)$$

$$h^0(x, h, \varphi) = 0 \text{ for } (x, h) = (0, 0)$$

$$f_e(x) = \frac{1}{2}(f(x) + f(-x)), \quad f_0(x) = \frac{1}{2}(f(x) - f(-x))$$

From formula (1.1), we have

$$|1 - j_{\alpha}(x)| \leq |1 - e_{\alpha}(x)| .$$

For $\alpha \geq \frac{-1}{2}$, we introduce the Bessel normalized function of the first kind j_{α} defined by

$$j_{\alpha}(x) = \Gamma(p + 1) \sum_{n=0}^{\infty} \frac{(-1)^n}{n!\Gamma(n + \alpha + 1)} \left(\frac{x}{2}\right)^{2n} \tag{1.3}$$

Moreover, from (1.3) we see that

$$\lim_{x \rightarrow 0} \frac{j_p(x) - 1}{x^2} \neq 0 ,$$

by consequence, there exist $c > 0$ and $\eta > 0$ satisfying

$$|x| \leq \eta \implies |j_{\alpha}(x) - 1| \geq c|x|^2 \tag{1.4}$$

In [3], we obtain

$$(\widehat{T_h f})(\lambda) = e_{\alpha}(\lambda h)\widehat{f}(\lambda) \tag{1.5}$$

2. Main Result

In this section we give the main result of this paper, We need first to define the Dunkl-Lipschitz class.

Definition 2.1. Let $\beta \in (0, 1)$. A function $f \in L_{2,\alpha}$ is said to be in the Dunkl-Lipschitz class, denote by $Lip(\beta, 2, \alpha)$, if

$$\|T_h f(x) + T_{-h} f(x) - 2f(x)\| = O(h^\beta), \text{ as } h \rightarrow 0.$$

Theorem 2.2. Let $f \in L_{2,\alpha}$. Then the following are equivalent

- (1) $f \in Lip(\beta, 2, \alpha)$.
- (2) $\int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda = O(r^{-2\beta})$ as $r \rightarrow +\infty$.

Proof . 1 \implies 2: Assume that $f \in Lip(\beta, 2, \alpha)$. Then we have

$$\|T_h f(x) + T_{-h} f(x) - 2f(x)\| = O(h^\beta), \text{ as } h \rightarrow 0.$$

From formulas (1.2) and (1.5), we have the Dunkl transform of $T_h f(x) + T_{-h} f(x) - 2f(x)$ is $2(j_\alpha(\lambda h) - 1)$.

Parseval's identity gives

$$\|T_h f(x) + T_{-h} f(x) - 2f(x)\| = \int_{-\infty}^{+\infty} |2(1 - j_\alpha(\lambda h))|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda.$$

From (1.4), we have

$$\int_{\frac{\eta}{2h} \leq |\lambda| \leq \frac{\eta}{h}} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \geq \frac{c^2 \eta^4}{16} \int_{\frac{\eta}{2h} \leq |\lambda| \leq \frac{\eta}{h}} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda.$$

There exists then a positive constant C such that

$$\begin{aligned} \int_{\frac{\eta}{2h} \leq |\lambda| \leq \frac{\eta}{h}} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda &\leq C \int_{-\infty}^{+\infty} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \\ &\leq Ch^{2\beta} \end{aligned}$$

For all $h > 0$, we have

$$\int_{r \leq |\lambda| \leq 2r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \leq Cr^{-2\beta}, \text{ for all } r > 0.$$

Furthermore, we obtain

$$\begin{aligned} \int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda &= \sum_{i=0}^{\infty} \int_{2^i r \leq |\lambda| \leq 2^{i+1} r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \\ &\leq C \sum_{i=0}^{\infty} (2^i r)^{-2\beta} \\ &\leq C r^{-2\beta} \end{aligned}$$

This prove that

$$\int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda = O(r^{-2\beta}) \text{ as } r \rightarrow +\infty.$$

2 \implies 1: Suppose now that

$$\int_{|\lambda| \geq r} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda = O(r^{-2\beta}) \text{ as } r \rightarrow +\infty.$$

We write

$$\begin{aligned} \int_{-\infty}^{+\infty} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda &= \int_{|\lambda| < \frac{1}{h}} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \\ &\quad + \int_{|\lambda| \geq \frac{1}{h}} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \end{aligned}$$

Firstly, we use the formula $|j_\alpha(x)| \leq 1$

$$\int_{|\lambda| \geq \frac{1}{h}} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda \leq 4 \int_{|\lambda| \geq \frac{1}{h}} |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda.$$

Then

$$\int_{|\lambda| \geq \frac{1}{h}} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda = O(h^{2\beta}) \text{ as } h \rightarrow 0.$$

Set

$$\phi(x) = \int_x^\infty |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda.$$

An integration by parts, we obtain

$$\begin{aligned} \int_0^x \lambda^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda &= \int_0^x -\lambda^2 \phi'(\lambda) d\lambda = -x^2 \phi(x) + 2 \int_0^x \lambda \phi(\lambda) d\lambda \\ &\leq 2 \int_0^x O(\lambda^{1-2\beta}) d\lambda \\ &= O(x^{2-2\beta}). \end{aligned}$$

We use the formula $1 - j_\alpha(x) = O(x^2)$, $0 \leq x \leq 1$ see [1]. Then

$$\begin{aligned}
\int_{-\infty}^{+\infty} |1 - j_\alpha(\lambda h)|^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda &= O(h^2) \int_{|\lambda| < \frac{1}{h}} \lambda^2 |\widehat{f}(\lambda)|^2 |\lambda|^{2\alpha+1} d\lambda + O(h^{2\beta}) \\
&= O(h^2 h^{-2+2\beta}) + O(h^{2\beta}) \\
&= O(h^{2\beta}) + O(h^{2\beta}) \\
&= O(h^{2\beta}).
\end{aligned}$$

and this end the proof. \square

References

- [1] V. A. Abilov and F. V. Abilova, *Approximation of Functions by Fourier-Bessel Sums*, Izv. Vyssh. Uchebn. Zaved., Mat., (8) (2001), 3–9.
- [2] H. Bateman and A. Erdélyi, *Higher Transcendental Functions*, vol. II (McGraw-Hill, New York, 1953; Nauka, Moscow, 1974).
- [3] E. S. Belkina and S. S. Platonov, *Equivalence of K -Functionals and Modulus of Smoothness Constructed by Generalized Dunkl Translations*, Izv. Vyssh. Uchebn. Zaved. Mat., (8) (2008) 3–15.
- [4] M. F. E. de Jeu, *The Dunkl transform*, Invent. Math., 113 (1993) 147–162.
- [5] C. F. Dunkl, *Differential-difference operators associated to reflection groups*, Trans. Amer. Math. Soc., 311 (1) (1989) 167–183.
- [6] C. F. Dunkl, *Hankel transforms associated to finite reflection groups. Hypergeometric functions on domains of positivity, Jack polynomials, and applications*, Contemp. Math., 138 (1992) 123–138.
- [7] L. Lapointe and L. Vinet, *Exact operator solution of the Calogero-Sutherland model*, J. Phys., 178 (1996) 425–452.
- [8] S. S. Platonov, *The Fourier transform of functions satisfying the Lipschitz condition on rank 1 symmetric spaces*, Siberian Math. J., 46 (6) (2005) 1108–1118.
- [9] M. Rösler, *Dunkl Operators: Theory and Applications*, Lect. Notes in Math., 1817 (2002) 93–135.
- [10] M. Rösler, *Positivity of the Dunkl intertwining operator*, Duke Math. J., 98 (3) (1999) 445–463.
- [11] E. C. Titchmarsh, *Introduction to the theory of Fourier Integrals*, Clarendon, Oxford, 1948, Komkniga, Moscow, 2005.