



# Weak subsequential continuity in fuzzy metric spaces and application

Said Beloul<sup>a,\*</sup>, Anita Tomar<sup>b</sup>, Ritu Sharma<sup>c</sup>

<sup>a</sup>Operators theory and PDE Laboratory, Department of Mathematics, Faculty of Exact Sciences, University of El Oued, P.O.Box789, El-Oued 39000, Algeria

<sup>b</sup>Department of Mathematics, Government Degree College Thatyur, Tehri Garhwal (Uttarakhand), India

<sup>c</sup>Department of Mathematics, V. S. K. C. Government P. G. College Dakpathar, Dehradun (Uttarakhand), India

(Communicated by Reza Saadati)

---

## Abstract

Compatibility of type (E) and weak subsequential continuity is utilized in a fuzzy metric space for the existence of a common fixed point. Illustrations and an application are stated to elucidate our outcomes.

*Keywords:* Compatibility of type (E), fuzzy metric space, dynamic programming functional equation, weak subsequential continuity.

*2010 MSC:* 47H10, 54H25.

---

## 1. Introduction

The fuzzy metric space is one of the consequential generalizations of metric space due to its interesting applications in stability theory, applied science, mathematical programming, engineering science, modelling theory, medical science, control theory, image processing and communication. The idea was initiated by Zadeh [20], where he defined the fuzzy sets. Later Kramosil and Michalek [7] familiarised with the fuzzy metric space which is further improved by George and Veeramanti [6] using the continuous t-norms [5, 6, 7, 8, 10, 11, 13, 14, 17, 20]. Now we establish common fixed point exploiting compatibility of type (E) and weak subsequential continuity to demonstrate the usefulness of these notions for contractive,  $\phi$ -contractive as well as an integral type contractive condition in the fuzzy metric space. In the sequel illustrations and an application to solve a functional equations is also stated to elucidate our outcomes.

---

\*Corresponding author

*Email addresses:* [beloulsaid@gmail.com](mailto:beloulsaid@gmail.com) (Said Beloul), [anitatmr@yahoo.com](mailto:anitatmr@yahoo.com) (Anita Tomar), [ritus4184@gmail.com](mailto:ritus4184@gmail.com) (Ritu Sharma)

*Received:* April 2020    *Accepted:* July 2020

## 2. Preliminaries

**Definition 2.1.** [2] Let  $\exists$  a sequence  $\{x_n\} \in X$  satisfying  $\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = z \in X$ . A pair  $(\mathcal{A}, \mathcal{H})$  over the standard metric space  $(X, d)$  is

1. weakly subsequentially continuous iff  $\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{A}z$  or  $\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{H}z$ .
2.  $\mathcal{H}$ -subsequentially continuous,  $\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{H}z$ .
3.  $\mathcal{A}$ -subsequentially continuous, iff  $\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{A}z$ .
4. compatible of type (E) [15] if

$$\lim_{n \rightarrow \infty} \mathcal{H}^2x_n = \lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{A}z$$

and

$$\lim_{n \rightarrow \infty} \mathcal{A}^2x_n = \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{H}z.$$

5.  $\mathcal{A}$ -compatible of type (E) [15] if

$$\lim_{n \rightarrow \infty} \mathcal{A}^2x_n = \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{H}z.$$

6.  $\mathcal{H}$ -compatible of type (E) [15] if

$$\lim_{n \rightarrow \infty} \mathcal{H}^2x_n = \lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{A}z.$$

Now we give an example of  $\mathcal{A}$ -subsequentially continuous and weakly subsequentially continuous mappings in the fuzzy metric space:

**Example 2.2.** Let  $X = [0, 2]$  and  $M(x, y, t) = \frac{t}{t+|x-y|}$  with continuous  $t$ -norm:  $a * b = ab, t > 0$ . We define  $\mathcal{A}$  and  $\mathcal{H}$  as:

$$\mathcal{A}x = \begin{cases} 1+x, & 0 \leq x \leq 1 \\ \frac{x+1}{2}, & 1 < x \leq 2, \end{cases} \quad \mathcal{H}x = \begin{cases} 1-x, & 0 \leq x \leq 1 \\ 2-x, & 1 < x \leq 2. \end{cases}$$

Observe that  $\mathcal{A}$  and  $\mathcal{H}$  are not continuous at 1.

We consider a sequence  $\{x_n\}$ , where  $x_n = \frac{1}{n}, n \geq 1$ .

So,

$$\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = 1$$

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \lim_{n \rightarrow \infty} \mathcal{A}(1 - \frac{1}{n}) = 2 = \mathcal{A}1,$$

i.e.,  $(\mathcal{A}, \mathcal{H})$  is  $\mathcal{A}$ -subsequentially continuous as well as weakly subsequentially continuous.

$\{y_n\}$  is a sequence, where  $y_n = 1 + \frac{1}{n}, n \geq 1$ , then

$$\lim_{n \rightarrow \infty} \mathcal{A}y_n = \lim_{n \rightarrow \infty} \mathcal{H}y_n = 1,$$

but

$$\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}y_n = \lim_{n \rightarrow \infty} \mathcal{H}(1 + \frac{1}{2n}) = 1 \neq \mathcal{H}1.$$

So,  $(\mathcal{A}, \mathcal{H})$  is not reciprocally continuous.

Observe that the weakly subsequentially continuous or  $\mathcal{A}$ -subsequentially continuous or  $\mathcal{H}$ -subsequentially continuous mappings are never reciprocally continuous maps [12] (see also Tomar and Karapinar [18]).

Now we give an example of compatible of type (E) in a fuzzy metric spaces:

**Example 2.3.** Let  $X = [0, \infty)$  and  $M(x, y, t) = \frac{t}{t+|x-y|}$  with the  $t$ -norm  $a * b = ab$ . We define  $\mathcal{A}$  and  $\mathcal{H}$  as follows:

$$\mathcal{A}x = \begin{cases} 2, & 0 \leq x \leq 2 \\ x+1, & x > 2, \end{cases} \quad \mathcal{H}x = \begin{cases} \frac{x+2}{2}, & 0 \leq x \leq 2 \\ 0, & x > 2. \end{cases}$$

If the sequence  $\{x_n\} \in X$  is defined by  $x_n = 2 - \frac{1}{n}$ , then

$$\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = 2,$$

$$\lim_{n \rightarrow \infty} \mathcal{A}^2x_n = \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = 2 = \mathcal{H}2,$$

$$\lim_{n \rightarrow \infty} \mathcal{H}^2x_n = \lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = 2 = \mathcal{A}2.$$

Hence,  $(\mathcal{A}, \mathcal{H})$  is compatible of type (E).

Observe that compatibility of type (E) implies  $\mathcal{A}$ -compatibility as well as  $\mathcal{H}$ -compatibility of type (E) but the reverse is not correct.

**Lemma 2.4.** [11] If in a fuzzy metric space  $(X, M, *)$ ,  $\exists$  a constant  $k \in (0, 1)$  satisfying

$$M(x, y, kt) \geq M(x, y, t),$$

$t > 0$  and fixed  $x, y \in X$ , then  $x = y$ .

### 3. Main results

Now we utilize the idea of weak subsequential continuity and compatibility of type (E) in a fuzzy metric space.

**Theorem 3.1.** Let  $(\mathcal{A}, \mathcal{H})$  and  $(\mathcal{B}, \mathcal{K})$  be compatible of type (E) as well as weakly subsequentially continuous pairs of a fuzzy metric space  $(X, M, *)$ . Then pairs  $(\mathcal{A}, \mathcal{H})$  and  $(\mathcal{B}, \mathcal{K})$  have a coincidence point. If:

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq \min \left\{ \begin{array}{l} M(\mathcal{A}x, \mathcal{B}y, t), M(\mathcal{A}x, \mathcal{H}x, t), \\ M(\mathcal{B}y, \mathcal{K}y, t), M(\mathcal{A}x, \mathcal{K}y, t), M(\mathcal{B}y, \mathcal{H}x, t) \end{array} \right\} \quad (3.1)$$

$k \in (0, 1)$ ,  $x, y \in X$  and  $t > 0$ , then  $\mathcal{A}, \mathcal{B}, \mathcal{H}$  and  $\mathcal{K}$  have a unique common fixed point in  $(X, M, *)$ .

**Proof .** As  $(\mathcal{A}, \mathcal{H})$  is weakly subsequentially continuous,  $\exists$  a sequence  $\{x_n\}$  in  $X$  such that

$$\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = z \in X$$

and

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{A}z$$

or

$$\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{H}z.$$

Also  $(\mathcal{A}, \mathcal{H})$  is compatible of type (E), So

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \lim_{n \rightarrow \infty} \mathcal{A}^2x_n = \mathcal{H}z$$

and

$$\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}^2x_n = \mathcal{A}z.$$

Consequently,  $\mathcal{A}z = \mathcal{H}z$ . Similarly  $(\mathcal{B}, \mathcal{K})$  is weakly subsequentially continuous  $\exists$  a sequence  $\{y_n\}$  such that

$$\lim_{n \rightarrow \infty} \mathcal{B}y_n = \lim_{n \rightarrow \infty} \mathcal{K}y_n = w \in X$$

and

$$\lim_{n \rightarrow \infty} \mathcal{B}\mathcal{K}y_n = \mathcal{B}w.$$

or

$$\lim_{n \rightarrow \infty} \mathcal{K}\mathcal{B}y_n = \mathcal{K}w.$$

Also  $(\mathcal{B}, \mathcal{K})$  is compatible of type (E)

$$\lim_{n \rightarrow \infty} \mathcal{B}\mathcal{K}y_n = \lim_{n \rightarrow \infty} \mathcal{B}^2y_n = \mathcal{K}w$$

$$\lim_{n \rightarrow \infty} \mathcal{K}\mathcal{B}y_n = \lim_{n \rightarrow \infty} \mathcal{K}^2y_n = \mathcal{B}w.$$

So,  $\mathcal{B}w = \mathcal{K}w$ .

We assert  $\mathcal{A}z = \mathcal{B}w$ . Using  $x = z$  and  $y = w$  in (3.1):

$$M(\mathcal{H}z, \mathcal{K}w, kt) \geq \min \left\{ \begin{array}{l} M(\mathcal{A}z, \mathcal{B}w, t), M(\mathcal{A}z, \mathcal{H}z, t), \\ M(\mathcal{B}w, \mathcal{K}w, t), M(\mathcal{A}z, \mathcal{K}w, t), M(\mathcal{B}w, \mathcal{H}z, t) \end{array} \right\}$$

$$M(\mathcal{A}z, \mathcal{B}w, kt) \geq \min\{M(\mathcal{A}z, \mathcal{B}w, t), 1, 1, M(\mathcal{A}z, \mathcal{B}w, t), M(\mathcal{A}z, \mathcal{B}w, t)\},$$

i.e.,

$$M(\mathcal{A}z, \mathcal{B}w, kt) \geq M(\mathcal{A}z, \mathcal{B}w, t).$$

From Lemma 2.4,  $\mathcal{A}z = \mathcal{B}w$ .

Now we prove  $z = \mathcal{A}z$ . Substituting  $x = x_n$  and  $y = w$  in (3.1):

$$M(\mathcal{H}x_n, \mathcal{K}w, kt) \geq \min \left\{ \begin{array}{l} M(\mathcal{A}x_n, \mathcal{B}w, t), M(\mathcal{A}x_n, \mathcal{H}x_n, t), \\ M(\mathcal{B}w, \mathcal{K}w, t), M(\mathcal{A}x_n, \mathcal{K}w, t), M(\mathcal{B}w, \mathcal{H}x_n, t) \end{array} \right\}.$$

Letting  $n \rightarrow \infty$ :

$$M(z, \mathcal{K}w, kt) \geq \min\{M(z, \mathcal{B}w, t), 1, 1, M(z, \mathcal{K}w, t), M(\mathcal{B}w, z, t)\},$$

i.e.,

$$M(z, \mathcal{A}z, kt) \geq M(z, \mathcal{A}z, t).$$

From Lemma 2.4,  $z = \mathcal{A}z = \mathcal{H}z$ .

Substituting  $x = x_n$  and  $y = y_n$  in (3.1):

$$M(\mathcal{H}x_n, \mathcal{K}y_n, kt) \geq \min \left\{ \begin{array}{l} M(\mathcal{A}x_n, \mathcal{B}y_n, t), M(\mathcal{A}x_n, \mathcal{H}x_n, t), \\ M(\mathcal{B}y_n, \mathcal{K}y_n, t), M(\mathcal{A}x_n, \mathcal{K}y_n, t), M(\mathcal{B}y_n, \mathcal{H}x_n, t) \end{array} \right\}.$$

Letting  $n \rightarrow \infty$ :

$$M(z, w, kt) \geq \min\{M(z, w, t), 1, 1, M(z, w, t), M(w, z, t)\},$$

i.e.,  $M(z, w, kt) \geq M(z, w, t)$ , i.e.,  $z = w$  (Lemma 2.4).

Hence,  $z$  is a common fixed point of  $\mathcal{A}, \mathcal{B}, \mathcal{H}$  and  $\mathcal{K}$ .

Suppose  $q$  is another common fixed point. Substituting  $x = z$  and  $y = q$  in (3.1):

$$M(\mathcal{H}z, \mathcal{K}q, kt) \geq \min\{M(\mathcal{A}z, \mathcal{B}q, t), M(\mathcal{A}z, \mathcal{H}q, t), M(\mathcal{B}q, \mathcal{K}q, t), M(\mathcal{A}z, \mathcal{K}q, t), M(\mathcal{B}q, \mathcal{H}z, t)\}$$

$$M(z, q, kt) \geq M(z, q, t).$$

Hence,  $z = q$ . Consequently,  $z$  is unique.  $\square$

**Example 3.2.** Let  $X = [0, 2]$  and  $M = \frac{t}{t+|x-y|}$  with  $t$ -norm defined by  $a * b = \min\{a, b\}$ ,  $x, y \in X$  and  $t > 0$ . Let

$$\mathcal{A}x = \begin{cases} x, & 0 \leq x \leq 1 \\ \frac{1}{2}, & 1 < x \leq 2, \end{cases} \quad \mathcal{B}x = \begin{cases} \frac{x+1}{2}, & 0 \leq x \leq 1 \\ 2, & 1 < x \leq 2, \end{cases}$$

$$\mathcal{H}x = \begin{cases} 1, & 0 \leq x \leq 1 \\ \frac{5}{4}, & 1 < x \leq 2, \end{cases} \quad \mathcal{K}x = \begin{cases} 2-x, & 0 \leq x \leq 1 \\ \frac{3}{4}, & 1 < x \leq 2. \end{cases}$$

Consider a sequence  $\{x_n\}$  defined as  $x_n = 1 - \frac{1}{n}$ ,  $n \geq 1$ .

Clearly,

$$\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = 1.$$

Also,

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{A}1$$

$$\lim_{n \rightarrow \infty} \mathcal{H}\mathcal{H}x_n = \lim_{n \rightarrow \infty} \mathcal{H}\mathcal{A}x_n = \mathcal{A}1$$

and

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{H}1,$$

Consider a sequence  $\{y_n\}$  defined by  $y_n = 1$ ,  $n \geq 1$ .

Clearly,

$$\lim_{n \rightarrow \infty} \mathcal{B}y_n = \lim_{n \rightarrow \infty} \mathcal{K}y_n = 1$$

and

$$\lim_{n \rightarrow \infty} \mathcal{B}\mathcal{K}y_n = \mathcal{B}1,$$

$$\lim_{n \rightarrow \infty} \mathcal{K}\mathcal{K}y_n = \lim_{n \rightarrow \infty} \mathcal{K}\mathcal{B}y_n = \mathcal{B}1$$

and

$$\lim_{n \rightarrow \infty} \mathcal{B}\mathcal{B}y_n = \lim_{n \rightarrow \infty} \mathcal{B}\mathcal{K}y_n = \mathcal{K}1,$$

i.e.,  $(\mathcal{A}, \mathcal{H})$  and  $(\mathcal{B}, \mathcal{K})$  are compatible of type (E) as well as weakly subsequentially continuous. Now,

1. When  $x, y \in [0, 1]$ :

$$M(\mathcal{H}x, \mathcal{K}y, t) = \frac{t}{t + |y - 1|} \geq \frac{t}{t + \frac{3}{2}|y - 1|} = M(\mathcal{B}y, \mathcal{K}y, t),$$

then for any  $k \in [\frac{2}{3}, 1)$ :

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq M(\mathcal{B}y, \mathcal{K}y, t).$$

2. When  $x \in [0, 1]$  and  $y \in (1, 2]$ :

$$M(\mathcal{H}x, \mathcal{K}y, t) = \frac{t}{t + \frac{1}{4}} \geq \frac{t}{t + 1} = M(\mathcal{B}y, \mathcal{H}x, t),$$

then for any  $k \in [\frac{1}{4}, 1)$ :

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq M(\mathcal{B}y, \mathcal{H}x, t).$$

3. When  $x \in (1, 2]$  and  $y \in [0, 1]$ :

$$M(\mathcal{H}x, \mathcal{K}y, t) = \frac{t}{t + |\frac{3}{4} - y|} \geq \frac{t}{t + |\frac{3}{2} - y|} = M(\mathcal{A}x, \mathcal{K}y, t),$$

then for any  $k \in [\frac{1}{2}, 1)$ :

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq M(\mathcal{A}x, \mathcal{K}y, t).$$

4. When  $x, y \in (1, 2]$ :

$$M(\mathcal{H}x, \mathcal{K}y, t) = \frac{t}{t + \frac{1}{2}} \geq \frac{t}{t + \frac{3}{2}} = M(\mathcal{A}x, \mathcal{B}y, t),$$

then for any  $k \in [\frac{1}{3}, 1)$ :

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq M(\mathcal{A}x, \mathcal{B}y, t).$$

Hence, for any  $k \in [\frac{2}{3}, 1)$  and for all  $x, y \in [0, \infty)$ , the inequality (3.1) holds.

Consequently, the hypotheses of Theorem 3.1 are verified and 1 is the unique common fixed point for  $\mathcal{A}, \mathcal{B}, \mathcal{H}$  and  $\mathcal{K}$ . It is interesting to observe that none of the mappings is continuous. Moreover, neither  $\mathcal{A}X \subseteq \mathcal{B}X$  nor  $\mathcal{H}X \subseteq \mathcal{K}X$ .

If  $\mathcal{A} = \mathcal{B}$  and  $\mathcal{H} = \mathcal{K}$ :

**Corollary 3.3.** Let  $(\mathcal{A}, \mathcal{H})$  be compatible of type (E) and weakly subsequentially continuous pair of a fuzzy metric space  $(X, M, *)$ . Then the pair  $(\mathcal{A}, \mathcal{H})$  has a coincidence point. If:

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq \min \left\{ \begin{array}{l} M(\mathcal{A}x, \mathcal{A}y, t), M(\mathcal{A}x, \mathcal{H}x, t), \\ M(\mathcal{A}y, \mathcal{H}y, t), M(\mathcal{A}x, \mathcal{H}y, t), M(\mathcal{A}y, \mathcal{H}x, t) \end{array} \right\} \tag{3.2}$$

$x, y \in X, k \in (0, 1)$  and  $t > 0$ , then  $\mathcal{A}$  and  $\mathcal{H}$  have a unique common fixed point.

**Corollary 3.4.** Corollary 3.3 is true even if the pair  $(\mathcal{A}, \mathcal{H})$  is  $\mathcal{A}$ -compatible of type (E) and  $\mathcal{A}$ -subsequentially continuous.

**Example 3.5.** Let  $X = [0, \infty)$  and  $M = \frac{t}{t+|x-y|}$  with  $t$ -norm,  $a * b = \min\{a, b\}$ ,  $x, y \in X$ ,  $t > 0$ . Define:

$$\mathcal{A}x = \begin{cases} 3x, & 0 \leq x \leq 1 \\ 2x - 1, & x > 1, \end{cases} \quad \mathcal{H}x = \begin{cases} \frac{x}{4}, & 0 \leq x \leq 1 \\ 0, & x > 1. \end{cases}$$

Consider a sequence  $\{x_n\}$ , where  $x_n = \frac{1}{n}$ ,  $n \geq 1$ .

Clearly,

$$\lim_{n \rightarrow \infty} \mathcal{A}x_n = \lim_{n \rightarrow \infty} \mathcal{H}x_n = 0.$$

Also we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n &= \mathcal{A}0 \\ \lim_{n \rightarrow \infty} \mathcal{A}^2x_n &= \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}x_n = \mathcal{H}0, \end{aligned}$$

i.e., the pair  $(\mathcal{A}, \mathcal{H})$  is  $\mathcal{A}$ -compatible of type (E) as well as  $\mathcal{A}$ -subsequentially continuous. Now,

1. When  $x, y \in [0, 1]$ :

$$M(\mathcal{H}x, \mathcal{H}y, t) = \frac{t}{t + \frac{1}{4}|x-y|} \geq \frac{t}{t + 3|x-y|} = M(\mathcal{A}x, \mathcal{A}y, t),$$

then for any  $k \in [\frac{1}{12}, 1)$ :

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq M(\mathcal{A}x, \mathcal{A}y, t).$$

2. When  $x \in [0, 1]$  and  $y > 1$ :

$$M(\mathcal{H}x, \mathcal{H}y, t) = \frac{t}{t + \frac{1}{4}x} \geq \frac{t}{t + \frac{11}{4}x} = M(\mathcal{A}x, \mathcal{H}x, t),$$

then for any  $k \in [\frac{1}{11}, 1)$ :

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq M(\mathcal{A}x, \mathcal{H}x, t).$$

3. When  $x \in (1, \infty)$  and  $y \in [0, 1]$ :

$$M(\mathcal{H}x, \mathcal{H}y, t) = \frac{t}{t + \frac{1}{4}y} \geq \frac{t}{t + \frac{11}{4}y} = M(\mathcal{A}y, \mathcal{H}y, t),$$

then for any  $k \in [\frac{1}{11}, 1)$ :

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq M(\mathcal{A}y, \mathcal{H}y, t).$$

4. When  $x, y \in (1, \infty)$ :

$$M(\mathcal{H}x, \mathcal{H}y, t) = \frac{t}{t+0} \geq \frac{t}{t+2|x-y|} = M(\mathcal{A}x, \mathcal{A}y, t),$$

then for any  $k \in (0, 1)$ :

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq M(\mathcal{A}x, \mathcal{A}y, t).$$

Hence, for any  $k \in [\frac{1}{11}, 1)$  the inequality ?? holds and as a result all the hypotheses of Corollary 3.4 are verified and 0 is the unique common fixed point for  $\mathcal{A}$  and  $\mathcal{H}$ . It is interesting to observe that both  $\mathcal{A}$  and  $\mathcal{H}$  are discontinuous and neither  $\mathcal{A}X \subseteq \mathcal{H}X$  nor  $\mathcal{H}X \subseteq \mathcal{A}X$ .

If  $\mathcal{A} = \mathcal{B}$  in Theorem 3.1:

**Corollary 3.6.** *Let  $(\mathcal{A}, \mathcal{H})$  and  $(\mathcal{A}, \mathcal{K})$  be compatible of type (E) and weakly subsequentially continuous pairs of a fuzzy metric space  $(X, M, *)$ . Then  $\mathcal{A}, \mathcal{H}$  and  $\mathcal{K}$  have a coincidence point. If*

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq \min \left\{ \begin{matrix} M(\mathcal{A}x, \mathcal{A}y, t), M(\mathcal{A}x, \mathcal{H}x, t), \\ M(\mathcal{A}y, \mathcal{K}y, t), M(\mathcal{A}x, \mathcal{K}y, t), M(\mathcal{A}y, \mathcal{H}x, t) \end{matrix} \right\} \tag{3.3}$$

*$x, y \in X$  and  $t > 0$ , then  $\mathcal{A}, \mathcal{H}$  and  $\mathcal{K}$  have a unique common fixed point.*

If  $\mathcal{H} = \mathcal{K}$ :

**Corollary 3.7.** *Let  $(\mathcal{A}, \mathcal{H})$  be compatible of type (E) and weakly subsequentially continuous pairs of a fuzzy metric space  $(X, M, *)$ . Then  $\mathcal{A}$  and  $\mathcal{H}$  have a coincidence point.*

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq \min \left\{ \begin{matrix} M(\mathcal{A}x, \mathcal{A}y, t), M(\mathcal{A}x, \mathcal{H}x, t), \\ M(\mathcal{A}y, \mathcal{H}y, t), M(\mathcal{A}x, \mathcal{H}y, t), M(\mathcal{A}y, \mathcal{H}x, t)t \end{matrix} \right\} \tag{3.4}$$

*$x, y \in X, k \in (0, 1)$  and  $t > 0$ , then  $\mathcal{A}$  and  $\mathcal{H}$  have a unique common fixed point.*

Now we utilize the weak subsequential continuity and compatibility of type (E) for  $\phi$ -contractive type mapping.

**Theorem 3.8.** *Theorem 3.1 remains true even if we replace 3.1 by*

$$M(\mathcal{H}x, \mathcal{K}y, kt) \geq \phi \left( \min \left\{ \begin{matrix} M(\mathcal{A}x, \mathcal{B}y, t), M(\mathcal{A}x, \mathcal{H}x, t), \\ M(\mathcal{B}y, \mathcal{K}y, t), M(\mathcal{A}x, \mathcal{K}y, t), M(\mathcal{B}y, \mathcal{H}x, t) \end{matrix} \right\} \right) \tag{3.5}$$

*$x, y \in X, k \in (0, 1)$  and  $t > 0$ ,  $\phi : [0, 1] \rightarrow [0, 1]$  is a lower semi continuous such that  $\phi(t) > t$ ,  $t \in (0, 1)$  with  $\phi(0) = 0$  and  $\phi(1) = 1$ .*

**Proof .** Theorem 3.8 follow the pattern of Theorem 3.1 as  $t > 0$ ,  $\phi(t) > t$ . $\square$  Now we exploit integral type contractive condition in a fuzzy metric space.

**Theorem 3.9.** *Theorem 3.1 remains true even if we replace 3.1 by*

$$\int_0^{M(\mathcal{H}x, \mathcal{K}y, kt)} \varphi(t)dt \geq \int_0^{m(x, y)} \varphi(t)dt, \tag{3.6}$$

where

$$m(x, y) = \min\{M(\mathcal{A}x, \mathcal{B}y, t), M(\mathcal{A}x, \mathcal{H}x, t), M(\mathcal{B}y, \mathcal{K}y, t), M(\mathcal{A}x, \mathcal{K}y, t), M(\mathcal{B}y, \mathcal{H}x, t)\}$$

*and  $x, y \in X, k \in (0, 1)$  and  $t > 0$ ,  $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is a Lebesgue integrable and summable and for each  $\varepsilon > 0$ ,  $\int_0^\varepsilon \varphi(t)dt > 0$ .*

**Proof .** Following Theorem 3.1,  $\mathcal{A}z = \mathcal{B}z$  and  $\mathcal{B}w = \mathcal{K}w$ .

We prove  $\mathcal{A}z = \mathcal{B}w$ . Taking  $x = z$  and  $y = w$  in (3.6) we get:

$$\int_0^{M(\mathcal{H}z, \mathcal{K}w, kt)} \varphi(t)dt \geq \int_0^{\min\{M(\mathcal{A}z, \mathcal{B}w, t), M(\mathcal{A}z, \mathcal{H}z, t), M(\mathcal{B}w, \mathcal{K}w, t), M(\mathcal{A}z, \mathcal{K}w, t), M(\mathcal{B}w, \mathcal{H}z, t)\}} \varphi(t)dt.$$



Since,  $\mathcal{A}z = \mathcal{H}z$  and  $\mathcal{B}w = \mathcal{K}w$ , we get:

$$\begin{aligned} \int_0^{M(\mathcal{A}z, \mathcal{B}w, kt)} \varphi(t) dt &\geq \int_0^{\min\{M(\mathcal{A}z, \mathcal{B}w, t), 1, 1, M(\mathcal{A}z, \mathcal{B}w, t), M(\mathcal{A}z, \mathcal{B}w, t)\}} \varphi(t) dt \\ &= \int_0^{M(\mathcal{A}z, \mathcal{B}w, t)} \varphi(t) dt, \end{aligned}$$

i.e.,

$$M(\mathcal{A}z, \mathcal{B}w, kt) \geq M(\mathcal{A}z, \mathcal{B}w, t).$$

From Lemma 2.4,  $\mathcal{A}z = \mathcal{B}w$ .

Substituting  $x = x_n$  and  $y = w$  in (3.6):

$$\int_0^{M(\mathcal{H}x_n, \mathcal{K}w, kt)} \varphi(t) dt \geq \int_0^{m(x_n, w)} \varphi(t) dt.$$

Letting  $n \rightarrow \infty$ :

$$\int_0^{M(z, \mathcal{K}w, kt)} \varphi(t) dt \geq \int_0^{M(z, \mathcal{B}w, t)} \varphi(s) ds,$$

i.e.,

$$\int_0^{M(z, \mathcal{A}z, kt)} \varphi(t) dt \geq \int_0^{M(z, \mathcal{A}z, t)} \varphi(t) dt,$$

i.e.,

$$M(z, \mathcal{A}z, kt) \geq M(z, \mathcal{A}z, t),$$

i.e.,  $z = \mathcal{A}z = \mathcal{H}z$  (from Lemma 2.4).

Nextly, taking  $x = x_n$  and  $y = w$  in (3.6):

$$\int_0^{M(\mathcal{H}x_n, \mathcal{K}y_n, kt)} \varphi(t) dt \geq \int_0^{m(x_n, y_n)} \varphi(t) dt.$$

Passing the limit, when  $n \rightarrow \infty$ :

$$\int_0^{M(z, w, kt)} \varphi(t) dt \geq \int_0^{m(z, w)} \varphi(t) dt = \int_0^{M(z, w, t)} \varphi(t) dt,$$

i.e.,

$$M(z, w, kt) \geq M(z, w, t).$$

Consequently,  $z$  is a common fixed point of  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{H}$  and  $\mathcal{K}$ .

If  $q$  is another fixed point, then using (3.6):

$$\int_0^{M(\mathcal{H}z, \mathcal{K}q, kt)} \varphi(t) dt \geq \int_0^{m(z, q)} \varphi(t) dt = \int_0^{M(z, q, t)} \varphi(t) dt.$$

Hence,  $z = q$  and  $z$  is unique.  $\square$

**Remark 3.10.** All results are true even if we replace compatibility of type (E) and weak subsequential continuity by any one of the following:

1.  $\mathcal{H}$  (or  $\mathcal{A}$ )-compatibility of type (E) and  $\mathcal{H}$  (or  $\mathcal{A}$ )-subsequential continuity,
2.  $\mathcal{A}$  (or  $\mathcal{H}$ )-compatibility of type (E) or compatibility of type (E) and subsequential continuity.

### 4. Application

We utilize Corollary 3.4 to solve functional equations arising in dynamic programming as an application. It was first studied by Bellman [1] using famous Banach fixed point theorem. Let  $W \subset X$  be state space and  $D \subset Y$  be decision space. Let  $B(W)$  be the set of bounded functions on  $W$ . Define:

$$M(h, k, t) = e^{-\frac{d(h,k)}{t}},$$

with t-norm  $a * b = \min\{a, b\}$ ,  $a, b \in [0, 1]$ , where  $d(h, k) = \|h(\tau) - k(\tau)\|_\infty = \sup_{\tau \in W} |h - k|_\tau$ . Then  $(X, M, *)$  is a fuzzy metric space.

**Theorem 4.1.** *Let  $\mathcal{H}$  and  $\mathcal{A}$  be self mappings of  $(X, M, *)$ . If the following hypotheses hold:*

- (a)  $H$  and  $K$  are bounded,
- (b)  $\exists \delta \in (0, 1)$  such that:

$$|H(x, y, f(\tau(x, y))) - K(x, y, g(\tau(x, y)))| \leq \delta(h, k, t),$$

where,

$$\delta(h, k, t) = \min\{M(\mathcal{A}x, \mathcal{A}y, t), M(\mathcal{A}x, \mathcal{H}x, t), M(\mathcal{A}y, \mathcal{H}y, t), M(\mathcal{A}x, \mathcal{H}y, t), M(\mathcal{A}y, \mathcal{H}x, t)\},$$

$x, y \in W$  and  $h, k \in B(W)$ ,

- (c)  $\exists$  a sequence  $\{h_n\} \in W$ , satisfying

$$\lim_{n \rightarrow \infty} \mathcal{A}h_n = \lim_{n \rightarrow \infty} \mathcal{H}h_n = h \in B(W),$$

$$\lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}h_n = \mathcal{A}h$$

and

$$\lim_{n \rightarrow \infty} \mathcal{A}^2h_n = \lim_{n \rightarrow \infty} \mathcal{A}\mathcal{H}h_n = \mathcal{H}h.$$

Then the system

$$\begin{cases} \mathcal{H}f(t) = \sup_{x \in W} \{u(x, t) + H(x, y, f(\tau(x, y)))\} \\ \mathcal{A}g(t) = \sup_{x \in W} \{u(x, y) + K(x, y, g(\tau(x, y)))\}, \end{cases} \tag{4.1}$$

has a unique bounded solution.

**Proof .** The system has a unique solution iff  $\mathcal{H}$  and  $\mathcal{A}$  have a unique common fixed point. For all  $h, k \in B(W)$  and  $\varepsilon > 0$ ,  $\exists y, z \in W$  such that

$$\mathcal{H}h < u(x, y) + H(x, y, h(\tau(x, y))) + \varepsilon \tag{4.2}$$

$$\mathcal{A}k < u(x, z) + K(x, z, k(\tau(x, z))) + \varepsilon \tag{4.3}$$

and since,

$$\mathcal{H}h \geq u(x, z) + H(x, z, h(\tau(x, z))), \tag{4.4}$$

$$\mathcal{A}k \geq u(x, y) + K(x, y, k(\tau(x, y))), \tag{4.5}$$

then from (4.2) and (4.5)

$$\begin{aligned} \mathcal{H}h - \mathcal{A}k &\leq H(x, y, h(\tau(x, y))) - K(x, y, k(\tau(x, y))) + \varepsilon \\ &\leq \delta(h, k) + \varepsilon. \end{aligned} \quad (4.6)$$

Also from (4.3) and (4.4)

$$\begin{aligned} \mathcal{H}h - \mathcal{A}k &> H(x, y, h(\tau(x, y))) - K(x, y, k(\tau(x, y))) - \varepsilon \\ &\geq -\delta d(h, k) - \varepsilon. \end{aligned} \quad (4.7)$$

Consequently, inequalities (4.5) and (4.7) implies that

$$\begin{aligned} d(\mathcal{H}h, \mathcal{A}k) &= \sup |\mathcal{H}h - \mathcal{A}k| \leq |H(x, y, h(\tau(x, y))) - K(x, y, k(\tau(x, y)))| + \varepsilon \\ &\leq \delta(h, k) + \varepsilon. \end{aligned}$$

Since,  $\varepsilon > 0$  is arbitrary

$$d(\mathcal{H}h, \mathcal{A}k) \leq d(h, k). \quad (4.8)$$

So,

$$e^{-\frac{d(\mathcal{H}h, \mathcal{A}k)}{t}} \geq e^{-\frac{d(h, k)}{t}}.$$

$\Rightarrow$

$$M(\mathcal{H}x, \mathcal{H}y, kt) \geq \delta(h, k, t).$$

The condition (c) implies that  $(\mathcal{A}, \mathcal{H})$  is  $\mathcal{A}$ -compatible of type (E) as well as  $\mathcal{A}$ -subsequentially continuous as a result all the hypotheses of Corollary 3.4 are verified and consequently,  $\mathcal{H}$  and  $\mathcal{A}$  have a unique common fixed point. Hence, (4.1) has a unique solution.  $\square$

## References

- [1] R. Bellman and E. S. Lee, *Functional equations in dynamic programming*, Aequ. Math. 17 (1978) 1–18.
- [2] S. Beloul, *Common fixed point theorems for weakly subsequentially continuous generalized contractions with applications*, Appl. Maths. E-Notes, 15 (2015) 173–186.
- [3] S. Beloul, *Common fixed point theorems for weakly subsequentially continuous mappings in fuzzy metric spaces via implicit relation*, TWMS J. App. Eng. Math. 8(1) (2018) 284–294.
- [4] S. Beloul and A. Tomar, *Integral type common fixed point theorems in modified intuitionistic fuzzy metric spaces*, Afrika Mat. 3 (2019) 1–16.
- [5] Y. J. Cho, S. Sedghi and N. Shobe, *Generalized fixed point theorems for Compatible mappings with some types in fuzzy metric spaces*, Chaos, Solitons Fract. 39 (2009) 2233–2244.
- [6] A. George and P. Veeramani, *On some results of analysis for fuzzy metric spaces*, Fuzzy Sets and Systems, 90 (1997) 365–368.
- [7] I. Kramosil and J. Michalek, *Fuzzy metric and statistical metric spaces*, Kybernetika, 11 (1975) 336–344.
- [8] S. Manro, A. Tomar and B. E. Rhoades, *Coincidence and common fixed point theorems in fuzzy metric spaces using a Meir-Keeler Type contractive condition*, Gazi Univ. J. Sci. 27(1) (2014) 669–678.
- [9] S. Manro and A. Tomar, *Common fixed point theorems using property (E.A.) and its variants involving quadratic terms*, Ann. Fuzzy Math. Inf. 7(3) (2014) 473–484.
- [10] S. Manro and A. Tomar, *Faintly compatible maps and existence of common fixed points in fuzzy metric space*, Ann. Fuzzy Math. Inf. 8(2) (2014) 223–230.
- [11] S. N. Mishra, N. Sharma and S. L. Singh, *Common fixed point of maps on fuzzy metric spaces*, Internat. J. Math. Math. Sci. 17 (1994) 253–258.

- [12] R. P. Pant, *A common fixed point theorem under a new condition*, Indian. J. Pure Appl. Math. 30(2) (1999) 147–152.
- [13] R. Rana, R. C. Dimri and A. Tomar, *Fixed point theorems in Fuzzy Metric spaces using implicit relations*, Int. J. Comp. Appl. 8(1) 2010 16–21.
- [14] D. O'Regan and M. Abbas, *Necessary and sufficient conditions for common fixed point theorems in fuzzy metric space*, Demonstr. Math. 42(4) (2009).
- [15] M. R. Singh and Y. M. Singh, *Compatible mappings of type (E) and common fixed point theorems of Meir-Keeler type*, Int. J. Math. Sci. Engg. Appl. 1(2) (2007) 299–315.
- [16] S. L. Singh and A. Tomar, *Weaker forms of commuting maps and existence of fixed points*, J. Korea. Soc. Math. Educ. ser. B: Pure Appl. Math. 10(3) (2003) 145–161.
- [17] S. L. Singh and A. Tomar, *Fixed point theorems in FM-spaces*, J. Fuzzy Math. 12(4) (2004).
- [18] A. Tomar and E. Karapinar, *On variants of continuity and existence of fixed point via Meir-Keeler contractions in MC-spaces*, J. Adv. Math. Stud. 9(2) (2016) 348–359.
- [19] A. Tomar, S. Upadhyay and R. Sharma, *Existence of common fixed point in symmetric space with application*, Elect. J. Math. Anal. Appl. 7(1) (2019) 116–129.
- [20] L. A. Zadeh, *Fuzzy sets*, Inf. Cont. 89 (1965) 338–353.