

# Approximation properties of bivariate generalized Baskakov-Kantorovich operators

Meenu Goyal<sup>a,\*</sup>, P. N. Agrawal<sup>b</sup>

<sup>a</sup>*School of Mathematics, Thapar Institute of Engineering and Technology, Patiala-147004, India*

<sup>b</sup>*Department of Mathematics, Indian Institute of Technology Roorkee-247667, India*

(Communicated by Madjid Eshaghi Gordji)

---

## Abstract

The purpose of this paper is to study the bivariate extension of the generalized Baskakov-Kantorovich operators and obtain results on the degree of approximation, Voronovskaja type theorems and their first order derivatives in polynomial weighted spaces. Furthermore, we illustrate the convergence of the bivariate operators to a certain function through graphics using Matlab algorithm. We also discuss the comparison of the convergence of the bivariate generalized Baskakov Kantorovich operators and the bivariate Szász-Kantorovich operators to the function through illustrations using Matlab.

Keywords: Rate of convergence, simultaneous approximation, bivariate modulus of continuity  
2020 MSC: 41A25, 41A28

---

## 1 Introduction

The approximation of functions by linear positive operators is an interesting area of research in approximation theory. Many new sequences and classes of linear positive operators have been considered in the literature and extensively studied by researchers. After the integral modifications of Bernstein polynomials by Kantorovich and Durrmeyer, the Kantorovich and Durrmeyer variants of several sequences of linear positive operators were defined and studied.

In [11], Erençin defined the Durrmeyer type modification of generalized Baskakov operators introduced by Mihešan [15], as

$$L_n^a(f; x) = \sum_{k=0}^{\infty} W_{n,k}^a(x) \frac{1}{B(k+1, n)} \int_0^{\infty} \frac{t^k}{(1+t)^{n+k+1}} f(t) dt, \quad x \geq 0,$$

where

$$W_{n,k}^a(x) = e^{\frac{-ax}{1+x}} \frac{P_k(n,a)}{k!} \frac{x^k}{(1+x)^{n+k}}, \quad P_k(n,a) = \sum_{i=0}^k \binom{n}{k} (n)_i a^{k-i}, \quad \text{and } (n)_0 = 1, (n)_i = n(n+1)\dots(n+i-1) \text{ for } i \geq 1 \text{ and}$$

---

\*Corresponding author

Email addresses: [meenu\\_rani@thapar.edu](mailto:meenu_rani@thapar.edu) (Meenu Goyal), [pnappfma@gmail.com](mailto:pnappfma@gmail.com) (P. N. Agrawal)

discussed some approximation properties. Very recently, Agrawal et al. [4] studied the simultaneous approximation and approximation of functions having derivatives of bounded variation by these operators.

Inspired by the above work, in [3], we proposed the Kantorovich modification of generalized Baskakov operators for the function  $f$  defined on  $C_\gamma[0, \infty) := \{f \in C[0, \infty) : f(t) = O(t^\gamma) \text{ as } t \rightarrow \infty, \text{ for some } \gamma > 0\}$  as follows :

$$K_n^a(f; x) = (n + 1) \sum_{k=0}^\infty W_{n,k}^a(x) \int_{\frac{k}{n+1}}^{\frac{k+1}{n+1}} f(t) dt, \quad a \geq 0. \tag{1.1}$$

In the present paper, our aim is to consider the bivariate extension of these operators. In this direction, Stancu [20] first introduced new linear positive operators in two and several dimensional variables. After that Barbosu [8] studied the bivariate extension of Stancu generalization of  $q$ -Bernstein operators. Dođru and Gupta [10] constructed a bivariate generalization of the Meyer-Konig and Zeller operators based on  $q$ -integers while in [2] Agratini presented two-dimensional extension of some univariate positive approximation processes expressed by series.

Many papers were published on approximation by modified Szász-Mirakyan and Baskakov operators for functions of one or two variables cf. [5, 6, 12, 13, 18, 19, 25] which deal with convergence, degree of approximation and Voronovskaja type theorems as well as convergence of partial derivatives of these operators. Wafi and Khatoon [21] defined the generalized Baskakov operators for functions of two variables in polynomial and exponential weighted spaces and discussed the rate of convergence and direct results. Later, in [22], the convergence of first order derivatives of these operators and a Voronovskaja type theorem were studied. For the recent background knowledge in this direction, readers can refer [1, 14, 16, 17, 23, 24].

## 2 Construction of the operators

Let  $I = [0, \infty) \times [0, \infty)$  and  $\omega_\gamma(x) = (1 + x^\gamma)^{-1}$  for  $\gamma \in \mathbb{N}^0$  (set of all non-negative integers). Further, for fixed  $\gamma_1, \gamma_2 \in \mathbb{N}^0$ , let  $\omega_{\gamma_1, \gamma_2}(x, y) = \omega_{\gamma_1}(x)\omega_{\gamma_2}(y)$ .

Then, for  $f \in C_{\gamma_1, \gamma_2}(I) := \{f \in C(I) : \omega_{\gamma_1, \gamma_2}(x, y)f(x, y) \text{ is bounded and uniformly continuous on } I\}$ , we define a bivariate extension of the operators (1.1) as follows:

$$K_{n_1, n_2}^a(f; x, y)$$

$$= (n_1 + 1)(n_2 + 1) \sum_{k_1=0}^\infty \sum_{k_2=0}^\infty W_{n_1, n_2, k_1, k_2}^a(x, y) \int_{\frac{k_2}{n_2+1}}^{\frac{k_2+1}{n_2+1}} \int_{\frac{k_1}{n_1+1}}^{\frac{k_1+1}{n_1+1}} f(u, v) du dv, \tag{2.1}$$

where

$$W_{n_1, n_2, k_1, k_2}^a(x, y) = \frac{x^{k_1} y^{k_2} p_{k_1}(n_1, a) p_{k_2}(n_2, a) e^{-\frac{ax}{1+x}} e^{-\frac{ay}{1+y}}}{k_1! k_2! (1+x)^{n_1+k_1} (1+y)^{n_2+k_2}}.$$

If  $f \in C_{\gamma_1, \gamma_2}(I)$  and if  $f(x, y) = f_1(x)f_2(y)$  for all  $(x, y) \in I$ , then

$$K_{n_1, n_2}^a(f(u, v); x, y) = K_{n_1}^a(f_1(u); x) K_{n_2}^a(f_2(v); y), \tag{2.2}$$

for  $(x, y) \in I$  and  $n_1, n_2 \in \mathbb{N}$ . The sup norm on  $C_{\gamma_1, \gamma_2}(I)$  is given by

$$\|f\|_{\gamma_1, \gamma_2} = \sup_{(x, y) \in I} |f(x, y)| \omega_{\gamma_1, \gamma_2}(x, y), \quad f \in C_{\gamma_1, \gamma_2}(I). \tag{2.3}$$

For  $f \in C_{\gamma_1, \gamma_2}(I)$ , we define the modulus of continuity

$$\omega(f, C_{\gamma_1, \gamma_2}; t, s) := \sup_{0 < h < t, 0 < \delta < s} \|\Delta_{h, \delta} f(\cdot, \cdot)\|_{\gamma_1, \gamma_2}, \quad t, s \geq 0, \tag{2.4}$$

where  $\Delta_{h, \delta} f(x, y) := f(x+h, y+\delta) - f(x, y)$  for  $(x, y) \in I$  and  $h, \delta > 0$ . Moreover, for fixed  $m \in \mathbb{N}$ , let  $C_{\gamma_1, \gamma_2}^m(I)$  be the space of all functions  $f \in C_{\gamma_1, \gamma_2}(I)$  having the partial derivatives  $\frac{\partial^k f}{\partial x^s \partial y^{k-s}} \in C_{\gamma_1, \gamma_2}(I), s = 1, 2, \dots, k; k = 1, 2, \dots, m$ .

The paper is organized as follows:

In Section 3 of this paper, we give some definitions and auxiliary results. In Section 4, we prove the main results of this paper wherein we study the degree of approximation, Voronovskaja type theorems and simultaneous approximation of first order derivatives for bivariate Baskakov Kantorovich operators  $K_{n_1, n_2}^a$ . The section 5 is devoted to the illustrations of the convergence of the operators  $K_{n_1, n_2}^a$  to a certain function and the comparison of convergence with bivariate Szász Kantorovich operators to the function using Matlab.

### 3 Auxiliary Results

**Lemma 3.1.** In [3], for the  $r$ th order central moment of  $K_n^a$ , defined as

$$u_{n,r}^a(x) := K_n^a((t-x)^r; x), \text{ we have}$$

- (i)  $u_{n,0}^a(x) = 1, u_{n,1}^a(x) = \frac{1}{n+1} \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right)$   
 and  $u_{n,2}^a(x) = \frac{1}{(n+1)^2} \left\{ (n+1)x^2 + (n-1)x + \frac{a^2x^2}{(1+x)^2} + 2ax \left( \frac{1-x}{1+x} \right) + \frac{1}{3} \right\};$
- (ii)  $u_{n,r}^a(x)$  is a rational function of  $x$  depending on the parameters  $a$  and  $r$ ;
- (iii) for each  $x \in (0, \infty), u_{n,r}^a(x) = O\left(\frac{1}{n^{\lfloor \frac{r+1}{2} \rfloor}}\right)$ , where  $\lfloor \beta \rfloor$  denotes the integer part of  $\beta$ .

**Lemma 3.2.** Let  $e_{ij} : I \rightarrow I, e_{ij} = x^i y^j, 0 \leq i, j \leq 2$  be two-dimensional test functions. Then the bivariate operators defined in (2.1) satisfy the following results:

- (i)  $K_{n_1, n_2}^a(e_{00}; x, y) = 1;$
- (ii)  $K_{n_1, n_2}^a(e_{10}; x, y) = \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right);$
- (iii)  $K_{n_1, n_2}^a(e_{01}; x, y) = \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right);$
- (iv)  $K_{n_1, n_2}^a(e_{20}; x, y) = \frac{1}{(n_1+1)^2} \left( n_1^2x^2 + n_1x^2 + 2n_1x + \frac{2an_1x^2}{1+x} + \frac{a^2x^2}{(1+x)^2} + \frac{2ax}{1+x} + \frac{1}{3} \right);$
- (v)  $K_{n_1, n_2}^a(e_{02}; x, y) = \frac{1}{(n_2+1)^2} \left( n_2^2y^2 + n_2y^2 + 2n_2y + \frac{2an_2y^2}{1+y} + \frac{a^2y^2}{(1+y)^2} + \frac{2ay}{1+y} + \frac{1}{3} \right);$
- (vi)  $K_{n_1, n_2}^a(e_{30}; x, y) = \frac{1}{(n_1+1)^3} \left( n_1^3x^3 + \frac{3n_1^2x^2}{2}(3+2x) + \frac{3n_1x^2}{2} + \frac{n_1x}{2}(4x^2+6x+5) + \frac{3ax^3n_1^2}{1+x} \right.$   
 $\left. + \frac{3an_1x^2}{1+x} \left\{ (3+x) + \frac{ax}{1+x} \right\} + \frac{ax}{1+x} \left\{ \frac{7}{2} + \frac{7}{2} \frac{ax}{1+x} + \frac{a^2x^2}{(1+x)^2} \right\} + \frac{1}{4} \right);$
- (vii)  $K_{n_1, n_2}^a(e_{03}; x, y) = \frac{1}{(n_2+1)^3} \left( n_2^3y^3 + \frac{3n_2^2y^2}{2}(3+2y) + \frac{3n_2y^2}{2} + \frac{n_2y}{2}(4y^2+6y+5) + \frac{3ay^3n_2^2}{1+y} \right.$   
 $\left. + \frac{3an_2y^2}{1+y} \left\{ (3+y) + \frac{ay}{1+y} \right\} + \frac{ay}{1+y} \left\{ \frac{7}{2} + \frac{7}{2} \frac{ay}{1+y} + \frac{a^2y^2}{(1+y)^2} \right\} + \frac{1}{4} \right).$

**Lemma 3.3.** For  $n_1, n_2 \in \mathbb{N}$ , we have

- (i)  $K_{n_1, n_2}^a(u-x; x, y) = \frac{1}{n_1+1} \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right);$
- (ii)  $K_{n_1, n_2}^a(v-y; x, y) = \frac{1}{n_2+1} \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right);$
- (iii)  $K_{n_1, n_2}^a((u-x)^2; x, y) = \frac{1}{(n_1+1)^2} \left( (n_1+1)x^2 + (n_1-1)x + \frac{a^2x^2}{(1+x)^2} + 2ax \left( \frac{1-x}{1+x} \right) + \frac{1}{3} \right);$
- (iv)  $K_{n_1, n_2}^a((v-y)^2; x, y) = \frac{1}{(n_2+1)^2} \left( (n_2+1)y^2 + (n_2-1)y + \frac{a^2y^2}{(1+y)^2} + 2ay \left( \frac{1-y}{1+y} \right) + \frac{1}{3} \right).$

**Remark 3.4.** For every  $x \in [0, \infty)$  and  $n \in \mathbb{N}$ , we have

$$K_{n_1, n_2}^a((u-x)^2; x, y) \leq \frac{\{\delta_{n_1}^a(x)\}^2}{n_1 + 1},$$

where  $\{\delta_{n_1}^a(x)\}^2 = \phi^2(x) + \frac{(1+a)^2}{n_1+1}$  and  $\phi(x) = \sqrt{x(1+x)}$ .

**Proof .** From Lemma 3.3 (iii), we have

$$K_{n_1, n_2}^a((u-x)^2; x, y) \leq \frac{1}{n_1 + 1} \left( x(1+x) + \frac{(1+a)^2}{n_1 + 1} \right) = \frac{\{\delta_{n_1}^a(x)\}^2}{n_1 + 1}.$$

□

**Lemma 3.5.** For every  $\gamma_1 \in \mathbb{N}^0$  there exist positive constants  $M_k(\gamma_1), k = 1, 2$  such that

- (i)  $\omega_{\gamma_1}(x) K_n^a\left(\frac{1}{\omega_{\gamma_1}(t)}; x\right) \leq M_1(\gamma_1),$
- (ii)  $\omega_{\gamma_1}(x) K_n^a\left(\frac{(t-x)^2}{\omega_{\gamma_1}(t)}; x\right) \leq M_2(\gamma_1) \frac{\{\delta_n^a(x)\}^2}{n+1},$

for all  $x \in \mathbb{R}^0 = \mathbb{R}_+ \cup \{0\}, \mathbb{R}_+ = (0, \infty)$  and  $n \in \mathbb{N}$ .

**Lemma 3.6.** For every  $\gamma_1, \gamma_2 \in \mathbb{N}^0$  there exist positive constant  $M_3(\gamma_1, \gamma_2)$ , such that

$$\| K_{n_1, n_2}^a(f; \cdot, \cdot) \|_{\gamma_1, \gamma_2} \leq M_3(\gamma_1, \gamma_2) \| f \|_{\gamma_1, \gamma_2} \tag{3.1}$$

for every  $f \in C_{\gamma_1, \gamma_2}(I)$  and for all  $n_1, n_2 \in \mathbb{N}$ .

**Proof .** From equation (2.2) and Lemma 3.5, we get

$$\begin{aligned} \omega_{\gamma_1, \gamma_2}(x, y) K_{n_1, n_2}^a\left(\frac{1}{\omega_{\gamma_1, \gamma_2}(u, v)}; x, y\right) &= \left(\omega_{\gamma_1}(x) K_{n_1}^a\left(\frac{1}{\omega_{\gamma_1}(u)}; x\right)\right) \left(\omega_{\gamma_2}(y) K_{n_2}^a\left(\frac{1}{\omega_{\gamma_2}(v)}; y\right)\right) \\ &\leq M_1(\gamma_1) M_2(\gamma_2), \text{ for all } (x, y) \in I \text{ and } n_1, n_2 \in \mathbb{N}. \end{aligned} \tag{3.2}$$

Taking supremum on the left side of the inequality of (3.2) and using (2.3), we obtain

$$\left\| K_{n_1, n_2}^a\left(\frac{1}{\omega_{\gamma_1, \gamma_2}(u, v)}; \cdot, \cdot\right) \right\|_{\gamma_1, \gamma_2} \leq M_4(\gamma_1, \gamma_2). \tag{3.3}$$

Now,  $\| K_{n_1, n_2}^a(f; \cdot, \cdot) \|_{\gamma_1, \gamma_2} \leq \| f \|_{\gamma_1, \gamma_2} \left\| K_{n_1, n_2}^a\left(\frac{1}{\omega_{\gamma_1, \gamma_2}(u, v)}; \cdot, \cdot\right) \right\|_{\gamma_1, \gamma_2}.$

From (3.3), we get the desired result. □

## 4 Main results

### 4.1 Local approximation

For  $f \in C_B(I)$  (the space of all bounded and uniformly continuous functions on  $I$ , let  $C_B^2(I) = \{f \in C_B(I) : f^{(p,q)} \in C_B(I), 0 \leq p+q \leq 2\}$ , where  $f^{(p,q)}$  is  $(p, q)$ th-order partial derivative with respect to  $x, y$  of  $f$ , equipped with the norm

$$\|f\|_{C_B^2(I)} = \|f\|_{C_B(I)} + \sum_{i=1}^2 \left\| \frac{\partial^i f}{\partial x^i} \right\|_{C_B(I)} + \sum_{i=1}^2 \left\| \frac{\partial^i f}{\partial y^i} \right\|_{C_B(I)}.$$

The Peetre’s  $K$ –functional of the function  $f \in C_B(I)$  is given by

$$\mathcal{K}(f; \delta) = \inf_{g \in C_B^2(I)} \{ \|f - g\|_{C_B(I)} + \delta \|g\|_{C_B^2(I)}, \delta > 0 \}.$$

It is also known that the following inequality

$$\mathcal{K}(f; \delta) \leq M_1 \{ \bar{\omega}_2(f; \sqrt{\delta}) + \min(1, \delta) \|f\|_{C_B(I)} \} \tag{4.1}$$

holds for all  $\delta > 0$  ([9], page 192). The constant  $M_1$  is independent of  $\delta$  and  $f$  and  $\bar{\omega}_2(f; \sqrt{\delta})$  is the second order modulus of continuity.

For  $f \in C_B(I)$ , the complete modulus of continuity for bivariate case is defined as follows:

$$\omega(f; \delta) = \sup \left\{ |f(u, v) - f(x, y)| : (u, v), (x, y) \in I \text{ and } \sqrt{(u - x)^2 + (v - y)^2} \leq \delta \right\}.$$

Further, the partial moduli of continuity with respect to  $x$  and  $y$  is defined as

$$\omega_1(f; \delta) = \sup \left\{ |f(x_1, y) - f(x_2, y)| : y \in \mathbb{R}^0 \text{ and } |x_1 - x_2| \leq \delta \right\},$$

$$\omega_2(f; \delta) = \sup \left\{ |f(x, y_1) - f(x, y_2)| : x \in \mathbb{R}^0 \text{ and } |y_1 - y_2| \leq \delta \right\}.$$

It is clear that they satisfy the properties of the usual modulus of continuity. The details of the modulus of continuity for the bivariate case can be found in [7].

Now, we find the order of approximation of the sequence  $K_{n_1, n_2}^a(f; x, y)$  to the function  $f(x, y) \in C_B(I)$  by Peetre’s  $K$ –functional.

**Theorem 1.** For the function  $f \in C_B(I)$ , the following inequality

$$\begin{aligned} |K_{n_1, n_2}^a(f; x, y) - f(x, y)| &\leq 4\mathcal{K}(f; M_{n_1, n_2}(x, y)) \\ &+ \omega \left( f; \sqrt{\left( \frac{1}{n_1 + 1} \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right) \right)^2 + \left( \frac{1}{n_2 + 1} \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right) \right)^2} \right) \\ &\leq M \left\{ \bar{\omega}_2 \left( f; \sqrt{M_{n_1, n_2}(x, y)} \right) + \min\{1, M_{n_1, n_2}(x, y)\} \|f\|_{C_B^2(I)} \right\} \\ &+ \omega \left( f; \sqrt{\left( \frac{1}{n_1 + 1} \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right) \right)^2 + \left( \frac{1}{n_2 + 1} \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right) \right)^2} \right) \end{aligned}$$

holds. The constant  $M > 0$  is independent of  $f$  and  $M_{n_1, n_2}(x, y)$ , where

$$M_{n_1, n_2}(x, y) = \frac{\{\delta_{n_1}^a(x)\}^2}{n_1 + 1} + \frac{\{\delta_{n_2}^a(y)\}^2}{n_2 + 1}.$$

**Proof .** We define the auxiliary operators as follows:

$$\bar{K}_{n_1, n_2}^a(f; x, y) = K_{n_1, n_2}^a(f; x, y) - f \left( \frac{1}{n_1 + 1} \left( n_1 x + \frac{ax}{1+x} + \frac{1}{2} \right), \frac{1}{n_2 + 1} \left( n_2 y + \frac{ay}{1+y} + \frac{1}{2} \right) \right) + f(x, y). \tag{4.2}$$

Then, from Lemma 3.3, we have  $\bar{K}_{n_1, n_2}^a((u - x); x, y) = 0$  and  $\bar{K}_{n_1, n_2}^a((v - y); x, y) = 0$ .

Let  $g \in C_B^2(I)$  and  $(u, v) \in I$ . Using the Taylor’s theorem, we have

$$g(u, v) - g(x, y) = \frac{\partial g(x, y)}{\partial x} (u - x) + \int_x^u (u - \alpha) \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} d\alpha + \frac{\partial g(x, y)}{\partial y} (v - y) + \int_y^v (v - \beta) \frac{\partial^2 g(x, \beta)}{\partial \beta^2} d\beta. \tag{4.3}$$

Operating  $\bar{K}_{n_1, n_2}^a$  on both sides of (4.3), we get

$$\begin{aligned} \bar{K}_{n_1, n_2}^a(g; x, y) - g(x, y) &= \bar{K}_{n_1, n_2}^a \left( \int_x^u (u - \alpha) \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} d\alpha; x, y \right) + \bar{K}_{n_1, n_2}^a \left( \int_y^v (v - \beta) \frac{\partial^2 g(x, \beta)}{\partial \beta^2} d\beta; x, y \right) \\ &= K_{n_1, n_2}^a \left( \int_x^u (u - \alpha) \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} d\alpha; x, y \right) \end{aligned}$$

$$\begin{aligned}
 & - \int_x^{\frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right)} \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right) - \alpha \right) \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} d\alpha + K_{n_1, n_2}^a \left( \int_y^v (v - \beta) \frac{\partial^2 g(x, \beta)}{\partial \beta^2} d\beta; x, y \right) \\
 & - \int_y^{\frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right)} \left( \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) - \beta \right) \frac{\partial^2 g(x, \beta)}{\partial \beta^2} d\beta.
 \end{aligned}$$

Hence,

$$\begin{aligned}
 |\overline{K}_{n_1, n_2}^a(g; x, y) - g(x, y)| & \leq K_{n_1, n_2}^a \left( \left| \int_x^u |u - \alpha| \left| \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} \right| d\alpha \right|; x, y \right) \\
 & + \left| \int_x^{\frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right)} \left| \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right) - \alpha \right| \left| \frac{\partial^2 g(\alpha, y)}{\partial \alpha^2} \right| d\alpha \right| \\
 & + K_{n_1, n_2}^a \left( \left| \int_y^v |v - \beta| \left| \frac{\partial^2 g(x, \beta)}{\partial \beta^2} \right| d\beta \right|; x, y \right) \\
 & + \left| \int_y^{\frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right)} \left| \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) - \beta \right| \left| \frac{\partial^2 g(x, \beta)}{\partial \beta^2} \right| d\beta \right| \\
 & \leq \left\{ K_{n_1, n_2}^a((u - x)^2; x, y) + \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right) - x \right)^2 \right\} \|g\|_{C_B^2(I)} \\
 & + \left\{ K_{n_1, n_2}^a((v - y)^2; x, y) + \left( \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) - y \right)^2 \right\} \|g\|_{C_B^2(I)} \\
 & \leq \left\{ \frac{1}{n_1+1} \{\delta_{n_1}^a(x)\}^2 + \left( \frac{1}{n_1+1} \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right) \right)^2 \right. \\
 & \left. + \frac{1}{n_2+1} \{\delta_{n_2}^a(y)\}^2 + \left( \frac{1}{n_2+1} \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right) \right)^2 \right\} \|g\|_{C_B^2(I)}.
 \end{aligned}$$

Thus, we get

$$|\overline{K}_{n_1, n_2}^a(g; x, y) - g(x, y)| \leq \left\{ \frac{2}{n_1+1} \{\delta_{n_1}^a(x)\}^2 + \frac{2}{n_2+1} \{\delta_{n_2}^a(y)\}^2 \right\} \|g\|_{C_B^2(I)}.$$

Also,

$$\begin{aligned}
 |\overline{K}_{n_1, n_2}^a(f; x, y)| & \leq |K_{n_1, n_2}^a(f; x, y)| + \left| f \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right), \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) \right) \right| + |f(x, y)| \\
 & \leq 3\|f\|_{C_B(I)}.
 \end{aligned} \tag{4.4}$$

Now, from equation (4.4), we get

$$\begin{aligned}
 |K_{n_1, n_2}^a(f; x, y) - f(x, y)| & \leq |\overline{K}_{n_1, n_2}^a(f - g; x, y)| + |\overline{K}_{n_1, n_2}^a(g; x, y) - g(x, y)| + |g(x, y) - f(x, y)| \\
 & + \left| f \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right), \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) \right) - f(x, y) \right| \\
 & \leq 3\|f - g\|_{C_B(I)} + \|f - g\|_{C_B(I)} + |\overline{K}_{n_1, n_2}^a(g; x, y) - g(x, y)| \\
 & + \left| f \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right), \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) \right) - f(x, y) \right| \\
 & \leq 4\|f - g\|_{C_B(I)} + \left\{ \frac{2}{n_1+1} \{\delta_{n_1}^a(x)\}^2 + \frac{2}{n_2+1} \{\delta_{n_2}^a(y)\}^2 \right\} \|g\|_{C_B^2(I)} \\
 & + \left| f \left( \frac{1}{n_1+1} \left( n_1x + \frac{ax}{1+x} + \frac{1}{2} \right), \frac{1}{n_2+1} \left( n_2y + \frac{ay}{1+y} + \frac{1}{2} \right) \right) - f(x, y) \right|
 \end{aligned}$$

$$\leq \left( 4\|f - g\|_{C_B(I)} + 2M_{n_1, n_2}(x, y)\|g\|_{C_B^2(I)} \right) + \omega\left(f; \sqrt{\left(\frac{1}{n_1 + 1}\left(-x + \frac{ax}{1 + x} + \frac{1}{2}\right)\right)^2 + \left(\frac{1}{n_2 + 1}\left(-y + \frac{ay}{1 + y} + \frac{1}{2}\right)\right)^2}\right).$$

Taking infimum on the right hand side over all  $g \in C_B^2(I)$  and using (4.1), we obtain

$$\begin{aligned} |K_{n_1, n_2}^a(f; x, y) - f(x, y)| &\leq 4\mathcal{K}(f; M_{n_1, n_2}(x, y)) \\ &\quad + \omega\left(f; \sqrt{\left(\frac{1}{n_1 + 1}\left(-x + \frac{ax}{1 + x} + \frac{1}{2}\right)\right)^2 + \left(\frac{1}{n_2 + 1}\left(-y + \frac{ay}{1 + y} + \frac{1}{2}\right)\right)^2}\right) \\ &\leq M\left\{\bar{\omega}_2\left(f; \sqrt{M_{n_1, n_2}(x, y)}\right) + \min\{1, M_{n_1, n_2}(x, y)\}\|f\|_{C_B^2(I)}\right\} \\ &\quad + \omega\left(f; \sqrt{\left(\frac{1}{n_1 + 1}\left(-x + \frac{ax}{1 + x} + \frac{1}{2}\right)\right)^2 + \left(\frac{1}{n_2 + 1}\left(-y + \frac{ay}{1 + y} + \frac{1}{2}\right)\right)^2}\right). \end{aligned}$$

Hence, the proof is completed.  $\square$

### 4.2 Rate of convergence of bivariate operators

**Theorem 2.** Suppose that  $f \in C_{\gamma_1, \gamma_2}^1(I)$  with  $\gamma_1, \gamma_2 \in \mathbb{N}^0$  then there exist a positive constant  $M_5(\gamma_1, \gamma_2)$  such that for all  $(x, y) \in I$  and  $n_1, n_2 \in \mathbb{N}$

$$\omega_{\gamma_1, \gamma_2}(x, y)|K_{n_1, n_2}^a(f; x, y) - f(x, y)| \leq M_5(\gamma_1, \gamma_2)\left\{\|f_x\|_{\gamma_1, \gamma_2} \frac{\delta_{n_1}^a(x)}{\sqrt{n_1 + 1}} + \|f_y\|_{\gamma_1, \gamma_2} \frac{\delta_{n_2}^a(y)}{\sqrt{n_2 + 1}}\right\}.$$

**Proof .** Let  $(x, y) \in I$  be a fixed point. Then, we have

$$\begin{aligned} f(t, z) - f(x, y) &= \int_x^t f_u(u, z)du + \int_y^z f_v(x, v)dv \quad \text{for } (t, z) \in I \\ K_{n_1, n_2}^a(f(t, z); x, y) - f(x, y) &= K_{n_1, n_2}^a\left(\int_x^t f_u(u, z)du; x, y\right) + K_{n_1, n_2}^a\left(\int_y^z f_v(x, v)dv; x, y\right). \end{aligned} \tag{4.5}$$

Now, by using (2.3), we get

$$\left|\int_x^t f_u(u, z)du\right| \leq \|f_x\|_{\gamma_1, \gamma_2} \left|\int_x^t \frac{du}{\omega_{\gamma_1, \gamma_2}(u, z)}\right| \leq \|f_x\|_{\gamma_1, \gamma_2} \left(\frac{1}{\omega_{\gamma_1, \gamma_2}(t, z)} + \frac{1}{\omega_{\gamma_1, \gamma_2}(x, z)}\right)|t - x|,$$

and analogously

$$\left|\int_y^z f_v(x, v)dv\right| \leq \|f_y\|_{\gamma_1, \gamma_2} \left(\frac{1}{\omega_{\gamma_1, \gamma_2}(x, z)} + \frac{1}{\omega_{\gamma_1, \gamma_2}(x, y)}\right)|z - y|.$$

By using these inequalities and from (2.2), we obtain for  $n_1, n_2 \in \mathbb{N}$

$$\begin{aligned} \omega_{\gamma_1, \gamma_2}(x, y)\left|K_{n_1, n_2}^a\left(\int_x^t f_u(u, z)du; x, y\right)\right| &\leq \omega_{\gamma_1, \gamma_2}(x, y)K_{n_1, n_2}^a\left(\left|\int_x^t f_u(u, z)du\right|; x, y\right) \\ &\leq \|f_x\|_{\gamma_1, \gamma_2} \omega_{\gamma_1, \gamma_2}(x, y)\left\{K_{n_1, n_2}^a\left(\frac{|t - x|}{\omega_{\gamma_1, \gamma_2}(t, z)}; x, y\right) + K_{n_1, n_2}^a\left(\frac{|t - x|}{\omega_{\gamma_1, \gamma_2}(x, z)}; x, y\right)\right\} \\ &= \|f_x\|_{\gamma_1, \gamma_2} \omega_{\gamma_2}(y)K_{n_2}^a\left(\frac{1}{\omega_{\gamma_2}(z)}; y\right) \times \left\{\omega_{\gamma_1}(x)K_{n_1}^a\left(\frac{|t - x|}{\omega_{\gamma_1}(t)}; x\right) + K_{n_1}^a(|t - x|; x)\right\}, \end{aligned} \tag{4.6}$$

and analogously

$$\omega_{\gamma_1, \gamma_2}(x, y) \left| K_{n_1, n_2}^a \left( \int_y^z f_v(x, v) dv; x, y \right) \right| \leq \|f_y\|_{\gamma_1, \gamma_2} \left\{ \omega_{\gamma_2}(y) K_{n_2}^a \left( \frac{|z-y|}{\omega_{\gamma_2}(z)}; y \right) + K_{n_2}^a(|z-y|; y) \right\}. \tag{4.7}$$

By the the Cauchy Schwarz inequality and Remark 3.4, we get for  $n_1 \in \mathbb{N}$

$$K_{n_1}^a(|t-x|; x) \leq (K_{n_1}^a((t-x)^2; x))^{1/2} (K_{n_1}^a(1; x))^{1/2} \leq \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}} \tag{4.8}$$

and

$$\omega_{\gamma_1}(x) K_{n_1}^a \left( \frac{|t-x|}{\omega_{\gamma_1}(t)}; x \right) \leq \omega_{\gamma_1}(x) \left( K_{n_1}^a \left( \frac{(t-x)^2}{\omega_{\gamma_1}(t)}; x \right) \right)^{1/2} \left( K_{n_1}^a \left( \frac{1}{\omega_{\gamma_1}(t)}; x \right) \right)^{1/2} \leq M_6(\gamma_1) \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}},$$

in view of Lemma 3.5. Analogously for  $n_2 \in \mathbb{N}$ , we have

$$K_{n_2}^a(|z-y|; y) \leq \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}}, \tag{4.9}$$

and

$$\omega_{\gamma_2}(y) K_{n_2}^a \left( \frac{|z-y|}{\omega_{\gamma_2}(z)}; y \right) \leq M_7(\gamma_2) \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}}. \tag{4.10}$$

From equations (4.5)-(4.10), we obtain

$$\omega_{\gamma_1, \gamma_2}(x, y) | K_{n_1, n_2}^a(f(t, z); x, y) - f(x, y) | \leq M_8(\gamma_1, \gamma_2) \left\{ \|f_x\|_{\gamma_1, \gamma_2} \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}} + \|f_y\|_{\gamma_1, \gamma_2} \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}} \right\},$$

for all  $n_1, n_2 \in \mathbb{N}$ . Thus the proof is completed.  $\square$

**Theorem 3.** Suppose that  $f \in C_{\gamma_1, \gamma_2}(I)$  with some  $\gamma_1, \gamma_2 \in \mathbb{N}^0$ . Then there exists a positive constant  $M_9(\gamma_1, \gamma_2)$  such that

$$\omega_{\gamma_1, \gamma_2}(x, y) | K_{n_1, n_2}^a(f(t, z); x, y) - f(x, y) | \leq M_9(\gamma_1, \gamma_2) \omega \left( f; C_{\gamma_1, \gamma_2}; \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}}, \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}} \right),$$

for all  $(x, y) \in I$  and  $n_1, n_2 \in \mathbb{N}$ .

**Proof .** Let  $f_{h, \delta}$  be the Steklov function of  $f \in C_{\gamma_1, \gamma_2}(I)$ , defined by the formula

$$f_{h, \delta}(x, y) := \frac{1}{h\delta} \int_0^h \int_0^\delta f(x+u, y+v) dv du, \tag{4.11}$$

$(x, y) \in I$  and  $h, \delta \in \mathbb{R}_+$ . From (4.11) it follows that

$$\begin{aligned} f_{h, \delta}(x, y) - f(x, y) &= \frac{1}{h\delta} \int_0^h \int_0^\delta \Delta_{u, v} f(x, y) dv du, \\ \frac{\partial}{\partial x} f_{h, \delta}(x, y) &= \frac{1}{h\delta} \int_0^\delta \Delta_{h, 0} f(x, y+v) dv \\ &= \frac{1}{h\delta} \int_0^\delta (\Delta_{h, v} f(x, y) - \Delta_{0, v} f(x, y)) dv, \\ \frac{\partial}{\partial y} f_{h, \delta}(x, y) &= \frac{1}{h\delta} \int_0^h \Delta_{0, \delta} f(x+u, y) du \\ &= \frac{1}{h\delta} \int_0^\delta (\Delta_{u, \delta} f(x, y) - \Delta_{u, 0} f(x, y)) du. \end{aligned}$$



Thus, from (2.3) and (2.4) we obtain

$$\| f_{h,\delta} - f \|_{\gamma_1,\gamma_2} \leq \omega(f, C_{\gamma_1,\gamma_2}; h, \delta), \tag{4.12}$$

$$\left\| \frac{\partial f_{h,\delta}}{\partial x} \right\|_{\gamma_1,\gamma_2} \leq 2h^{-1}\omega(f, C_{\gamma_1,\gamma_2}; h, \delta), \tag{4.13}$$

$$\left\| \frac{\partial f_{h,\delta}}{\partial y} \right\|_{\gamma_1,\gamma_2} \leq 2\delta^{-1}\omega(f, C_{\gamma_1,\gamma_2}; h, \delta). \tag{4.14}$$

For  $h, \delta \in \mathbb{R}_+$ , we can write

$$\begin{aligned} \omega_{\gamma_1,\gamma_2}(x, y) | K_{n_1,n_2}^a(f(t, z); x, y) - f(x, y) | \leq & \omega_{\gamma_1,\gamma_2}(x, y) \{ K_{n_1,n_2}^a(f(t, z) - f_{h,\delta}(t, z); x, y) \\ & + | K_{n_1,n_2}^a(f_{h,\delta}(t, z); x, y) - f_{h,\delta}(x, y) | \\ & + | f_{h,\delta}(x, y) - f(x, y) | \} := R_1 + R_2 + R_3. \end{aligned} \tag{4.15}$$

By (2.3), Lemma 3.6 and (4.12) it follows that

$$\begin{aligned} R_1 \leq \| K_{n_1,n_2}^a(f - f_{h,\delta}; \dots) \|_{\gamma_1,\gamma_2} & \leq M_{10}(\gamma_1, \gamma_2) \| f - f_{h,\delta} \|_{\gamma_1,\gamma_2} \\ & \leq M_{10}(\gamma_1, \gamma_2) \omega(f, C_{\gamma_1,\gamma_2}; h, \delta), \end{aligned}$$

and

$$R_3 \leq \omega(f, C_{\gamma_1,\gamma_2}; h, \delta).$$

By using Theorem 2 and (4.13) and (4.14), we get

$$\begin{aligned} R_2 \leq M_{11}(\gamma_1, \gamma_2) & \left\{ \left\| \frac{\partial f_{h,\delta}}{\partial x} \right\|_{\gamma_1,\gamma_2} \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}} + \left\| \frac{\partial f_{h,\delta}}{\partial y} \right\|_{\gamma_1,\gamma_2} \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}} \right\} \\ & \leq 2M_{11}(\gamma_1, \gamma_2) \omega(f, C_{\gamma_1,\gamma_2}; h, \delta) \left\{ h^{-1} \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}} + \delta^{-1} \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}} \right\}. \end{aligned}$$

Consequently, we drive from (4.15)

$$\omega_{\gamma_1,\gamma_2}(x, y) | K_{n_1,n_2}^a(f(t, z); x, y) - f(x, y) | \leq M_{12}(\gamma_1, \gamma_2) \omega(f, C_{\gamma_1,\gamma_2}; h, \delta) \left\{ 1 + h^{-1} \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}} + \delta^{-1} \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}} \right\}$$

for all  $(x, y) \in I, n_1, n_2 \in \mathbb{N}$  and  $h, \delta \in \mathbb{R}_+$ . On choosing  $h = \frac{\delta_{n_1}^a(x)}{\sqrt{n_1+1}}$  and  $\delta = \frac{\delta_{n_2}^a(y)}{\sqrt{n_2+1}}$ , we immediately obtain the required result.  $\square$

As a consequence of Theorem 3, we have

**Theorem 4.** Let  $f \in C_{\gamma_1,\gamma_2}(I)$  with some  $\gamma_1, \gamma_2 \in \mathbb{N}^0$ . Then for every  $(x, y) \in I$ ,

$$\lim_{n_1, n_2 \rightarrow \infty} K_{n_1, n_2}^a(f; x, y) = f(x, y).$$

**Theorem 5. (Voronovskaja type theorem)** Let  $f \in C_{\gamma_1,\gamma_2}^2(I)$ . Then for every  $(x, y) \in I$ ,

$$\begin{aligned} \lim_{n \rightarrow \infty} n \{ K_{n,n}^a(f; x, y) - f(x, y) \} = & \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right) f_x(x, y) + \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right) f_y(x, y) \\ & + \frac{x}{2}(1+x)f_{xx}(x, y) + \frac{y}{2}(1+y)f_{yy}(x, y). \end{aligned}$$

**Proof .** Let  $(x, y) \in I$  be fixed. By Taylor formula, we have

$$\begin{aligned}
 f(u, v) &= f(x, y) + f_x(x, y)(u - x) + f_y(x, y)(v - y) \\
 &\quad + \frac{1}{2}\{f_{xx}(x, y)(u - x)^2 + 2f_{xy}(x, y)(u - x)(v - y) + f_{yy}(x, y)(v - y)^2\} \\
 &\quad + \psi(u, v; x, y)\sqrt{(u - x)^4 + (v - y)^4},
 \end{aligned}$$

where  $\psi(\cdot, \cdot; x, y) \equiv \psi(\cdot, \cdot) \in C_{\gamma_1, \gamma_2}(I)$  and  $\psi(x, y) = 0$ . Thus, we get

$$\begin{aligned}
 K_{n,n}^a(f(u, v); x, y) &= f(x, y) + f_x(x, y)K_n^a(u - x; x) + f_y(x, y)K_n^a(v - y; y) \\
 &\quad + \frac{1}{2}\{f_{xx}(x, y)K_n^a((u - x)^2; x) + 2f_{xy}(x, y)K_n^a(u - x; x)K_n^a(v - y; y) \\
 &\quad + f_{yy}(x, y)K_n^a((v - y)^2; y)\} + K_{n,n}^a(\psi(u, v)\sqrt{(u - x)^4 + (v - y)^4}; x, y).
 \end{aligned}$$

Hence, using Lemma 3.1, we have

$$\begin{aligned}
 \lim_{n \rightarrow \infty} n\{K_{n,n}^a(f(u, v); x, y) - f(x, y)\} &= f_x(x, y)\left(-x + \frac{ax}{1+x} + \frac{1}{2}\right) + f_y(x, y)\left(-y + \frac{ay}{1+y} + \frac{1}{2}\right) \\
 &\quad + \frac{1}{2}\{x(1+x)f_{xx}(x, y) + y(1+y)f_{yy}(x, y)\} \\
 &\quad + \lim_{n \rightarrow \infty} nK_{n,n}^a(\psi(u, v)\sqrt{(u - x)^4 + (v - y)^4}; x, y). \tag{4.16}
 \end{aligned}$$

Applying the Hölder’s inequality, we have

$$\begin{aligned}
 |K_{n,n}^a(\psi(u, v)\sqrt{(u - x)^4 + (v - y)^4}; x, y)| &\leq \{K_{n,n}^a(\psi^2(u, v); x, y)\}^{1/2} \{K_{n,n}^a(((u - x)^4 + (v - y)^4); x, y)\}^{1/2} \\
 &\leq \{K_{n,n}^a(\psi^2(u, v); x, y)\}^{1/2} \{K_n^a((u - x)^4; x) + K_n^a((v - y)^4; y)\}^{1/2}.
 \end{aligned}$$

By Theorem 4

$$\lim_{n \rightarrow \infty} K_{n,n}^a(\psi^2(u, v); x, y) = \psi^2(x, y) = 0,$$

and from Lemma 3.1 (iii), for each  $(x, y) \in I$ ,  $K_n^a((u - x)^4; x) = O\left(\frac{1}{n^2}\right)$  and  $K_n^a((v - y)^4; y) = O\left(\frac{1}{n^2}\right)$ . Hence

$$\lim_{n \rightarrow \infty} nK_{n,n}^a(\psi(u, v)\sqrt{(u - x)^4 + (v - y)^4}; x, y) = 0. \tag{4.17}$$

By combining (4.16) and (4.17), we obtain the desired result.  $\square$

### 4.3 Simultaneous approximation

**Theorem 6.** Let  $f \in C_{\gamma_1, \gamma_2}^1(I)$ . Then for every  $(x, y) \in \mathbb{R}_+^2 = \mathbb{R}_+ \times \mathbb{R}_+$ ,

$$\lim_{n \rightarrow \infty} \left(\frac{\partial}{\partial \omega} K_{n,n}^a(f; \omega, y)\right)_{\omega=x} = \frac{\partial f}{\partial x}(x, y), \tag{4.18}$$

$$\lim_{n \rightarrow \infty} \left(\frac{\partial}{\partial \nu} K_{n,n}^a(f; x, \nu)\right)_{\nu=y} = \frac{\partial f}{\partial y}(x, y). \tag{4.19}$$

**Proof .** We shall prove only (4.18) because the proof of (4.19) is similar. By the Taylor formula for  $f \in C_{\gamma_1, \gamma_2}^1(I)$ , we have

$$f(u, v) = f(x, y) + f_x(x, y)(u - x) + f_y(x, y)(v - y) + \psi(u, v; x, y)\sqrt{(u - x)^2 + (v - y)^2} \text{ for } (u, v) \in I,$$

where  $\psi(u, v; x, y) \equiv \psi(\cdot, \cdot) \in C_{\gamma_1, \gamma_2}(I)$  and  $\psi(x, y) = 0$ . Operating  $K_{n,n}^a(\cdot; \cdot, y)$  to the above inequality and then by using Lemma 3.3, we get

$$\begin{aligned} \left(\frac{\partial}{\partial \omega} K_{n,n}^a(f(u, v); \omega, y)\right)_{\omega=x} &= f(x, y) \left(\frac{\partial}{\partial \omega} K_{n,n}^a(1; \omega, y)\right)_{\omega=x} + f_x(x, y) \left(\frac{\partial}{\partial \omega} K_{n,n}^a(u-x; \omega, y)\right)_{\omega=x} \\ &\quad + f_y(x, y) \left(\frac{\partial}{\partial \omega} K_{n,n}^a(v-y; \omega, y)\right)_{\omega=x} \\ &\quad + \left(\frac{\partial}{\partial \omega} K_{n,n}^a(\psi(u, v; x, y)\sqrt{(u-x)^2 + (v-y)^2}; \omega, y)\right)_{\omega=x}, \text{ for } (u, v) \in I \\ &= f_x(x, y) \left\{ \frac{\partial}{\partial \omega} \left( \frac{1}{n+1} \left( n\omega + \frac{a\omega}{1+\omega} + \frac{1}{2} \right) \right) \right\}_{\omega=x} \\ &\quad + f_y(x, y) \left\{ \frac{\partial}{\partial \omega} \left( \frac{1}{n+1} \left( ny + \frac{ay}{1+y} + \frac{1}{2} \right) \right) \right\}_{\omega=x} + E, \text{ say} \\ &= \frac{n}{n+1} f_x(x, y) + E. \end{aligned}$$

It is sufficient to prove that  $E \rightarrow 0$  as  $n \rightarrow \infty$ .

$$\begin{aligned} E &= (n+1)^2 \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \left(\frac{\partial}{\partial \omega} W_{n,n,k_1,k_2}^a(\omega, y)\right)_{\omega=x} \int_{\frac{k_2}{n+1}}^{\frac{k_2+1}{n+1}} \int_{\frac{k_1}{n+1}}^{\frac{k_1+1}{n+1}} \psi(u, v)\sqrt{(u-x)^2 + (v-y)^2} du dv \\ &= (n+1)^2 \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \frac{\{(k_1-nx)(1+x) - ax\}}{x(1+x)^2} W_{n,n,k_1,k_2}^a(x, y) \\ &\quad \times \int_{\frac{k_2}{n+1}}^{\frac{k_2+1}{n+1}} \int_{\frac{k_1}{n+1}}^{\frac{k_1+1}{n+1}} \psi(u, v)\sqrt{(u-x)^2 + (v-y)^2} du dv \\ &= \frac{n(n+1)^2}{x(1+x)} \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} \left(\frac{k_1}{n} - x\right) W_{n,n,k_1,k_2}^a(x, y) \int_{\frac{k_2}{n+1}}^{\frac{k_2+1}{n+1}} \int_{\frac{k_1}{n+1}}^{\frac{k_1+1}{n+1}} \psi(u, v)\sqrt{(u-x)^2 + (v-y)^2} du dv \\ &\quad - \frac{a}{(1+x)^2} K_{n,n}^a(\psi(u, v)\sqrt{(u-x)^2 + (v-y)^2}; x, y) \\ &= E_1 + E_2, \text{ say.} \end{aligned}$$

First, we estimate  $E_1$  by using Schwarz inequality.

$$\begin{aligned} E_1 &\leq \frac{n}{x(1+x)} \left( \sum_{k_1=0}^{\infty} W_{n,k_1}^a(x) \left(\frac{k_1}{n} - x\right)^2 \right)^{1/2} \\ &\quad \times \left( (n+1)^2 \sum_{k_1=0}^{\infty} \sum_{k_2=0}^{\infty} W_{n,n,k_1,k_2}^a(x, y) \int_{\frac{k_2}{n+1}}^{\frac{k_2+1}{n+1}} \int_{\frac{k_1}{n+1}}^{\frac{k_1+1}{n+1}} \psi^2(u, v)((u-x)^2 + (v-y)^2) du dv \right)^{1/2} \\ &\leq \frac{n}{x(1+x)} \left( \sum_{k_1=0}^{\infty} W_{n,k_1}^a(x) \left(\frac{k_1}{n} - x\right)^2 \right)^{1/2} \{ K_{n,n}^a(\psi^4(u, v); x, y)(K_n^a((u-x)^4; x) \\ &\quad + 2K_n^a((u-x)^2; x)(K_n^a((v-y)^2; y) + K_n^a((v-y)^4; y)) \}^{1/4} \end{aligned}$$

$$|E_1| \leq M_{12}(x, y) \{ K_{n,n}^a(\psi^4(u, v); x, y) \}^{1/4}, \text{ in view of Lemma 3.1.}$$

From Theorem 4, we obtain

$$\lim_{n \rightarrow \infty} K_{n,n}^a(\psi^4(u, v); x, y) = \psi^4(x, y) = 0, \text{ for } (x, y) \in \mathbb{R}_+^2.$$

To estimate  $E_2$ , proceeding in a manner similar to the estimate of  $E_1$ , we get  $E_2 \rightarrow 0$  as  $n \rightarrow \infty$ . Combining the estimates of  $E_1$  and  $E_2$ , it follows that  $E \rightarrow 0$  as  $n \rightarrow \infty$ . This completes the proof.  $\square$




Figure 1: The Convergence of  $K_{100,100}^{10}(f; x, y)$  (red) to  $f(x, y)$  (yellow)

Similarly, we can prove the following theorem:

**Theorem 7.** Let  $f \in C_{\gamma_1, \gamma_2}^3(I)$ . Then for every  $(x, y) \in \mathbb{R}_+^2$ , we have

$$\begin{aligned} \lim_{n \rightarrow \infty} n \left\{ \left( \frac{\partial}{\partial \omega} K_{n,n}^a(f; \omega, y) \right)_{\omega=x} - \frac{\partial f}{\partial x}(x, y) \right\} &= \left( -1 + \frac{a}{(1+x)^2} \right) f_x(x, y) + \left( 1 + \frac{ax}{1+x} \right) f_{xx}(x, y) \\ &+ \left( -y + \frac{ay}{1+y} + \frac{1}{2} \right) f_{xy}(x, y) + \frac{y}{2}(1+y) f_{xyy}(x, y) \\ &+ \frac{x}{2}(1+x) f_{xxx}(x, y) \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} n \left\{ \left( \frac{\partial}{\partial \nu} K_{n,n}^a(f; x, \nu) \right)_{\nu=y} - \frac{\partial f}{\partial y}(x, y) \right\} &= \left( -1 + \frac{a}{(1+y)^2} \right) f_y(x, y) + \left( 1 + \frac{ay}{1+y} \right) f_{yy}(x, y) \\ &+ \left( -x + \frac{ax}{1+x} + \frac{1}{2} \right) f_{xy}(x, y) + \frac{x}{2}(1+x) f_{xxy}(x, y) \\ &+ \frac{y}{2}(1+y) f_{yyy}(x, y). \end{aligned}$$

### 5 Numerical Examples

In the following, we give some numerical results regarding the approximation properties of bivariate generalized Baskakov-Kantorovich operators  $K_{n_1, n_2}^a(f; x, y)$  using Matlab algorithms for construction of operators.

Let us consider the function  $f : I \rightarrow \mathbb{R}^0, f(x, y) = x^2y^2 - 9xy^2 + 4x^2$ . The convergence of the bivariate generalized Baskakov-Kantorovich operators to the function  $f$  is illustrated in Example 1.

**Example 1.** For  $n_1 = n_2 = 100; n_1 = n_2 = 500$  and  $a = 10$ , the convergence of the bivariate generalized Baskakov-Kantorovich operators  $K_{n_1, n_2}^a(f; x, y)$  (red) to the function  $f(x, y) = x^2y^2 - 9xy^2 + 4x^2$  (yellow) is illustrated in figures 1 and 2 respectively. We observe that as the values of  $n_1$  and  $n_2$  increase, the error in the approximation of the function by the operators becomes smaller.




Figure 2: The Convergence of  $K_{500,500}^{10}(f; x, y)$  (red) to  $f(x, y)$  (yellow)




Figure 3. The Comparison of bivariate Szász-Kantorovich (blue) and bivariate generalized Baskakov-Kantorovich  $K_{100,100}^{10}(f; x, y)$  (red) to  $f(x, y)$  (yellow)




Figure 4. The Comparison of bivariate Szász-Kantorovich (blue) and bivariate generalized Baskakov-Kantorovich  $K_{500,500}^{10}(f; x, y)$  (red) to  $f(x, y)$  (yellow)

**Example 2.** For  $n_1 = n_2 = 100$ ;  $n_1 = n_2 = 500$  and  $a = 10$ , the comparison of the bivariate generalized Baskakov-Kantorovich operators  $K_{n_1, n_2}^a(f; x, y)$  (red) and the bivariate Szász-Kantorovich operators (blue) to the function  $f(x, y) = x^2y^2 - 9xy^2 + 4x^2$  (yellow) is illustrated in figures 5 and 5 respectively. It is observed that the error in the approximation of  $f$  by the bivariate Szász-Kantorovich operators is smaller than the bivariate generalized Baskakov-Kantorovich operators.

## References

- [1] A.M. Acu, T. Acar, C.V. Muraru and V.A. Radu, *Some approximation properties by a class of bivariate operators*, Math. Meth. Appl. Sci. **42** (2019), 5551–5565.
- [2] O. Agratini, *Bivariate positive operators in polynomial weighted spaces*, Abstr. Appl. Anal. **2013** (2013), 8.
- [3] P.N. Agrawal and M. Goyal, *Generalized Baskakov Kantorovich operators*, Filomat 31(19) (2017) 6131–6151.
- [4] P.N. Agrawal, V. Gupta, and A. Sathish Kumar, *Generalized Baskakov-Durrmeyer type operators*, Rend. Circ. Mat. Palermo **63** (2014), no. 2, 193–209.
- [5] F. Altomare and M. Campiti, *Korovkin-Type Approximation Theory and its Applications*, de Gruyter Studies in Mathematics, 17. Walter de Gruyter and Co., Berlin 1994.
- [6] F. Altomare and M. Cappelletti Montano and V. Leonessa, *On a generalization of Szász-Mirakjan-Kantorovich operators*, Results Math. **63** (2013), no. 3-4, 837–863.
- [7] G.A. Anastassiou and S. Gal, *Approximation Theory. Moduli of Continuity and Global Smoothness Preservation*, Birkhäuser, Boston, 2000.
- [8] D. Barbosu, *Some generalized bivariate Bernstein operators*, Math. Notes (Miskolc) **1** (2000), no. 1, 3–10.
- [9] P.L. Butzer and H. Berens, *Semi-groups of Operators and Approximation*, Springer, New York, 1967.
- [10] O. Dođru and V. Gupta, *Korovkin-type approximation properties of bivariate  $q$ -Meyer-König and Zeller operators*, Calcolo **43** (2006), no. 1, 51–63.
- [11] A. Erençin, *Durrmeyer type modification of generalized Baskakov operators*, Appl. Math. Comput. **218** (2011), no. 8, 4384–4390.
- [12] B. Firlej and L. Rempulska, *Approximation of functions of several variables by some operators of the Szász-Mirakjan type*, Fasc. math. 27 (1997) 15–27.
- [13] M. Gurdek, L. Rempulska and M. Skorupka, *The Baskakov operators for functions of two variables*, Collect. Math. **50** (1999), no. 3, 289–302.
- [14] A. Kajla and M. Goyal, *Modified Bernstein-Kantorovich operators for functions of one and two variables*, Rend. Circ. Mat. di Palermo II **67** (2018), no. 2, 379–395.
- [15] V. Miheşan, *Uniform approximation with positive linear operators generated by generalized Baskakov method*, Automat. Comput. Appl. Math. **7** (1998), no. 1, 34–37.
- [16] M. Nasiruzzaman, *Approximation properties by Szász-Mirakjan operators to bivariate functions via Dunkl analogue*, Iran. J. Sci. Technol. Trans. A, Sci. **45** (2021), 259–269.
- [17] F. Özger, *Weighted statistical approximation properties of univariate and bivariate  $\lambda$ -Kantorovich operators*. Filomat **33** (2019), no. 11, 3473–3486.
- [18] L. Rempulska and M. Skorupka, *On convergence of first derivatives of certain Szász-Mirakjan type operators*, Rend. Mat. Appl. **19** (1999), no. 2, 269–279.
- [19] M. Skorupka, *Approximation of functions of two variables by some linear positive operators*, Matematiche (Catania) **50** (1995), no. 2, 323–336.
- [20] DD. Stancu, *A new class of uniform approximating polynomial operators in two and several variables*. G. Alexits, S. B. Stechkin(eds.): Proc. Conf. Consecutive Theory Functions, Budapest: Akad. Kiadó, 1972, pp. 443–455.
- [21] A. Wafi and S. Khatoun, *Approximation by generalized Baskakov operators for functions of one and two variables in exponential and polynomial weight spaces*, Thai. J. Math. **2** (2004) 53–66.

- [22] A. Wafi and S. Khatoon, *Convergence and Voronovskaja-type theorems for derivatives of generalized Baskakov operators*, Cent. Eur. J. Math. **6** (2008), no. 2, 325–334.
- [23] R. Yadav, R. Meher and V.N. Mishra, *Results on bivariate Szász-Mirakjan type operators in polynomial weight spaces*, Math. Eng. Sci. Aerosp. **11** (2020), no. 4, 939–957.
- [24] R. Yadav, R. Meher and V.N. Mishra, *Quantitative estimations of bivariate summation-integral-type operators*, Math. Meth. Appl. Sci. **42** (2019), 7172–7191.
- [25] G. You and P. Xaun, *Weighted approximation by multidimensional Baskakov operators*, J. Math. Res. Expos. **20** (2000), no. 1, 43–50.