

A new univalent integral operator defined by the Opoola differential operator

Bitrus Sambo^{a,*}, Timothy Oloyede Opoola^b

^aDepartment of Mathematics, Gombe State University, P.M.B. 127, Gombe, Nigeria

^bDepartment of Mathematics, University of Ilorin, P.M.B. 1515, Ilorin, Nigeria

(Communicated by Mugur Alexandru Acu)

Abstract

In this investigation, using Opoola differential operator ($D^m(\mu, \beta, t)f(z)$), a new integral operator: $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) : A^n \rightarrow A$ is defined in the unit disk, $U = \{z \in C : |z| < 1\}$; and we investigated the Univalence conditions of this generalized operator. Finally, a number of corollaries and remarks which show the extension of our results are presented.

Keywords: Analytic functions, Univalent functions starlike functions, convex functions, close-to-convex functions, Integral operator

2020 MSC: 30C45, 30C50, 30C55

1 Introduction and definitions

Let A denote the family of functions, $f(z)$ that are analytic in the unit disk, $U = \{z \in C : |z| < 1\}$. Let S be the subclass of $f(z) \in A$ that are of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k \quad (1.1)$$

which are univalent in U . Let S^* , C and K denote respectively the subclasses of S known as class of starlike functions with respect to the origin, class of convex functions and class of Close-to-convex functions. Let A and S denote the classes of all functions as defined above. Next, we define some well known subclasses of A , denoted by A_2 , K , K_2 , $K_{2,\delta}$ and $S(p)$ respectively as follow:

$$A_2 \subset A = \left\{ f \in A : f(z) = z + \sum_{k=3}^{\infty} a_k z^k \quad (z \in U, a_2 = 0) \right\}$$

$$K \subset A = \left\{ f \in A : \left| \frac{z^2 f'(z)}{(f(z))^2} - 1 \right| < 1 \quad z \in U \right\}$$

$$K_2 \subset K = \{f \in K : f''(0) = 0 \quad z \in U\}$$

*Corresponding author

Email addresses: bitrussambo3@gmail.com (Bitrus Sambo), opoola.to@unilorin.edu.ng (Timothy Oloyede Opoola)

$$K_{2,\delta} \subset K_2 = \left\{ f \in K_2 : \left| \frac{z^2 f'(z)}{(f(z))^2} - 1 \right| < 1 - \delta, \quad 0 \leq \delta < 1 \quad z \in U \right\}$$

$$S(p) \subset A = \left\{ f \in A : \left| \left(\frac{z}{f(z)} \right)'' \right| \leq p, \quad 0 < p \leq 2, \quad p \in \Re \quad z \in U \right\}$$

In [22], Opoola introduced the following differential operator:

$$D^m(\mu, \beta, t)f(z) : A \rightarrow A$$

$$D^0(\mu, \beta, t)f(z) = f(z)$$

$$D^1(\mu, \beta, t)f(z) = zD_t f(z) = tzf'(z) - z(\beta - \mu)t + [1 + (\beta - \mu - 1)t]f(z)$$

$$D^m(\mu, \beta, t)f(z) = zD_t(D^{m-1}(\mu, \beta, t)f(z)), \quad m \in N \quad (1.2)$$

If $f(z)$ is given by (1.1), then from (1.2), we see that

$$D^m(\mu, \beta, t)f(z) = z + \sum_{k=2}^{\infty} [1 + (k + \beta - \mu - 1)t]^m a_k z^k \quad (1.3)$$

where, $0 \leq \mu \leq \beta$, $t \geq 0$ and $m \in N_0 = N \cup 0$.

Remark 1.1. (i) when $\beta = \mu, t = \lambda$, $D^n(\mu, \mu, \lambda)f(z) = D_\lambda^n f(z)$ is the Al-Oboudi Differential operator in [1].

(ii) when $\beta = \mu, t = 1$, $D^n(\mu, \mu, 1)f(z) = D^n f(z)$ is the Salagean Differential operator introduced in [30].

Alexander in [2] introduced and investigated a univalent Integral operator and was continued by Libera in [18]. In [4], Bernardi generalized the work of Libera where he gave the conditions for which the operator is starlike, convex and close-to-convex in the unit disk, U . Being an interesting area of research, many results have been obtained by many scholars. We refer interested readers to the works in [3, 5, 7, 8, 13, 14, 16, 17, 19, 23, 24, 25, 27, 28, 31] for more information on different univalent integral operators.

Motivated by the works of Bulut [9, 10, 11, 12, 15], where a differential operator was used to define integral operators; we generalize Bulut's work and others by using Opoola differential operator and defined a generalized Integral operator, and investigated the conditions for which the operator is univalent in the unit disk, $U = \{z \in C : |z| < 1\}$. The operator and the results generalize some existing results.

Now we introduce a new general integral operator by means of Opoola differential operator.

Definition 1.2. Let $n \in N, m \in N_0, t, \beta \geq 0, 0 \leq \mu \leq \beta, \sigma \in C$ with $\Re(\sigma) > 0$ and $\alpha_j \in C, (j = 1, 2, \dots, n)$, we define the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) : A^n \rightarrow A$ by

$$I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) := \left\{ \sigma \int_0^z u^{(\sigma-1)} \prod_{j=1}^n \left(\frac{D^m(\mu, \beta, t)f_j(u)}{u} \right)^{\alpha_j} du \right\}^{\frac{1}{\sigma}}, \quad z \in U \quad (1.4)$$

where $f_1, \dots, f_n \in A$ and $D^m(\mu, \beta, t)f_j(z)$ is the Opoola differential operator defined in (1.3).

Remark 1.3. (i) For $n \in N, \sigma = 1, \alpha_j \in C, j = 1, 2, \dots, n$ in (1.4), we have the integral operator

$$I_{t,\beta,\mu}^m(f_1, \dots, f_n)(z) := \left\{ \int_0^z \prod_{j=1}^n \left(\frac{D^m(\mu, \beta, t)f_j(u)}{u} \right)^{\alpha_j} du \right\}, \quad z \in U \quad (1.5)$$

where $f_1, \dots, f_n \in A$ and $D^m(\mu, \beta, t)f_j(z)$ is the Opoola differential operator defined in (1.3).

(ii) For $n = 1, \sigma = 1, \alpha \in C$ in (1.4), we have the integral operator

$$I_{t,\beta,\mu}^m f(z) := \left\{ \int_0^z \left(\frac{D^m(\mu, \beta, t)f(u)}{u} \right)^\alpha du \right\}, \quad z \in U \quad (1.6)$$

where $f \in A$ and $D^m(\mu, \beta, t)f(z)$ is the Opoola differential operator defined in (1.3).

(iii) For $n \in N, m \in N_0, \sigma = 1, \alpha_j \in C, \beta = \mu, t = \lambda$ and $D^m(\mu, \mu, \lambda)f_j(z) = D_\lambda^m f_j(z), j = 1, 2, \dots, n$, we have the integral operator

$$I(f_1, \dots, f_n)(z) := \int_0^z \left(\frac{D_\lambda^m f_1(u)}{u} \right)^{\alpha_1} \dots \left(\frac{D_\lambda^m f_n(u)}{u} \right)^{\alpha_n} du, \quad z \in U \tag{1.7}$$

which was introduced by Bulut, $D_\lambda^m f_j(z)$ is the Al-Oboudi differential operator in [15].

(iv) For $n \in N, \sigma, \alpha_j \in C, D^0(\mu, \beta, t)f_j(z) = f_j(z) \in S, j = 1, 2, \dots, n$, we have the integral operator

$$I_\sigma(f_1, \dots, f_n)(z) := \left\{ \sigma \int_0^z \left(\frac{f_1(u)}{u} \right)^{\alpha_1} \dots \left(\frac{f_n(u)}{u} \right)^{\alpha_n} du \right\}^{\frac{1}{\sigma}}, \quad z \in U \tag{1.8}$$

in [6].

(v) For $n \in N, \sigma = 1, \alpha_j \in C, D^0(\mu, \beta, t)f_j(z) = f_j(z) \in S, j = 1, 2, \dots, n$, we have the integral operator

$$I(f_1, \dots, f_n)(z) := \left\{ \int_0^z \left(\frac{f_1(u)}{u} \right)^{\alpha_1} \dots \left(\frac{f_n(u)}{u} \right)^{\alpha_n} du \right\}, \quad z \in U \tag{1.9}$$

in [6].

(vi) For $m = 0, n = 1, \sigma = 1, \alpha_1 = 1, \alpha_2 = \alpha_3 = \dots = \alpha_n = 0$, and $D^0(\mu, \beta, t)f_1(z) = f(z) \in A$, we have Alexander integral operator

$$I(f)(z) := \int_0^z \frac{f(u)}{u} du, \quad z \in U \tag{1.10}$$

in [2].

(vii) For $m = 0, n = 1, \sigma = 1, \alpha[0, 1], \alpha_2 = \alpha_3 = \dots = \alpha_n = 0$, and $D^0(\mu, \mu, t)f_1(z) = f(z) \in A$, we have the integral operator

$$I(f)(z) := \int_0^z \left(\frac{f(u)}{u} \right)^\alpha du \quad z \in U, \tag{1.11}$$

in [20].

2 Relevant Lemmas

In order to prove our main results, we need the following Lemmas.

Lemma 2.1 (General Schwarz Lemma). [21] Let the analytic function $f(z)$ be regular in the disk

$$U_R = \{z \in C : |z| < R\},$$

with $|f(z)| \leq M$ for fixed M . If $f(z)$ has one zero and multiplicity order bigger than m for $z = 0$, then

$$|f(z)| \leq \frac{M}{R^m} |z|^m \quad (z \in U_R).$$

The equality can hold only if $f(z) = e^{i\theta} \frac{M}{R^m} z^m$, where θ is a constant.

Lemma 2.2. [29] Let α be a complex number with $Re\alpha > 0$ and $f(z) \in A$. If $f(z)$ satisfies

$$\frac{1 - |z|^{2\Re\alpha}}{\Re\alpha} \left| \frac{zf''(z)}{f'(z)} \right| \leq 1 \quad (z \in U),$$

then, for any complex number β with $\Re(\beta) \geq \Re(\alpha)$, the integral operator

$$F_\beta(z) := \left\{ \beta \int_0^z t^{\beta-1} f'(t) dt \right\}^{\frac{1}{\beta}} = z + \dots$$

is analytic and univalent in U .

Lemma 2.3. [26] Let $\alpha \in C$, ($\Re(\alpha) > 0$) and $c \in C(|c| \leq 1 : c \neq -1)$. Suppose also that the function $f(z)$ given by (1.1) is analytic in U . If

$$\left| c|z|^{2\alpha} + (1 - |z|^{2\alpha}) \frac{zf''(z)}{\alpha f'(z)} \right| \leq 1 \quad (z \in U),$$

then the function $F_\alpha(z)$ defined by

$$F_\alpha(z) := \left\{ \alpha \int_0^z t^{\alpha-1} f'(t) dt \right\}^{\frac{1}{\alpha}} = z + \dots$$

is analytic and univalent in U .

Lemma 2.4. [32] If a function $f(z) \in S(p)$, then $\left| \frac{z^2 f'(z)}{(f(z))^2} - 1 \right| \leq p|z|^2$.

3 Main Results

Theorem 3.1. Let $\alpha_1, \dots, \alpha_n, \sigma \in C, m \in N_0, t \geq 0, \beta \geq 0, 0 \leq \mu \leq \beta$, and each of the functions $f_j \in A, j = 1, \dots, n$, and

$$\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))} - 1 \right| \leq 1, \quad z \in U \quad (3.1)$$

and $\Re(\sigma) \geq \sum_{i=1}^n |\alpha_i| > 0$, then the integral operator $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent class S .

Proof . Since $f_j \in A, j = 1, \dots, n$ by (3.1), we have

$$\begin{aligned} \frac{(D^m(\mu, \beta, t)f_j(z))}{z} &= \frac{z + \sum_{k=2}^{\infty} [1 + (k + \beta - \mu - 1)t]^m a_{k,j} z^k}{z} \\ &= 1 + \sum_{k=2}^{\infty} [1 + (k + \beta - \mu - 1)t]^m a_{k,j} z^{k-1} \end{aligned}$$

for all $z \in U$ and $m \in N_0$. Let us define a function $h(z)$ by

$$h(z) = \int_0^z \prod_{j=1}^n \left(\frac{(D^m(\mu, \beta, t)f_j(u))}{u} \right)^{\alpha_j} du. \quad (3.2)$$

So that by Fundamental Theorem of Calculus, we have

$$h'(z) = \prod_{j=1}^n \left(\frac{(D^m(\mu, \beta, t)f_j(z))}{z} \right)^{\alpha_j} \quad (3.3)$$

for all $z \in U$. This equality implies that

$$\begin{aligned} \ln h'(z) &= \alpha_1 \ln \left(\frac{(D^m(\mu, \beta, t)f_1(z))}{z} \right) + \dots + \alpha_n \ln \left(\frac{(D^m(\mu, \beta, t)f_n(z))}{z} \right) \\ &= \alpha_1 [\ln(D^m(\mu, \beta, t)f_1(z)) - \ln z] + \dots + \alpha_n [\ln(D^m(\mu, \beta, t)f_n(z)) - \ln z]. \end{aligned} \quad (3.4)$$

By differentiating the above equality logarithmical, we have

$$\frac{h''(z)}{h'(z)} = \alpha_1 \left[\frac{(D^m(\mu, \beta, t)f_1(z))'}{D^m(\mu, \beta, t)f_1(z)} - \frac{1}{z} \right] + \dots + \alpha_n \left[\frac{(D^m(\mu, \beta, t)f_n(z))'}{D^m(\mu, \beta, t)f_n(z)} - \frac{1}{z} \right]. \quad (3.5)$$

Then

$$\frac{zh''(z)}{h'(z)} = \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right]. \quad (3.6)$$

So by the condition of the Theorem, we find

$$\begin{aligned} \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \right) \\ &\leq \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \leq \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \leq 1 \end{aligned} \tag{3.7}$$

since $\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \leq 1$, and $\Re(\sigma) \geq \sum_{j=1}^n |\alpha_j|$. Finally by applying Lemma 2.2, for the function $h(z)$, we proved that $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)(z) \in S$. \square

Remark 3.2. If we set $\beta = \mu, \sigma = 1$ in Theorem 3.1, then we have [15, Theorem 2.3].

Remark 3.3. If we set $m = 0, \sigma = 1$ in Theorem 3.1, then we have [6, Theorem 1].

Corollary 3.4. Let $\alpha_j > 0, \sigma \in C, m \in N_0, t \geq 0, \beta \geq 0, 0 \leq \mu \leq \beta$, and each of the functions $f_j \in A, j = 1, 2, \dots, n$. If $f_j \in A, j = 1, 2, \dots, n$, satisfies the inequality (3.1) and $\Re(\sigma) \geq \sum_{i=1}^n \alpha_j$, then the integral operator $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)$ defined by (1.4) is in the univalent class S .

Remark 3.5. If we set $\beta = \mu, \sigma = 1$ in Corollary 3.4, we get [15, Corollary 2.5].

Theorem 3.6. Let $M_j \geq 1$ and suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$ satisfies the inequality

$$\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| \leq 1, \quad z \in U, m \in N_0. \tag{3.8}$$

Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq \sum_{i=1}^n |\alpha_j| (2M_j + 1) > 0$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M_j, z \in U, j = 1, 2, \dots, n$, then the integral operator $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)$ defined by (1.4) is in the univalent function class S .

Proof . We know from the proof of Theorem 3.1 that

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \right). \tag{3.9}$$

So by the imposed condition, we find

$$\frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \left| \frac{zh''(z)}{h'(z)} \right| \leq \frac{1 - |z|^{2\Re(\rho)}}{\Re(\rho)} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} \right| + 1 \right) \tag{3.10}$$

By Schwarz Lemma, we get $|D^m(\mu, \beta, t)g_i| \leq M_j |z|, R = 1$, therefore

$$\begin{aligned} \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} \right| M_j + 1 \right) \\ &\leq \frac{1 - |z|^{2\Re(\sigma)}}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| M_j + M_j + 1 \right) \\ &\leq \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| (2M_j + 1) \leq 1 \end{aligned} \tag{3.11}$$

since $\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| \leq 1$ and $\Re(\sigma) \geq \sum_{j=1}^n |\alpha_j| (2M_j + 1)$. By applying Lemma 2.2, for function $h(z)$, we prove that $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n) \in S$. \square

Corollary 3.7. Let $M_j \geq 1, \alpha_j > 0$ and suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$, satisfies the inequality (3.8). Also, let $\alpha, \sigma \in C$ with $\Re(\sigma) \geq \sum_{i=1}^n |\alpha_i| (2M_i + 1) > 0$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M_j, z \in U, j = 1, \dots, n$, then for any complex number σ with $\Re(\sigma) \geq \Re(\alpha)$, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Corollary 3.8. Let $M \geq 1$ and suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$, satisfies the inequality (3.8). Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq (2M + 1) \sum_{i=1}^n |\alpha_i| > 0$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M, z \in U, j = 1, 2, \dots, n$, then for any complex number σ with $\Re(\sigma) \geq \Re(\alpha)$, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Corollary 3.9. Suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$ satisfies the inequality (3.8). Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq 3 \sum_{i=1}^n |\alpha_i| > 0$. If $|D^m(\mu, \beta, t)f_j(z)| \leq 1, z \in U, j = 1, 2, \dots, n$, then for any complex number σ with $\Re(\sigma) \geq \Re(\alpha)$, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (3.10) is in the univalent function class S .

Remark 3.10. In Corollary 3.9

- (i) If we set $\sigma = 1, \beta = \mu$, then we obtain Theorem 2.6.
- (ii) If we set $\sigma = 1, \alpha_j > 0, j = 1, 2, \dots, n$, then we have [15, Corollary 2.8].

Theorem 3.11. Suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$, satisfies the inequality

$$\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \leq 1, \quad z \in U. \quad (3.12)$$

Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$, with $\Re(\sigma) \geq \sum_{i=1}^n |\alpha_i| > 0$, and let $c \in C$ be such that $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j|$, then the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent class S .

Proof . We know from the proof of Theorem 3.1 that

$$\frac{zh''(z)}{h'(z)} = \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right]. \quad (3.13)$$

We have that

$$c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} = c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{1}{\sigma} \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right]. \quad (3.14)$$

Then

$$\begin{aligned} \left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| &= \left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{1}{\sigma} \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right] \right| \\ &\leq |c| + \left| \frac{1 - |z|^{2\sigma}}{\sigma} \right| \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \right) \\ &\leq |c| + \frac{1}{|\sigma|} \sum_{j=1}^n |\alpha_j| \leq |c| + \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \leq 1 \end{aligned} \quad (3.15)$$

since $\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \leq 1, \Re(\sigma) \geq \sum_{j=1}^n |\alpha_j|$ and $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j|$. Finally, applying Lemma 2.3 for the function $h(z)$, we prove that $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) \in S$. \square

Corollary 3.12. Suppose that each of the functions $f_j \in A, j \in \{1, \dots, n\}$, satisfies the inequality (3.12). Also, let $\alpha_j > 0, \sigma \in C$, with $\Re(\sigma) \geq \sum_{i=1}^n \alpha_i$, and let $c \in C$ be such that $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n \alpha_j$, then the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent class S .

Theorem 3.13. Let $M_j \geq 1$ and suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$, satisfies the inequality

$$\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| \leq 1, \quad z \in U, m \in N_0. \tag{3.16}$$

Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq \sum_{i=1}^n |\alpha_j|(2M_j + 1) > 0$, and let $c \in C$ be such that $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j|(2M_j + 1)$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M_j \quad (z \in U, j = 1, 2, \dots, n)$, then, for any complex number σ with $\Re(\sigma) \geq \Re(\alpha)$, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Proof . We know from the proof of Theorem 3.1 that

$$\frac{zh''(z)}{h'(z)} = \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right]. \tag{3.17}$$

Therefore,

$$c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} = c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{1}{\sigma} \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right]. \tag{3.18}$$

Then

$$\begin{aligned} \left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| &= \left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{1}{\sigma} \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right] \right| \\ &\leq |c| + \left| \frac{1 - |z|^{2\sigma}}{\sigma} \right| \sum_{j=1}^n |\alpha_j| \left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right| \\ &\leq |c| + \left| \frac{1 - |z|^{2\sigma}}{\sigma} \right| \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} \right| + 1 \right). \end{aligned} \tag{3.19}$$

By Schwarz Lemma, we get $|D^m(\mu, \beta, t)g_i| \leq M_j |z|, R = 1$, therefore

$$\begin{aligned} \left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| &\leq |c| + \frac{1}{|\sigma|} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} \right| M_j + 1 \right) \\ &\leq |c| + \frac{1}{|\sigma|} \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| M_j + M_j + 1 \right) \\ &\leq |c| + \frac{1}{|\sigma|} \sum_{j=1}^n |\alpha_j| (2M_j + 1) \\ &\leq |c| + \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| (2M_j + 1) \leq 1, \end{aligned} \tag{3.20}$$

since $\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| \leq 1$ and $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j|(2M_j + 1)$. Finally, applying Lemma 2.3 for function $h(z)$, we proved that $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) \in S$. \square

Corollary 3.14. Let $M_j \geq 1, \alpha_j > 0$ and suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$ satisfies the inequality (3.16). Also, let $\sigma \in C$ with $\Re(\sigma) \geq \sum_{i=1}^n \alpha_j(2M_j + 1) > 0$, and let $c \in C$ be such that $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n \alpha_j(2M_j + 1)$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M_j, z \in U, j = 1, \dots, n$, then the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Corollary 3.15. Let $M \geq 1$, and suppose that each of the functions $f_j \in A, j = 1, \dots, n$ satisfies the inequality (3.16) Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq (2M + 1) \sum_{i=1}^n |\alpha_j| > 0$, and let $c \in C$ be such that $|c| \leq 1 - \frac{(2M_j+1)}{\Re(\sigma)} \sum_{j=1}^n \alpha_j$. If $|D^m(\mu, \beta, t)f_j(z)| \leq M, z \in U, j = 1, 2, \dots, n$, then the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Corollary 3.16. Suppose that each of the functions $f_j \in A, j = 1, 2, \dots, n$ satisfies the inequality (3.16). Also, let $\alpha_1, \dots, \alpha_n, \sigma \in C$ with $\Re(\sigma) \geq 3 \sum_{i=1}^n |\alpha_i| > 0$, and let $c \in C$ be such that $|c| \leq 1 - \frac{3}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j|$. If $|D^m(\mu, \beta, t)f_j(z)| \leq 1, z \in U, j = 1, 2, \dots, n$, then the integral operator $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Theorem 3.17. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M_j, M_j \geq 1$, for all $j = 1, 2, \dots, n, D^m(\mu, \beta, t)f_j(z) \in S(p_j)$, for $j = 1, 2, \dots, n$, with $\alpha_1, \dots, \alpha_n, \sigma \in C$ such that

$$\Re(\sigma) \geq \sum_{j=1}^{\infty} |\alpha_j| \frac{[(2M_j - 1)p_j + 2] M_j}{2M_j - 1}.$$

Also, let $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \frac{[(2M_j - 1)p_j + 2] M_j}{2M_j - 1}$, then the integral operator $I_{t, \beta, \mu}^{m, \sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S .

Proof . Since $D^m(\mu, \beta, t)f_j(z) \in S(p_j)$, so by Lemma 2.4, we have

$$\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| \leq p_j |z|^2, \quad z \in U \quad (3.21)$$

From $|D^m(\mu, \beta, t)f_j(z)| \leq M_j$, for any $j = 1, \dots, n$ and by using Lemma 2.1, we get $|D^m(\mu, \beta, t)f_j(z)| \leq M_j |z|, R = 1$. From the proof of Theorem 3.1, we have that

$$\frac{zh''(z)}{h'(z)} = \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right] \quad (3.22)$$

and so

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} \right| + 1 \right). \quad (3.23)$$

By Schwarz Lemma, we get $|D^m(\mu, \beta, t)g_i| \leq M_j |z|, R = 1$, therefore

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} \right| M_j + 1 \right) \quad (3.24)$$

and

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| M_j + M_j + 1 \right). \quad (3.25)$$

Thus,

$$\begin{aligned} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \sum_{j=1}^n |\alpha_j| \left(p_j |z|^2 M_j + M_j + 1 \right) \leq \sum_{j=1}^n |\alpha_j| (p_j M_j + 2M_j) \\ &\leq \sum_{j=1}^n |\alpha_j| (p_j M_j + 2M_j + 4M_j^2 + 8M_j^3 + \dots) \end{aligned} \quad (3.26)$$

because $2M_j, 4M_j^2, 8M_j^3, \dots > 1$. Which implies that

$$\begin{aligned} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \sum_{j=1}^n |\alpha_j| \left(p_j M_j + \frac{2M_j}{2M_j - 1} \right) \\ &\leq \sum_{j=1}^n |\alpha_j| \frac{[(2M_j - 1)p_j + 2] M_j}{2M_j - 1}. \end{aligned} \quad (3.27)$$

Now, from

$$\left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| \leq |c| + \frac{1}{|\sigma|} \left| \frac{zh''(z)}{h'(z)} \right| \leq |c| + \frac{1}{\Re(\sigma)} \left| \frac{zh''(z)}{h'(z)} \right| \tag{3.28}$$

This implies

$$\left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| \leq |c| + \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \frac{[(2M_j - 1)p_j + 2] M_j}{2M_j - 1} \leq 1 \tag{3.29}$$

Since $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \frac{[(2M_j - 1)p_j + 2] M_j}{2M_j - 1}$, by Lemma 2.3, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is in the univalent function class S . \square

Corollary 3.18. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M, M_j = M \geq 1$, for $j = 1, 2, \dots, n, D^m(\mu, \beta, t)f_j(z) \in S(p_j)$, for $j = 1, 2, \dots, n$, with

$$\Re(\sigma) \geq \sum_{j=1}^{\infty} |\alpha_j| \frac{[(2M - 1)p_j + 2] M}{2M - 1}$$

where σ, α_j are complex numbers. If $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| \frac{[(2M - 1)p_j + 2] M}{2M - 1}$, then $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) \in S$.

Corollary 3.19. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M, M_j = M \geq 1$, for $j = 1, 2, \dots, n, D^m(\mu, \beta, t)f_j(z) \in S(p_j)$, for $j = 1, 2, \dots, n, |\alpha_j| = |\alpha|$ with

$$\Re(\sigma) \geq |\alpha| \sum_{j=1}^{\infty} \frac{[(2M - 1)p_j + 2] M}{2M - 1}$$

where σ, α are complex numbers. If $|c| \leq 1 - \frac{|\alpha|}{\Re(\sigma)} \sum_{j=1}^n \frac{[(2M - 1)p_j + 2] M}{2M - 1}, M \geq 1$, then by Lemma 2.3 the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) \in S$.

Theorem 3.20. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M_j, M_j \geq 1$, for all $(j = 1, \dots, n), D^m(\mu, \beta, t)f_j(z) \in K_{2,\delta_j}$ for $(j = 1, \dots, n)$, such that

$$\Re(\sigma) \geq \sum_{j=1}^{\infty} |\alpha_j| [(1 - \delta_j) + (n(n + 1))/2] M_j$$

where σ, α_j are complex numbers. If $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| [(1 - \delta_j) + (n(n + 1))/2] M_j, M_j \geq 1$ then, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is univalent in U .

Proof . Using the proof of Theorem 3.1, we have

$$\frac{zh''(z)}{h'(z)} = \sum_{j=1}^n \alpha_j \left[\frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} - 1 \right] \tag{3.30}$$

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z(D^m(\mu, \beta, t)f_j(z))'}{D^m(\mu, \beta, t)f_j(z)} \right| + 1 \right) \tag{3.31}$$

By Schwarz Lemma, we get $|D^m(\mu, \beta, t)g_i| \leq M_j |z|, R = 1$, therefore

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} \right| M_j + 1 \right) \tag{3.32}$$

and

$$\left| \frac{zh''(z)}{h'(z)} \right| \leq \sum_{j=1}^n |\alpha_j| \left(\left| \frac{z^2(D^m(\mu, \beta, t)f_j(z))'}{(D^m(\mu, \beta, t)f_j(z))^2} - 1 \right| M_j + M_j + 1 \right). \quad (3.33)$$

Thus,

$$\begin{aligned} \left| \frac{zh''(z)}{h'(z)} \right| &\leq \sum_{j=1}^n |\alpha_j| ((1 - \delta_j)M_j + M_j + 1) \leq \sum_{j=1}^n |\alpha_j| ((1 - \delta_j)M_j + 2M_j) \\ &\leq \sum_{j=1}^n |\alpha_j| ((1 - \delta_j)M_j + M_j + 2M_j + 3M_j + \dots + nM_j) \\ &\leq \sum_{j=1}^n |\alpha_j| [(1 - \delta_j)M_j + (n(n + 1)/2) M_j]. \end{aligned} \quad (3.34)$$

Now, we evaluate the expression

$$\left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| \leq |c| + \frac{1}{|\sigma|} \left| \frac{zh''(z)}{h'(z)} \right| \leq |c| + \frac{1}{\Re(\sigma)} \left| \frac{zh''(z)}{h'(z)} \right|. \quad (3.35)$$

Then

$$\left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| \leq |c| + \frac{1}{|\sigma|} \left| \frac{zh''(z)}{h'(z)} \right| \leq |c| + \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| [(1 - \delta_j)M_j + (n(n + 1)/2) M_j]$$

This implies that

$$\left| c|z|^{2\sigma} + (1 - |z|^{2\sigma}) \frac{zh''(z)}{\sigma h'(z)} \right| \leq 1.$$

Since $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| [(1 - \delta_j) + (n(n + 1)/2)] M_j$ by Lemma 2.3, the integral operator, $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z) \in S$. \square

Corollary 3.21. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M, M \geq 1$ for all $(j = 1, 2, \dots, n)$, $D^m(\mu, \beta, t)f_j(z) \in K_{2,\delta_j}$, for $j = 1, 2, \dots, n$, such that

$$\Re(\sigma) \geq \sum_{j=1}^{\infty} |\alpha_j| [(1 - \delta_j) + (n(n + 1))/2] M$$

where σ, α_j are complex numbers. If $|c| \leq 1 - \frac{1}{\Re(\sigma)} \sum_{j=1}^n |\alpha_j| [(1 - \delta_j) + (n(n + 1))/2] M, M \geq 1$ then, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is univalent in U .

Corollary 3.22. Let c be a complex number, $|D^m(\mu, \beta, t)f_j(z)| \leq M, M \geq 1, \alpha_1 = \dots = \alpha_n = \alpha$ for all $(j = 1, 2, \dots, n)$, $D^m(\mu, \beta, t)f_j(z) \in K_{2,\delta}$ for $j = 1, \dots, n$, such that

$$\Re(\sigma) \geq |\alpha| \sum_{j=1}^{\infty} [(1 - \delta_j) + (n(n + 1))/2] M$$

where σ, α_j are complex numbers. If $|c| \leq 1 - \frac{|\alpha|}{\Re(\sigma)} \sum_{j=1}^n [(1 - \delta_j) + (n(n + 1))/2] M, M \geq 1$ then, the integral operator $I_{t,\beta,\mu}^{m,\sigma}(f_1, \dots, f_n)(z)$ defined by (1.4) is univalent in class S .

Conclusion

In conclusion, a new generalized integral operator is defined in the unit disk U and conditions for univalence of the integral operators in U are investigated. Results obtained generalize some existing results.

Acknowledgments

The authors sincerely appreciate the efforts of the reviewers and the editor for their suggestions that added values to this paper.

References

- [1] F.M. Al-Oboudi, *On univalent functions defined by a Sălăgean differential operator*, Int. J. Math. Math. Sci. **2004** (2004), 1429–1436.
- [2] J.W. Alexander, *Functions which map the interior of the unit circle upon simple regions*, Ann. Math. **17** (1915), no. 1, 12–22.
- [3] C. Barbatu and D. Breaz, *Univalence criteria for some general integral operators*, An. St. Univ. Ovidius Const. **29** (2021), no. 1, 37–52.
- [4] S.D. Bernardi, *Convex and starlike univalent functions*, Trans. Amer. Math. Soc. **135** (1969), 429–446.
- [5] D. Breaz, N. Breaz, and H.M. Srivastava, *An extension of the univalent condition for a family of integral operators*, Appl. Math. Lett. **22** (2009), no. 1, 41–44.
- [6] D. Breaz and N. Breaz, *Two integral operators*, Studia Univ. Babeş-Bolyai Math. **47** (2002), 13–20.
- [7] D. Breaz and N. Breaz, *Univalence of an integral operator*, Mathematica **70** (2005), 35–38.
- [8] D. Breaz and N. Breaz, *An integral Univalent operator*, Acta Math. Univ. Comenianae. New Ser. **76** (2007), no. 2, 137–142.
- [9] S. Bulut, *Univalence preserving integral operators defined by generalized Al-Oboudi differential Operator*, An St. Univ. Ovidius Constanta **17** (2009), no. 1, 37–50.
- [10] S. Bulut, *Univalence condition for a new generalization of the family of integral operators*, Acta Univ. Apulensis Math. Inf. **18** (2009), 71–78.
- [11] S. Bulut, *A new univalent integral operator defined by Al-Oboudi differential operator*, Gen. Math. **18** (2010), no. 2, 85–93.
- [12] S. Bulut, *An integral univalent operator defined by generalized Al-Oboudi differential operator on the classes $T_j, T_{j,\mu}, S_j(p)$* , Novi Sad J. Math. **40** (2010), no. 1, 43–53.
- [13] C. Barbatu and D. Breaz, *Some Univalence conditions of a certain general integral operator*, Eur. J. Pure Appl. Math. **13** (2020), no. 5, 1285–1299.
- [14] S. Bulut and D. Breaz, *Univalency and convexity conditions for a general integral operator*, Chin. J. Math. **2014** (2014), 4 pages.
- [15] S. Bulut, *Sufficient conditions for univalence of an integral operator defined by Al-Oboudi differential operator*, J. Inequal. Appl. **2008** (2008), 5 pages.
- [16] E. Deniz, D. Raducanu, and H. Orhan, *On the univalence of an integral operator defined by Hadamard product*, Appl. Math. Lett. **24** (2012), 179–184.
- [17] I. Faisal and M. Darus, *A study of Ahlfors' univalence criteria for a space of analytic functions: Criteria II*, Math. Comput. Model. **55** (2012), 1466–1470.
- [18] R.J. Libera, *Some classes of regular univalent functions*, Proc. Amer. Math. Soc. **16** (1965), 755–758.
- [19] S.S. Miller, P.T. Mocanu, and M.O. Reade, *Bezilevic functions and generalized convexity*, Rev. Roumaine. Math. Pure Appl. **19** (1974), 213–224.
- [20] S.S. Miller, P.T. Mocanu and M.O. Reade, *Starlike integral operators*, Pacific J. Math. **79** (1978), no. 1, 157–168.
- [21] Z. Nehari, *Conformal Mapping*, Dover, New York, NY, USA, 1975.
- [22] T.O. Opoola, *On a subclass of univalent functions defined by a generalized differential operator*, Int. J. Math. Anal. **11** (2017), no.18, 869–876.

-
- [23] G.I. Oros, G. Oros, and D. Breaz, *Sufficient conditions for univalence of an integral operator*, J. Inequal. Appl. **2008** (2008), 7 pages.
- [24] A. Oprea, D. Breaz, and H.M. Srivastava, *Univalence conditions for a new family of integral operators*, Filomat **30** (2016), no. 5, 1243–1251.
- [25] A. Oprea and D. Breaz, *Univalence conditions for a general integral operator*, An. St. Univ. Ovidius Constanta **23** (2015), no. 1, 213–224.
- [26] V. Pescar, *A new generalization of Ahlfors's and Becker's criterion of univalence*, Bull. Malays. Math. Soc. (Ser. 2) **19** (1996), 53–54.
- [27] V. Pescar, *New criteria for Univalence of certain integral operators*, Demonstr. Math. **33** (2000), 51–54.
- [28] V. Pescar, *On the Univalence of some integral operators*, J. Indian Acad. Math. **27** (2005), 239–243.
- [29] N. Pascu, *An improvement on Becker's univalence criterion*, Proc. Commem. Session Stoilow, Brasov, 1987, pp. 43–48.
- [30] G.S. Sălăgean, *Subclasses of Univalent Functions*, Complex Anal. Fifth Roman. Seminar, part I (Brucharest, 1983), Lecture Notes in Mathematics, vol. 1013, Springer, Berlin, 1983, pp 362–372.
- [31] N. Seenivasagan and D. Breaz, *Certain sufficient conditions for univalence*, Gen. Math. **15** (2007), no. 4, 7–15.
- [32] V. Singh, *On class of univalent functions*, Int. J. Maths. Math Soc. **23** (2000).