

On fuzzy C^* -algebras

Ali Reza Khoddami*, Fateme Mohammadi

Faculty of Mathematical Sciences, Shahrood University of Technology, P. O. Box 3619995161-316, Shahrood, Iran

(Communicated by Abasalt Bodaghi)

Abstract

In this paper, we show that there is no fuzzy C^* -algebra norm according to the definitions contained in [1, 2, 3]. Based on our claim, some parts of [Theorems 2 and 3, 2], [Corollaries 1 and 2, 2] and [Theorem 4, 3] are incorrect.

Keywords: Fuzzy normed space, Fuzzy normed algebra, Fuzzy normed $*$ -algebra, Fuzzy C^* -algebra norm
2020 MSC: Primary 46S40; Secondary 03E72

1 Introduction

We are well aware that the definition of an induced fuzzy C^* -algebra was introduced by Park et al. at the first time [4]. As far as we know, the definition of a fuzzy C^* -algebra norm was introduced by Ansari et al. at the first time [1]. This definition followed by Lo'lo' et al. [2]. Also some results according to this definition were presented by Movahednia and De la Sen [3]. This paper aims to inform the authors and the readers to avoid following an incorrect definition.

Definition 1.1 ([1, 2, 3, 4]). Let X be a real vector space. A function $N : X \times \mathbb{R} \rightarrow [0, 1]$ is called a fuzzy norm on X if for all $x, y \in X$ and all $s, t \in \mathbb{R}$

1. $N(x, t) = 0$ for all $t \leq 0$,
2. $N(x, t) = 1$ for all $t > 0$ if and only if $x = 0$,
3. $N(cx, t) = N(x, \frac{t}{|c|})$ for all $c \neq 0$,
4. $N(x + y, s + t) \geq \min(N(x, s), N(y, t))$,
5. $N(x, \cdot)$ is a non-decreasing function on \mathbb{R} and $\lim_{t \rightarrow \infty} N(x, t) = 1$,
6. For $x \neq 0$, $N(x, \cdot)$ is continuous on \mathbb{R} .

The pair (X, N) is called a fuzzy normed vector space.

Definition 1.2 ([1, 2, 3, 4]). Let (X, N) be a fuzzy normed vector space.

1. A sequence $\{x_n\}$ in X is said to be convergent if there exists an $x \in X$ such that

$$\lim_{n \rightarrow \infty} N(x_n - x, t) = 1$$

for all $t > 0$.

*Corresponding author

Email addresses: khoddami.alireza@shahroodut.ac.ir (Ali Reza Khoddami), fa.mohammadi36@gmail.com (Fateme Mohammadi)

2. A sequence $\{x_n\}$ in X is called Cauchy if for each $\epsilon > 0$ and each $t > 0$ there exists an $n_0 \in \mathbb{N}$ such that for all $n > n_0$ and all $p \geq 1$, we have $N(x_{n+p} - x_n, t) > 1 - \epsilon$. If each Cauchy sequence is convergent, then the fuzzy norm is said to be complete and the fuzzy normed space is called a fuzzy Banach space.

Definition 1.3 ([1, 2, 3, 4]). Let X be an algebra and (X, N) a fuzzy normed space. The fuzzy normed space (X, N) is called a fuzzy normed algebra if

$$N(xx', st) \geq N(x, s)N(x', t)$$

for all $x, x' \in X, s, t \in \mathbb{R}^+$. A complete fuzzy normed algebra is called a fuzzy Banach algebra.

Definition 1.4 ([1, 2, 3]). Let A be an algebra over \mathbb{C} . An involution on A is a mapping

$$\begin{aligned} * : A &\longrightarrow A \\ a &\longrightarrow a^* \end{aligned}$$

such that

1. $(\alpha a + \beta b)^* = \bar{\alpha}a^* + \bar{\beta}b^*$ for all $a, b \in A$ and $\alpha, \beta \in \mathbb{C}$.
2. $(ab)^* = b^*a^*$ for all $a, b \in A$.
3. $(a^*)^* = a$ for all $a \in A$.

A complex algebra with an involution is a $*$ -algebra.

Definition 1.5 ([1, 2, 3, 4]). Let A be a $*$ -algebra and (A, N) a fuzzy normed algebra. The fuzzy normed algebra (A, N) is called a fuzzy normed $*$ -algebra if,

$$N(a^*, t) = N(a, t)$$

for all $a \in A$ and for all $t \in \mathbb{R}^+$. A complete fuzzy normed $*$ -algebra is called a fuzzy Banach $*$ -algebra.

Definition 1.6 ([1, 2, 3]). Let (A, N) be a fuzzy Banach $*$ -algebra. Then (A, N) is called a fuzzy C^* -algebra if,

$$N(a^*a, st) = N(a^*, s)N(a, t)$$

for all $a \in A$ and for all $s, t \in \mathbb{R}^+$.

Theorem 1.7. There is no a fuzzy C^* -algebra norm according to Definition 1.6.

Proof . We suppose by contradiction that N is a fuzzy C^* -algebra norm on A in the sense Definition 1.6. Let $0 \neq a \in A$. Since $\lim_{t \rightarrow \infty} N(a^*a, t) = 1$, there exists an $M > 0$ such that $N(a^*a, t) > \frac{1}{2}$ for all $t \geq M$. So by Definition 1.6, for all $n \in \mathbb{N}$ we have, $\frac{1}{2} < N(a^*a, M) = N(a^*a, \frac{M}{n} \cdot n) = N(a^*, \frac{M}{n})N(a, n)$. Hence

$$\frac{N(a^*a, M)}{N(a, n)} = N(a^*, \frac{M}{n}) = N(a, \frac{M}{n})$$

for all $n \in \mathbb{N}$. By parts (6) and (5) of Definition 1.1, $N(a, \cdot)$ is continuous at $t = 0$ and $\lim_{n \rightarrow \infty} N(a, n) = 1$. So we can conclude that $N(a^*a, M) = \lim_{n \rightarrow \infty} \frac{N(a^*a, M)}{N(a, n)} = \lim_{n \rightarrow \infty} N(a, \frac{M}{n}) = 0$. Hence $N(a^*a, M) = 0$, that is a contradiction. \square

Example 1.8. Let $(A, *, \|\cdot\|)$ be a C^* -algebra. Define $N : A \times \mathbb{R} \longrightarrow [0, 1]$ by

$$N(a, t) = \begin{cases} 0, & t \leq 0 \\ \frac{t}{t + \|a\|}, & t > 0. \end{cases}$$

Then (A, N) is a fuzzy Banach $*$ -algebra but $N(a^*a, 2 \times 3) \neq N(a^*, 2)N(a, 3)$.

References

- [1] N. Ansari, M.H. Hooshmand, M.E. Gordji, and K. Jahedi, *Stability of fuzzy orthogonally $*$ – n –derivation in orthogonally fuzzy C^* –algebras*, Int. J. Nonlinear Anal. Appl. **12** (2021), no. 1, 533–540.
- [2] P. Lóló, E. Movahednia, and M. De la Sen, *Hyers stability and multi-fuzzy Banach algebra*, Mathematics **10** (2022), 1–13.
- [3] E. Movahednia and M. De la Sen, *A new approach to involution in fuzzy C^* –algebra via functional inequality and Python implementation*, Axioms **12** (2023), 1–14.
- [4] C. Park, J.R. Lee, T.M. Rassias, and R. Saadati, *Fuzzy $*$ –homomorphisms and fuzzy $*$ –derivations in induced fuzzy C^* –algebras*, Math. Comput. Model. **54** (2011), 2027–2039.