

# An algebraic method to obtain analytical solutions for a class of fractional partial differential equations

Ameneh Manouchehrian, Ahmad Haghbin\*

Department of Mathematics, Gorgan Branch, Islamic Azad University, Gorgan, Iran

(Communicated by Mugur Alexandru Acu)

---

## Abstract

In this study, we apply an algebraic approach to solve a class of fractional partial differential equations (FPDEs) by utilizing conformable fractional derivatives. This method has been successfully utilized to study and achieve analytical solutions for Drinfeld-Sokolov-Wilson equations. In this approach, we use a suitable fractional transformation and the principles of fractional calculus to simplify fractional partial differential equations into ordinary differential equations.

Keywords: fractional differential equations, Drinfeld-Sokolov-Wilson equations, conformable fractional derivative, analytical solution

2020 MSC: 26A33, 26D15, 34A08, 34A12

---

## 1 Introduction

In recent decades, fractional differential equations (FDEs) have been used widely to model physical phenomena, especially uncommon phenomena and complex natural processes that cannot be efficiently described by classical calculus [15, 18]. These equations have applications in various fields, such as mathematical biology, fluid mechanics, nonlinear optics, image processing, plasma physics and so on. We need to find solutions for these equations in order to research and describe these phenomena. It is difficult to solve FDEs. Hence, numerous researchers have developed and implemented numerical and analytical techniques to solve these equations in recent years. Some of these techniques are such as, Finite difference method [1], Adomian decomposition method [3], differential transform method [16], variational iteration method [5, 8],  $\frac{G'}{G^2}$ -expansion method [11], Fan-sub equation method [12], Lie group method [9], exp-expansion method [4], first integral method [2] and so on. In this paper, we study a class of fractional partial differential equations, namely Drinfeld-Sokolov-Wilson (DSW) in the following form [10]:

$$D_t^\alpha u + n v D_x^\beta v = 0, \tag{1.1}$$

$$D_t^\alpha v + q D_x^{3\alpha} v + r u D_x^\beta v + s v D_x^\beta u = 0. \tag{1.2}$$

where  $p, q, r$ , and  $s$  are non-zero parameters. Also,  $D_t^\alpha$  and  $D_x^\beta$  represent fractional partial derivative operators, as described in section 2. When  $\alpha = \beta = 1$ , equations (1.1) and (1.2) are called classical DSW equations. These equations play an important role in modeling shallow water flow and fluid dynamics. Equations (1.1) and (1.2) were

---

\*Corresponding author

Email addresses: [manouchehriana@gmail.com](mailto:manouchehriana@gmail.com) (Ameneh Manouchehrian), [ah.haghbin@iaua.ac.ir](mailto:ah.haghbin@iaua.ac.ir) (Ahmad Haghbin)

first introduced by Drinfeld and Sokolov and then, studied by Wilson. Researchers have utilized some techniques to study this equation such as, Galerkin method [20], F-expansion method [6], simplest equation method[14] and so on. Our aim in this paper is to find analytical solutions for the DSW equations using an algebraic method, as described in section 3.

The rest of our work is organized as follows. In section 2, we present definition of the conformable derivative with its properties. Description of method and its applications to the space-time fractional differential equations are described in section 3. Then the mentioned method is applied to the DSW equations in section 4. Graphical presentation of some solutions are shown in section 5 Discussion and conclusions are presented in section 6.

## 2 Definition of the conformable derivative with its properties

In this section, we illustrate the definition of the conformable fractional derivative(CFD) and some its important properties of order  $\gamma$  with respect to the independent variable  $\zeta$  as follows[13].

For a function  $h : [0, \infty] \rightarrow \mathbb{R}$ , the CFD of  $h$  of order  $\alpha$  is defined by

$$D^\gamma h(\zeta) = \lim_{\eta \rightarrow 0} \frac{h(\zeta + \eta \zeta^{1-\gamma}) - h(\zeta)}{\eta}. \quad (2.1)$$

Some well-known properties to this newly defined fractional derivative are as follows. If  $g$  and  $h \neq 0$  be two functions  $\gamma$ -differentiable,  $\gamma \in (0, 1]$  and  $a, b \in \mathbb{R}$ . Then, we have

- (1)  $D^\gamma \{ag(\zeta) + bh(\zeta)\} = aD^\gamma g(\zeta) + bD^\gamma h(\zeta)$ ,
- (2)  $D^\gamma \{g(\zeta)h(\zeta)\} = g(\zeta)D^\gamma h(\zeta) + h(\zeta)D^\gamma g(\zeta)$ ,
- (3)  $D^\gamma \left\{ \frac{g(\zeta)}{h(\zeta)} \right\} = \frac{h(\zeta)D^\gamma g(\zeta) - g(\zeta)D^\gamma h(\zeta)}{h^2(\zeta)}$ ,
- (4)  $D^\alpha C = 0$ , for all constant functions  $f(z) = C$ ,
- (5)  $D^\gamma(g)(\zeta) = \zeta^{1-\gamma} \frac{dg}{d\zeta}$ .

Also, CFD of some special functions are as follows.

- (a)  $D^\gamma(\zeta^r) = r\zeta^{r-\gamma}$  for all  $r \in \mathbb{R}$ ,
- (b)  $D^\gamma(1) = 0$ ,
- (c)  $D^\gamma(e^{c\zeta}) = c\zeta^{1-\gamma}e^{c\zeta}$ ,  $c \in \mathbb{R}$ ,
- (d)  $D^\gamma(\sin b\zeta) = b\zeta^{1-\gamma} \cos b\zeta$ ,  $b \in \mathbb{R}$ ,
- (e)  $D^\gamma(\cos b\zeta) = -b\zeta^{1-\gamma} \sin b\zeta$ ,  $b \in \mathbb{R}$ ,
- (f)  $D^\gamma\left(\frac{1}{\gamma}\zeta^\gamma\right) = 1$ .

The proofs of these properties can be seen in[13]. Let  $\gamma \in (n, n+1]$ , and  $h$  be an  $\gamma$ -differentiable att  $t > 0$ . Then the CFD of  $h$  of order  $\gamma$  is defined as

$$D^\gamma(h(t)) = \lim_{\eta \rightarrow 0} \frac{h^{([\gamma]-1)}(t + \eta t^{([\gamma]-\gamma)}) - h^{([\gamma]-1)}(t)}{\eta}. \quad (2.2)$$

where  $[\gamma]$  is the smallest integer greater than or equal to  $\gamma$ .

## 3 Description of method and its applications to the space-time FDEs

In this section, we outline the main steps of this method for solving FDEs. For a given FDE in two variables  $x$  and  $t$  we have

$$P(u, u_x, u_t, D_t^\alpha u, D_x^\alpha u, \dots) = 0, \quad 0 < \alpha < 1, \quad (3.1)$$

where  $D_t^\alpha u$  and  $D_x^\alpha u$  are CFDs of  $u$ ,  $u = u(x, t)$  is an unknown function and  $P$  is a polynomial in  $u$  and its various partial derivatives, in which the highest order derivatives and nonlinear terms are involved. We take the travelling wave transformation

$$\eta = \frac{\omega}{\beta} x^\beta - \frac{\lambda}{\alpha} t^\alpha, \quad (3.2)$$

where  $\omega$  and  $\lambda$  are nonzero constants to be determined later. Substituting (3.2) into (3.1), we reduce (3.1) to the following ordinary differential equation

$$\tilde{N}(U, U', U'', U''', \dots) = 0. \quad (3.3)$$

Here prime denotes the derivative with respect to  $\eta$ . Exact solutions for this equation can be constructed as a finite series

$$\Phi(\xi) = \frac{\sum_{j=0}^{\eta_1} A_j \Theta(\xi)}{\sum_{j=0}^{\eta_2} B_j \Theta(\xi)}, \quad (3.4)$$

where the positive constants  $\eta_1$  and  $\eta_2$  can be calculated by considering the homogeneous balance between the highest order derivatives and the highest nonlinear terms of  $\Phi(\xi)$  in equation (3.3), and  $A_j$  ( $0 \leq j \leq \eta_1$ ),  $B_j$  ( $0 \leq j \leq \eta_2$ ) are constants to be found later and  $A_{\eta_1}, B_{\eta_2} \neq 0$ . Here  $\Theta = \Theta(\xi)$  satisfies the following ODE

$$\Theta'(\xi) = p + \Theta^2(\xi) \quad (3.5)$$

where  $p$  is a constant and which has the following special solutions [17].

Case1: When  $p < 0$ ,

$$\Theta_1(\xi) = -\sqrt{-p} \tanh(\sqrt{-p} \xi), \quad (3.6)$$

$$\Theta_2(\xi) = -\sqrt{-p} \coth(\sqrt{-p} \xi), \quad (3.7)$$

$$\Theta_3(\xi) = -\sqrt{-p} \tanh(2\sqrt{-p} \xi) \pm i\sqrt{-p} \operatorname{sech}(2\sqrt{-p} \xi), \quad (3.8)$$

$$\Theta_4(\xi) = -\sqrt{-p} \coth(2\sqrt{-p} \xi) \pm \sqrt{-p} \operatorname{csch}(2\sqrt{-p} \xi), \quad (3.9)$$

$$\Theta_5(\xi) = -\frac{1}{2} \left( \sqrt{-p} \tanh\left(\frac{\sqrt{-p}}{2} \xi\right) + \sqrt{-p} \coth\left(\frac{\sqrt{-p}}{2} \xi\right) \right). \quad (3.10)$$

Case2: When  $p > 0$ ,

$$\Theta_6(\xi) = \sqrt{p} \tan(\sqrt{p} \xi), \quad (3.11)$$

$$\Theta_7(\xi) = -\sqrt{p} \cot(\sqrt{p} \xi), \quad (3.12)$$

$$\Theta_8(\xi) = -\sqrt{p} \tan(2\sqrt{p} \xi) \pm \sqrt{p} \sec(2\sqrt{p} \xi), \quad (3.13)$$

$$\Theta_9(\xi) = -\sqrt{p} \cot(2\sqrt{p} \xi) \pm \sqrt{p} \csc(2\sqrt{p} \xi), \quad (3.14)$$

$$\Theta_{10}(\xi) = \frac{1}{2} \left( \sqrt{p} \tan\left(\frac{\sqrt{p}}{2} \xi\right) - \sqrt{p} \cot\left(\frac{\sqrt{p}}{2} \xi\right) \right). \quad (3.15)$$

Case3: When  $p = 0$ ,

$$\Theta_{11}(\xi) = -\frac{1}{\xi + d}. \quad (3.16)$$

where  $d$  is a constant. Now, this method for obtaining exact solutions of FDEs consists from the following two main steps:

- Step (1). By substituting (3.4) with Eq.(3.5) into (3.3) and collecting all terms with the same powers of  $\Theta(\xi)$  together, the left hand side of Eq.(3.3) is converted into a polynomial. After setting each coefficient of this polynomial to zero, we obtain a set of algebraic equations in terms of  $A_j$  ( $j = 0, 1, 2, \dots, \eta_1$ ),  $B_j$  ( $j = 0, 1, 2, \dots, \eta_2$ ),  $\kappa$  and  $\omega$ .
- Step (2). Solving the system of algebraic equations and then substituting the results and the general solutions (3.6)-(3.16) into (3.4), it gives travelling wave solutions of (3.3).

## 4 Application

In this section, we consider the DSW system as follows[10, 7]

$$D_t^\alpha u + nvD_x^\beta v = 0, \quad (4.1)$$

$$D_t^\alpha v + qD_x^{3\alpha} v + ruD_x^\beta v + svD_x^\beta u = 0. \quad (4.2)$$

where  $u$  and  $v$  are functions of space variable  $x$  and time variable  $t$  and  $0 < \alpha, \beta \leq 1$ . For obtaining exact solutions of (4.1) and (4.2), We take the traveling wave transformation

$$\begin{aligned} u(x, t) &= U(\eta), \quad v(x, t) = V(\eta), \\ \eta &= \frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha, \end{aligned} \quad (4.3)$$

where  $\omega$  and  $\lambda$  are constants which should to be determined later. Then equations (4.1) and (4.2) are reduced into two nonlinear ODEs

$$-\lambda U' + n\omega VV' = 0, \quad (4.4)$$

$$-\lambda V + q\omega^3 V''' + r\omega UV' + s\omega VU' = 0, \quad (4.5)$$

By integrating of Eq.(4.4) with respect to  $\eta$ , we obtain

$$U = \frac{n\omega}{2\lambda}V^2. \quad (4.6)$$

Substituting equation (4.6) into equation (4.5) yields

$$q\omega^3 V''' + n\omega^2 \left(\frac{r+2s}{2\lambda}\right) V^2 V' - \lambda V' = 0. \quad (4.7)$$

By integrating of equation (4.7) with respect to  $\eta$ , we get

$$q\omega^3 V'' + n\omega^2 \left(\frac{r+2s}{6\lambda}\right) V^3 - \lambda V = 0. \quad (4.8)$$

Balancing  $V''$  with  $V^3$  in (4.8) along with (3.4), we get the below:

$$\eta_1 - \eta_2 + 2 = 3(\eta_1 - \eta_2) \implies \eta_1 = \eta_2 + 1, \quad (4.9)$$

therefore, solution of (4.8) can be expressed as follows.

Type 1:  $\eta_1 = 1$  and  $\eta_2 = 0$ ,

$$V(\eta) = \frac{A_0 + A_1 \Theta(\eta)}{B_0}, \quad (4.10)$$

Substituting (4.10) into (4.8) and making use of equation (3.5) and equating each coefficient of this polynomial to zero, we obtain a set of nonlinear algebraic equations for  $A_0, A_1, B_0, p, \kappa$  and  $\omega$ . Solving obtained system using *Mathematica*, we obtain

$$\bullet \text{Set 1 : } A_0 = 0, \quad A_1 = A_1, \quad B_0 = \frac{i\sqrt{np(r+2s)}\omega}{\sqrt{6}\lambda} A_1, \quad p = \frac{\lambda}{2q\omega^3}. \quad (4.11)$$

By using of the (4.10), (4.11) and cases (3.6)-(3.10) respectively, we get

$$\begin{cases} v_1(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \tanh(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_1(x, t) = \frac{3\lambda}{(r+2s)\omega} \tanh^2(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_2(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \coth(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_2(x, t) = \frac{3\lambda}{(r+2s)\omega} \coth^2(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_3(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\tanh(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{isech}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_3(x, t) = \frac{3\lambda}{(r+2s)\omega} [-\tanh(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{isech}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_4(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\coth(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{csch}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_4(x, t) = \frac{3\lambda}{(r+2s)\omega} [-\coth(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{csch}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_5(x, t) = -\frac{\sqrt{6}\lambda}{2\sqrt{n(r+2s)}\omega} \tanh(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \coth(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_5(x, t) = \frac{3\lambda}{4(r+2s)\omega} \tanh(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \coth(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))^2. \end{cases}$$

By using of the (4.10), (4.11) and cases (3.11)-(3.15) respectively, we get

$$\begin{cases} v_6(x, t) = -\frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_6(x, t) = -\frac{3\lambda}{(r+2s)\omega} \tan^2(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_7(x, t) = \frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_7(x, t) = -\frac{3\lambda}{(r+2s)\omega} \cot^2(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_8(x, t) = \frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_8(x, t) = -\frac{3\lambda}{(r+2s)\omega} [-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_9(x, t) = \frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_9(x, t) = -\frac{3\lambda}{(r+2s)\omega} [-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_{10}(x, t) = -\frac{i\sqrt{6}\lambda}{2\sqrt{n(r+2s)}\omega} [\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_{10}(x, t) = \frac{3\lambda}{4(r+2s)\omega} [\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

Type 2:  $\eta_1 = 2$  and  $\eta_2 = 1$ ,

$$V(\eta) = \frac{A_0 + A_1 \Theta(\eta) + A_2 \Theta^2(\eta)}{B_0 + B_1 \Theta(\eta)}, \quad (4.12)$$

Substituting (4.12) into (4.8) and making use of equation (3.5) and equating each coefficient of this polynomial to zero, we obtain a set of nonlinear algebraic equations for  $A_0, A_1, A_2, B_0, B_1, p, \kappa$  and  $\omega$ . Solving obtained system using *Mathematica*, we obtain

$$\bullet \text{Set 1 : } A_0 = A_0, A_1 = 0, A_2 = \frac{1}{p}A_0, B_0 = 0, B_1 = \frac{\sqrt{n(r+2s)}\omega}{\sqrt{3p}\lambda}A_0, p = -\frac{\lambda}{4q\omega^3}. \quad (4.13)$$

By using of the (4.12), (4.13) and cases (3.11)-(3.15) respectively, we get

$$\begin{cases} v_1(x, t) = \frac{\sqrt{3}\lambda}{\sqrt{n(r+2s)}\omega} [\tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_1(x, t) = \frac{3\lambda}{2(r+2s)\omega} [\tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_2(x, t) = -\frac{\sqrt{3}\lambda}{\sqrt{n(r+2s)}\omega} [\tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_2(x, t) = \frac{3\lambda}{2(r+2s)\omega} [\tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_3(x, t) = \frac{\sqrt{3}\lambda}{\sqrt{n(r+2s)}\omega} \left( \frac{1+[-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right), \\ u_3(x, t) = \frac{3\lambda}{2(r+2s)\omega} \left( \frac{1+[-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right)^2. \end{cases}$$

$$\begin{cases} v_4(x, t) = \frac{\sqrt{3}\lambda}{\sqrt{n(r+2s)}\omega} \left( \frac{1+[-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right), \\ u_4(x, t) = \frac{3\lambda}{2(r+2s)\omega} \left( \frac{1+[-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right)^2. \end{cases}$$

$$\begin{cases} v_5(x, t) = \frac{\sqrt{3}\lambda}{\sqrt{n(r+2s)}\omega} \left( \frac{2+[\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right), \\ u_5(x, t) = \frac{3\lambda}{2(r+2s)\omega} \left( \frac{2+[\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2}{[\tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]} \right)^2. \end{cases}$$

$$\bullet \text{Set 2 : } A_0 = 0, A_1 = A_1, A_2 = A_2, B_0 = \frac{i\sqrt{np(r+2s)}\omega}{\sqrt{6}\lambda}A_1, \\ B_1 = \frac{i\sqrt{np(r+2s)}\omega}{\sqrt{6}\lambda}A_2, p = \frac{\lambda}{2q\omega^3}. \quad (4.14)$$

By using of the (4.12), (4.14) and cases (3.6)-(3.10) respectively, we get

$$\begin{cases} v_1(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \tanh(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_1(x, t) = \frac{3\lambda}{(r+2s)\omega} \tanh^2(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_2(x, t) = -\frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \coth(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_2(x, t) = \frac{3\lambda}{(r+2s)\omega} \coth^2(\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_3(x, t) = \frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\tanh(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{isech}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_3(x, t) = \frac{3\lambda}{(r+2s)\omega} [-\tanh(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{isech}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_4(x, t) = \frac{\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\coth(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{csch}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_4(x, t) = \frac{3\lambda}{(r+2s)\omega} [-\coth(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \operatorname{csch}(2\sqrt{-p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_5(x, t) = -\frac{\sqrt{6}\lambda}{2\sqrt{n(r+2s)}\omega} [\tanh(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \coth(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_5(x, t) = \frac{3\lambda}{4(r+2s)\omega} [\tanh(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) + \coth(\frac{\sqrt{-p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

By using of the (4.12), (4.14) and cases (3.11)-(3.15) respectively, we get

$$\begin{cases} v_6(x, t) = -\frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \tan(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_6(x, t) = -\frac{3\lambda}{(r+2s)\omega} \tan^2(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_7(x, t) = -\frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} \cot(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_7(x, t) = -\frac{3\lambda}{(r+2s)\omega} \cot^2(\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)). \end{cases}$$

$$\begin{cases} v_8(x, t) = -\frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_8(x, t) = -\frac{3\lambda}{(r+2s)\omega} [-\tan(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \sec(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_9(x, t) = -\frac{i\sqrt{6}\lambda}{\sqrt{n(r+2s)}\omega} [-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))], \\ u_9(x, t) = -\frac{3\lambda}{(r+2s)\omega} [-\cot(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) \pm \csc(2\sqrt{p}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))]^2. \end{cases}$$

$$\begin{cases} v_{10}(x, t) = -\frac{i\sqrt{6}\lambda}{2\sqrt{n(r+2s)}\omega} \tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)), \\ u_{10}(x, t) = -\frac{3\lambda}{4(r+2s)\omega} \tan(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha)) - \cot(\frac{\sqrt{p}}{2}(\frac{\omega}{\beta}x^\beta - \frac{\lambda}{\alpha}t^\alpha))^2. \end{cases}$$

## 5 Graphical Presentation of Specific Solutions

In this section, we provide a visual representation of certain solutions corresponding to equations (4.1) and (4.2), utilizing the relationships (4.10) and (4.11). The solutions are depicted in three-dimensional space, and are categorized based on the parameter  $p$ . Specifically, we present solutions for the cases where  $p < 0$  ((3.6)-(3.10)), as well as those where  $p > 0$  ((3.11)-(3.15)). These graphical representations aim to offer an intuitive insight into the behavior of the solutions under specific parameter conditions. The figures labeled from (1) and (2) correspond to various solution scenarios and illustrate the dynamic nature of the system.




Figure 1: In cases  $p = -0.6, n = 3, q = 2, r = 2, s = 1, \lambda = -0.3, \omega = 0.5$






Figure 2: In cases  $p = 0.6, n = 3, q = 2, r = 2, s = 1, \lambda = 0.3, \omega = 0.5$

## 6 Concluding Remarks

In this study, we have employed an extended algebraic method to investigate a specific class of fractional differential equations, namely, the Drinfeld-Sokolov-Wilson system. Our findings demonstrate the effectiveness of the applied method, indicating its potential applicability in solving a wide range of nonlinear evolution equations. It is noteworthy that computational and programming tasks were performed using *Mathematica* throughout this research endeavor.

## References

- [1] M. Cui, *Compact finite difference method for the fractional diffusion equation*, J. Comput. Phys. **228** (2009), no. 20, 7792–7804.
- [2] M. Ekici, M. Mirzazadeh, M. Eslami, Q. Zhou, S.P. Moshokoa, A. Biswas, and M. Belic, *Optical soliton perturbation with fractional-temporal evolution by first integral method with conformable fractional derivatives*, Optik **127** (2016), no. 22, 10659–10669.
- [3] A.M.A. El-Sayed, S.H. Behiry, and W.E. Raslan, *Adomian's decomposition method for solving an intermediate fractional advection–dispersion equation*, Comput. Math. Appl. **59** (2010), no. 5, 1759–1765.
- [4] F. Ferdous, M.G. Hafez, A. Biswas, M. Ekici, Q. Zhou, M. Alifiras, and M. Belic, *Oblique resonant optical solitons with Kerr and parabolic law nonlinearities and fractional temporal evolution by generalized  $\exp(-\Phi(\xi))$ -expansion*, Optik **178** (2019), 439–448.
- [5] J. He, *Variational iteration method for delay differential equations*, Commun. Nonlinear Sci. Numer. Simul. **2** (1997), no. 4, 235–236.
- [6] Y. He, Y. Long, and S. Li, *Exact solutions of the Drinfel'd-Sokolov-Wilson equation using the F-expansion method combined with Exp-function method*, Math. Forum **5** (2010), 3231–3242.
- [7] R. Hirota, B. Grammaticos, and A. Ramani, *Soliton structure of the Drinfel'd-Sokolov-Wilson equation*, J. Math. Phys. **27** (1986), no. 6, 1499–1505.
- [8] M. Inc, *The approximate and exact solutions of the space-and time-fractional Burgers equations with initial conditions by variational iteration method*, J. Math. Anal. Appl. **345** (2008), no. 1, 476–484.
- [9] H. Jafari, N. Kadkhoda, M. Azadi, and M. Yaghoobi, *Group classification of the time-fractional Kaup-Kupershmidt equation*, Sci. Iran. **24** (2017), no. 1, 302–307.
- [10] H.M. Jaradat, S. Al-Shara, Q.J. Khan, M. Alquran, and K. Al-Khaled, *Analytical solution of time-fractional Drinfeld-Sokolov-Wilson system using residual power series method*, IAENG Int. J. Appl. Math. **46** (2016), no. 1, 64–70.
- [11] N. Kadkhoda, *Application of  $\frac{G'}{G^2}$ -expansion method for solving fractional differential equations*, Int. J. Appl. Comput. Math. **3** (2017), no. 1, 1415–1424.
- [12] N. Kadkhoda and H. Jafari, *Application of fractional sub-equation method to the space-time fractional differential equations*, Int. J. Adv. Appl. Math. Mech. **4** (2017), no. 2, 1–6.
- [13] R. Khalil, M. Al Horani, A. Yousef, and M. Sababheh, *A new definition of fractional derivative*, J. Comput. Appl. Math. **264** (2014), 65–70.
- [14] K. Khan, M.A. Akbar, and M.N. Alam, *Traveling wave solutions of the nonlinear Drinfel'd-Sokolov-Wilson equation and modified Benjamin-Bona-Mahony equations*, J. Egypt. Math. Soc. **21** (2013), no. 3, 233–240.
- [15] A.A. Kilbas, H.M. Srivastava, and J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Vol. 204, Elsevier, 2006.
- [16] Z. Odibat and S. Momani, *A generalized differential transform method for linear partial differential equations of fractional order*, Appl. Math. Lett. **21** (2008), no. 2, 194–199.
- [17] M.S. Osman, H. Rezazadeh, M. Eslami, A. Neirameh, and M. Mirzazadeh, *Analytical study of solitons to Benjamin-Bona-Mahony-Peregrine equation with power law nonlinearity by using three methods*, Univ. Polit. Buch. Sci. Bull. Ser. A Appl. Math. Phys. **80** (2018), no. 4, 267–278.

- 
- [18] I. Podlubny, *An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*, Math. Sci. Eng. **198** (1999), 340.
- [19] H. Rezazadeh, *New solitons solutions of the complex Ginzburg-Landau equation with Kerr law nonlinearity*, Optik, **167** (2018), 218–227.
- [20] M. Santillana and C. Dawson, *A numerical approach to study the properties of solutions of the diffusive wave approximation of the shallow water equations*, Comput. Geosci. **14** (2010), no. 1, 31–53.