

Novel concepts of connectivity in vague incidence graphs with application

Nastaran Farhang Rad, Yahya Talebi*

Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Babolsar, Iran

(Communicated by Reza Saadati)

Abstract

Vague sets (VSs) being the most advanced form of fuzzy sets has more capacity to analyze the network state more intelligently. It is proven that VS is more useful to solve many real life problems having uncertainties. Fuzzy graphs (FGs) are efficient mathematical models for analyzing many problems of daily life. One of the most widely used types of FG is vague incidence graph (VIG). VIGs play an important role in various fields such as computer science, psychology, medicine, and political sciences and are used to find effective people in an organization or social institution. They can be used to describe the problems which cannot be handled through FGs and VGs. So, in this paper, specific ideas analogous to vague cut vertices and vague bridges in VGs, vague incidence cut-vertices, and vague incidence bridges are explored. The notion of vague incidence gain and vague incidence less for vague incidence paths and pairs of vertices is also initiated.

Keywords: Vague set, vague graph, vague incidence graph, cut-vertices, bridges
2020 MSC: 05C99, 03E72

1 Introduction

Fuzzy graphs models are beneficial mathematical tools for addressing the combinatorial problems in various fields involving algebra, computing, environmental science, and optimization. In 1965 [36], fuzzy set theory was first introduced by Zadeh. Fuzzy set theory is a highly influential mathematical tool for solving approximate reasoning related problems. Gaa and Buehrer [13] presented and structured the vague set theory. The vague sets describe more possibilities than fuzzy sets. Rosenfeld [30] has defined the concept of a fuzzy graph. Brualdi and Massey [8] introduced the concept of an incidence graph. The fuzzification of the incidence graphs has been proposed by Dinesh [9, 10]. Mordeson et al. [14, 15, 16, 17] defined new concepts in fuzzy incidence graphs. Ramakrishna [23] recommended the vague graph notion and evaluated some of its features. Akram et al. [1, 3, 2] presented new definitions of FGs. Borzooei and Rashmanlou [6, 5, 7, 4] investigated different results on VG. Poulik et al. [11, 21, 22] studied randic index and connectivity concepts in bipolar fuzzy graphs. Samanta et al. [31, 32] defined fuzzy competition graphs and some properties of bipolar fuzzy graphs. Rashmanlou et al. [26, 29, 25, 28, 24] studied new concepts in VGs. Talebi et al. [33, 34, 35] investigated isomorphism on VGs and defined interval-valued intuitionistic fuzzy competition graph. Some properties of interval-valued intuitionistic fuzzy graphs have been given by Borzooei et al. [27].

*Corresponding author

Email addresses: farhang_nastaran@yahoo.com (Nastaran Farhang Rad), talebi@umz.ac.ir (Yahya Talebi)

Connectivity in fuzzy graphs has many application to several fields. Connectivity arises in facility location problems, where the number of facilities (e.g., hospital, fire stations) is fixed and one attempts to minimize the distance that a person needs to travel to the closest facility. Fang et al. [12] presented the connectivity index and wiener index in fuzzy incidence graphs. Nazeer et al. [18, 19] given the notion of order, size, domination, and strong pair domination in fuzzy incidence graphs. Hence, in this paper, norel concepts of vague incidence graphs such as vague incidence gain path, vague incidence loss path, vague incidence balanced path, vague incidence optimal path, vague incidence walk, vague incidence bridges, and vague incidence cut-vertices are introduced.

2 Preliminaries

A FG is the form $y = (\tau, \nu)$ which is a pair of mapping $\tau : V \rightarrow [0, 1]$ and $\nu : V \times V \rightarrow [0, 1]$ which is defined as

$$\nu(p, q) \leq \tau(p) \wedge \tau(q), \quad \forall p, q \in \nu,$$

and ν is a symmetric fuzzy relation on τ and \wedge denotes minimum.

Definition 2.1. [13] A VS z is a pair (t_z, f_z) on set V where t_z and f_z are taken as real valued functions defined on $V \rightarrow [0, 1]$, so that $t_z(p) + f_z(p) \leq 1$, for all $p \in V$.

Definition 2.2. [23] $G = (z, w)$ is named a VG on G^* , which $z = (t_z, f_z)$ is a VS on $E \subseteq V \times V$, so that $t_w(pq) \leq \min(t_z(p), t_z(q))$ and $f_w(pq) \geq \max(f_z(p), f_z(q))$, $\forall pq \in E$.

Definition 2.3. [9] Let $G = (V, E)$ be a graph then, $G^* = (V, E, I)$ is called an incidence graph, so that $I \subseteq V \times E$. If $V = \{p, q\}$, $E = \{pq\}$ and $I = \{(p, pq)\}$, then (V, E, I) is an incidence graph even though $(q, pq) \notin I$. If $(p, pq), (q, pq), (q, qz), (z, qz) \in I$ then pq or qz are called adjacent edges.

Definition 2.4. [10] Let $G^* = (V, E, I)$ be an incidence graph and σ be a fuzzy subset of V and μ , a fuzzy subset of E .

Let ψ be a fuzzy subset of I . If $\psi(v, pq) \leq \sigma(v) \wedge \mu(pq)$, for all $v \in V$ and $pq \in E$, then ψ is called a fuzzy incidence of graph G^* and $G = (\sigma, \mu, \psi)$ is called a fuzzy incidence graph (FIG) of G^* .

3 Vague incidence graph

Definition 3.1. $G = (M, N, P)$ is called a VIG of underlying crisp graph $G^* = (V, E, I)$, if

$$\begin{aligned} M &= \{ \langle t_M(v), f_M(v) \rangle \mid v \in V \}, \\ N &= \{ \langle t_N(pq), f_N(pq) \rangle \mid pq \in E \}, \\ P &= \{ \langle t_P(v, pq), f_P(v, pq) \rangle \mid (v, pq) \in I \}, \end{aligned}$$

such that


$$\begin{aligned} t_N(pq) &\leq t_M(p) \wedge t_M(q), \quad f_N(pq) \geq f_M(p) \vee f_M(q), \\ t_P(v, pq) &\leq t_M(v) \wedge t_N(pq), \\ f_P(v, pq) &\geq f_M(v) \vee f_N(pq), \quad \forall v \in V, pq \in E, \end{aligned}$$

and

$$0 \leq t_M(v) + f_M(v) \leq 1, \quad 0 \leq t_N(pq) + f_N(pq) \leq 1, \quad 0 \leq t_P(v, pq) + f_P(v, pq) \leq 1.$$

Example 3.2. Consider an incidence graph $G^* = (V, E, I)$ so that $V = \{p, q, r, s\}$, $E = \{pq, qr, qs, rs, ps\}$ and $I = \{(p, pq), (q, pq), (q, qr), (r, qr), (q, qs), (s, qs), (r, rs), (s, rs), (s, ps), (p, ps)\}$ as shown in Figure 1. It is easy to show that $G = (M, N, P)$ is a VIG of G^* , as shown in Figure 2, where:

$$\begin{aligned} M &= \left\{ \frac{p}{(0.2, 0.4)}, \frac{q}{(0.3, 0.5)}, \frac{r}{(0.4, 0.6)}, \frac{s}{(0.5, 0.5)} \right\}, \\ N &= \left\{ \frac{pq}{(0.2, 0.6)}, \frac{qr}{(0.3, 0.6)}, \frac{qs}{(0.2, 0.7)}, \frac{rs}{(0.4, 0.6)}, \frac{ps}{(0.2, 0.8)} \right\}, \\ P &= \left\{ \frac{(p, pq)}{(0.2, 0.6)}, \frac{(q, qp)}{(0.2, 0.7)}, \frac{(q, qr)}{(0.2, 0.6)}, \frac{(r, qr)}{(0.3, 0.7)}, \frac{(q, qs)}{(0.2, 0.8)}, \frac{(s, qs)}{(0.2, 0.7)}, \frac{(r, rs)}{(0.3, 0.7)}, \frac{(s, rs)}{(0.4, 0.6)}, \frac{(s, ps)}{(0.2, 0.8)}, \frac{(p, ps)}{(0.2, 0.8)} \right\}. \end{aligned}$$


 Figure 1: Incidence graph G^*

Definition 3.3. The support of a VIG $G = (M, N, P)$ is defined as $G^* = (M^*, N^*, P^*)$, where:

$$M^* = \text{Support of } M = \{v \in V : t_M(v) > 0, f_M(v) > 0\}$$

$$N^* = \text{Support of } N = \{vw \in E : t_N(vw) > 0, f_N(vw) > 0\}$$

$$P^* = \text{Support of } P = \{(v, vw) \in I : t_P(v, vw) > 0, f_P(v, vw) > 0\}.$$

Definition 3.4. A partial vague subgraph is called a partial vague incidence subgraph (PVIS) $H = (M', N', P')$ of a VIG $G = (M, N, P)$ if $t_{P'}(v_i, v_i w_j) \leq t_P(v_i, v_i w_j)$ and $f_{P'}(v_i, v_i w_j) \geq f_P(v_i, v_i w_j)$, for all $(v_i, v_i w_j) \in P^*$.

Definition 3.5. A vague subgraph is called a vague incidence subgraph (VIS) $H = (M', N', P')$ of a VIG $G = (M, N, P)$ if $t_{P'}(v_i, v_i w_j) = t_P(v_i, v_i w_j)$ and $f_{P'}(v_i, v_i w_j) = f_P(v_i, v_i w_j)$, for all $(v_i, v_i w_j)$ is the set of incidence pair of H .

Example 3.6. Figure 3 and Figure 4 are examples of PVIS and VIS of VIG given in Figure 2.

Definition 3.7. A strong VG, G is called strong VIG if $t_P(v_i, v_i w_j) = \wedge\{t_M(v_i), t_N(v_i w_j)\}$ and $f_P(v_i, v_i w_j) = \vee\{f_M(v_i), f_N(v_i w_j)\}$, for each $t_P(v_i, v_i w_j)$ and $f_P(v_i, v_i w_j)$ in P^* .

Definition 3.8. A complete VG, G is called complete VIG if $t_P(v_i, v_i w_j) = \wedge\{t_M(v_i), t_N(v_i w_j)\}$ and $f_P(v_i, v_i w_j) = \vee\{f_M(v_i), f_N(v_i w_j)\}$, for each $v_i, v_j \in M^*$.

Definition 3.9. If $vw \in N^*$ then vw is called an edge of the VIG $G = (M, N, P)$ and if $(v, vw), (w, vw) \in P^*$ then (v, vw) and (w, vw) are incidence paths (IPs) of $G = (M, N, P)$.

Definition 3.10. An vague incidence path

$$P : q_0, (q_0, q_0 q_1), q_0 q_1, (q_1, q_0 q_1), q_1, (q_1, q_1 q_2), q_1 q_2, (q_2, q_1 q_2), q_2, \dots, q_{n-1}, (q_{n-1}, q_{n-1} q_n), q_{n-1} q_n, (q_n, q_{n-1} q_n), q_n$$

is a sequence of different nodes so that either one of the given below condition is satisfied:

- (i) $t_P(v_i, v_i w_j) > 0$ and $f_P(v_i, v_i w_j) > 0$, for some i, j ,
- (ii) $t_P(v_i, v_i w_j) > 0$ and $f_P(v_i, v_i w_j) = 0$, for some i, j ,
- (iii) $t_P(v_i, v_i w_j) = 0$ and $f_P(v_i, v_i w_j) > 0$, for some i, j .

Definition 3.11. A sequence

$$P : q_0, (q_0, q_0 q_1), q_0 q_1, (q_1, q_0 q_1), q_1, (q_1, q_1 q_2), q_1 q_2, (q_2, q_1 q_2), q_2, \dots, q_{n-1}, (q_{n-1}, q_{n-1} q_n), q_{n-1} q_n, (q_n, q_{n-1} q_n), q_n$$

in G is called vague incidence walk (VIW). If $q_0 = q_n$ then a VIW is closed.




Figure 2: VIG G




Figure 3: PVIS of VIG provided in Figure 2

Example 3.12. Let $G = (M, N, P)$ be a VIG provided in Figure 5 where

$$\begin{aligned}
 M &= \{(p, 0.2, 0.4), (q, 0.5, 0.2), (r, 0.2, 0.2), (s, 0.1, 0.4)\}, \\
 N &= \{(pq, 0.2, 0.6), (pr, 0.2, 0.5), (ps, 0.1, 0.7), (qr, 0.2, 0.3), (rs, 0.2, 0.8)\}, \\
 P &= \left\{ \left((p, pq), 0.2, 0.8 \right), \left((q, pq), 0.2, 0.6 \right), \left((p, pr), 0.1, 0.5 \right), \left((r, pr), 0.2, 0.6 \right), \left((p, ps), 0.1, 0.7 \right), \right. \\
 &\quad \left. \left((s, ps), 0.1, 0.8 \right), \left((q, qr), 0.1, 0.4 \right), \left((r, qr), 0.2, 0.4 \right), \left((r, rs), 0.1, 0.8 \right), \left((s, rs), 0.1, 0.8 \right) \right\}.
 \end{aligned}$$

A VIW $P_1 : p, (p, pr), pr(r, pr), r(r, rs), rs, (s, rs), s, (s, sp), sp(p, sp), p$ in Figure 5 is closed because its beginning and final node is similar but it is not a VIP because all nodes are not different.




Figure 4: VIS of VIG given in Figure 2




Figure 5: A VIG G

Definition 3.13. Let $G = (M, N, P)$ be a VIG for any q-r VIP

$$P : q = q_1, (q_1, q_1q_2), q_1q_2, (q_2, q_1q_2), q_2, \dots, q_{n-1}, (q_{n-1}, q_{n-1}q_n), q_{n-1}q_n, (q_n, q_{n-1}q_n), q_n = r$$

in G , we define

$$\wedge \left\{ t_p(q_1, q_1q_2), t_p(q_2, q_1q_2), t_p(q_2, q_2q_3), t_p(q_3, q_2q_3), \dots, t_p(q_{n-1}, q_{n-1}q_n), t_p(q_n, q_{n-1}q_n) \right\}$$

as the vague incidence gain (VIG) of P . VIG of P is expressed by $G_{VI}(p)$ and

$$\vee \left\{ f_p(q_1, q_1q_2), f_p(q_2, q_1q_2), f_p(q_2, q_2q_3), f_p(q_3, q_2q_3), \dots, f_p(q_{n-1}, q_{n-1}q_n), f_p(q_n, q_{n-1}q_n) \right\}$$

as the vague incidence loss (VIL) of P . VIL of P is denoted by $L_{VI}(p)$.

Example 3.14. In Figure 5, for a VIP $P = psr$, $G_{VI}(p) = \wedge(0.1, 0.1, 0.1, 0.1) = 0.1$ and $L_{VI}(p) = \vee(0.7, 0.8, 0.8, 0.8) = 0.8$.

Definition 3.15. A VIP, P is called a vague incidence gain path (VIGP) if $G_{VI}(p) > L_{VI}(p)$ and vague incidence loss path (VILP) if $G_{VI}(p) < L_{VI}(p)$.

Example 3.16. In Figure 5, $P : psr$ is a VILP because $G_{VI}(p) = 0.1 < L_{VI}(p) = 0.8$.

Definition 3.17. A VIG is named as connected if there exists a VIP between each pair of nodes.

Definition 3.18. Let r and s be nodes in a connected VIG G. Among all r-s VIP in G, a VIP whose VIG is greater than or equal to of any other r-s VIP in G is called a maximum r-s VIGP. It is shown by $\max(r - s)_{VIGP}$. Similarly, a r-s VIP whose VIL is smaller than or equal to that of any other r-s VIP in G is known as minimum r-s VILP. It is expressed by $\min(r - s)_{VILP}$. That is a VIP, P is $\max(r - s)_{VIGP}$ if $G_{VI}(p) > G_{VI}(p^\diamond)$ and is a $\min(r - s)_{VILP}$ if $L_{VI}(p) \leq L_{VI}(p^\diamond)$, where p^\diamond is any r-s VIP in G.

Example 3.19. Table 1 shows the vague incidence max gain and vague incidence min loos between each pair of nodes of a VIG given in Figure 5.

Table 1: Name of employees in a hospital and their services.

Vertices	Max-VIG	Max-VIGP	Min-VIL	Min-VILP
p-q	0.2	pq	0.6	prq
p-r	0.1	Any path	0.6	pr
p-s	0.1	Any path	0.8	Any path
q-r	0.1	Any path	0.4	qr
q-s	0.1	Any path	0.8	Any path
r-s	0.1	Any path	0.8	Any path

Definition 3.20. In a VIG, a r-s VIP, P is called a vague incidence balanced path (VIBP) if $G_{VI}(p) = L_{VI}(p)$ and p is called vague incidence optimal path if P is a $\max(r - s)_{VIGP}$ and $\min(r - s)_{VILP}$.

Example 3.21. Figure 5 does not have any VIBP because for each VIP $G_{VI}(p) \neq L_{VI}(p)$ but $r-s : r, (r, rs), rs, (s, rs), s$ is a VIOP.

Definition 3.22. Assume $G = (M, N, P)$ is a VIG and let r and s be any two nodes of G. The VIG of r and s is defined as VIG of $\max(r - s)_{VIGP}$. It is presented by $G_{VI}(r, s)$. In the same way, the VIL of r and s is defined as VIL of $\min(r - s)_{VILP}$. It is denoted by $L_{VI}(r, s)$.

Example 3.23. In Figure 5, $G_{VI}(p, s) = 0.1$, $L_{VI}(p, s) = 0.8$, $G_{VI}(q, r) = 0.1$, $L_{VI}(q, r) = 0.4$.

Definition 3.24. Let H be a VIS of G, r and s be any two nodes of H, then a VIG of r and s in H is the VIG of $\max(r - s)_{VIGP}$ strictly belongs to H and it is shown by $G_{VIH}(r, s)$. In a similar way, VIL of r and s in H is the VIL of $\min(r - s)_{VILP}$ strictly belongs to H and is denoted by $L_{VIH}(r, s)$. If there dose not exist $\max(r - s)_{VIGP}$ and $\min(r - s)_{VILP}$ in H then $G_{VIH}(r, s) = 0$ and $L_{VIH}(r, s) = 0$.

Example 3.25. Let $G = (M, N, P)$ be a VIG provided in Figure 6 which is a VIS of VIG shown in Figure5 which,

$$\begin{aligned}
 M &= \{(p, 0.2, 0.4), (q, 0.5, 0.2), (r, 0.2, 0.2), (s, 0.2, 0.4)\} \\
 N &= \{(pq, 0.2, 0.6), (pr, 0.2, 0.5)\} \\
 P &= \{((p, pq), 0.2, 0.8), ((q, pq), 0.2, 0.6), ((p, pr), 0.1, 0.5), ((r, pr), 0.2, 0.6)\}.
 \end{aligned}$$

It can be seen that $G_{VIH}(r, s) = 0$ and $L_{VIH}(r, s) = 0$




Figure 6: VIS of Figure 5

Proposition 3.26. If H is a VIS of a VIG $G = (M, N, P)$ then, $G_{VIH}(r, s) \leq G_{VI}(r, s)$ and $L_{VIH}(r, s) \leq L_{VI}(r, s)$, for each pairs of nodes r and s .

Proof . Let H be a VIS of G having same number of nodes, edges and VPs with equal membership degree and non-membership degree of nodes, edges and VPs. This implies that $G_{VIH}(r, s) = G_{VI}(r, s)$ and $L_{VIH}(r, s) = L_{VI}(r, s)$, for each pair of nodes r and s in H . Now, if H has less nuber of nodes, edges and VPs then membership degree and non-membership degree of nodes, edges, and VPs will less this implies $G_{VIH}(r, s) < G_{VI}(r, s)$ and $L_{VIH}(r, s) < L_{VI}(r, s)$ so, $G_{VIH}(r, s) \leq G_{VI}(r, s)$ and $L_{VIH}(r, s) \leq L_{VI}(r, s)$. \square

Definition 3.27. Let $G = (M, N, P)$ be a VIG with n nodes, $\{r_1, r_2, \dots, r_n\}$. The vague incidence gain loss matrix (VIGLM) is defined as $M = [G_{VI_{i,j}}, L_{VI_{i,j}}]$ which $G_{VI_{i,j}} = G_{VI}(r_i, r_j)$ and $L_{VI_{i,j}} = L_{VI}(r_i, r_j)$, for $i \neq j$ and $G_{VI_{i,j}} = t_M(r_i)$, $L_{VI_{i,j}} = f_M(r_i)$, for $i = j$.

Example 3.28. VIGLM of G shown in Figure5 is given as below.

$$\begin{bmatrix} (0.2, 0.4) & (0.2, 0.6) & (0.1, 0.6) & (0.1, 0.8) \\ (0.2, 0.6) & (0.5, 0.2) & (0.1, 0.4) & (0.1, 0.8) \\ (0.1, 0.6) & (0.1, 0.4) & (0.2, 0.2) & (0.1, 0.8) \\ (0.1, 0.8) & (0.1, 0.8) & (0.1, 0.8) & (0.2, 0.4) \end{bmatrix}$$

It is clear from the matrix that VIGLM of a VIG is a symmetric matrix.

Definition 3.29. Suppose that $G = (M, N, P)$ is a VIG. A node $q \in M^*$ is called as a vague incidence cut-node (C_{VI}) if there exists two nodes $r, s \in M^*$ of the type $q \neq r \neq s$ so that $G_{VI_{(G-q)}}(r, s) < G_{VI_G}(r, s)$ and $L_{VI_{(G-q)}}(r, s) > L_{VI_G}(r, s)$. A node q is called a vague incidence gain cut-node (C_{VIL}) if $L_{VI_{(G-q)}}(r, s) < L_{VI_G}(r, s)$ and is called a vague incidence loss cut-node (C_{VIG}) if $G_{VI_{(G-q)}}(r, s) > G_{VI_G}(r, s)$.

Example 3.30. In Figure7 , p is a C_{VIL} because $L_{VI_{(G-p)}}(q, r) = 0.9 > L_{VI_G}(q, r) = 0.84$ and r is a C_{VIG} because $G_{VI_{(G-r)}}(q, s) = 0 < G_{VI_G}(q, s) = 0.01$.

Theorem 3.31. A node q in a VIG $G = (M, N, P)$ is a C_{VI} if and only if q is a node in each $\max(r, s)_{VIGP}$ and in each $\min(r, s)_{VILP}$, for some r and s in M^* .




Figure 7: VIG having

Proof . Let $G = (M, N, P)$ be a VIG. Suppose that q is a C_{VI} . According to the definition of C_{VI} , there exists nodes r and s in G so that $q \neq r \neq s$ and (i) : $G_{VI(G-q)}(r, s) < G_{VI_G}(r, s)$ and (ii) : $L_{VI(G-q)}(r, s) > L_{VI_G}(r, s)$. From (i) it is clear that the deleting of q from G deletes all $\max(r, s)_{VIGP}$ and from (ii) it can be seen that the deletion of q deletes each $\min(r, s)_{VILP}$. So, q is in each $\max(r, s)_{VIGP}$ and $\min(r, s)_{VILP}$.

Conversely, suppose that q is in each $\max(r, s)_{VIGP}$ and in each $\min(r, s)_{VILP}$. Then, the deletion of q from G results in the deletion of all $\max(r, s)_{VIGP_s}$ and $\min(r, s)_{VILP_s}$. This shows the G_{VI} will lessen and L_{VI} will enhance between r and s . So, $G_{VI(G-q)}(r, s) < G_{VI_G}(r, s)$ and $L_{VI(G-q)}(r, s) > L_{VI_G}(r, s)$. Therefore, q is a C_{VI} . \square

Definition 3.32. Let $G = (M, N, P)$ be a VIG. An edge $e = rs$ in G is called a vague incidence bridge (B_{VI}) if $G_{VI(G-e)}(p, q) < G_{VI_G}(p, q)$ and $L_{VI(G-e)}(p, q) > L_{VI_G}(p, q)$, for some $p, q \in M^*$.

Example 3.33. Let $G = (M, N, P)$ be a VIG shown in Figure8 which,

$$M = \{(p, 0.2, 0.5), (q, 0.3, 0.3), (r, 0.1, 0.6), (s, 0.4, 0.1)\},$$

$$N = \{(pq, 0.04, 0.73), (pr, 0.07, 0.63), (qr, 0.1, 0.7), (qs, 0.2, 0.4), (rs, 0.1, 0.8)\},$$

$$P = \left\{ \left((p, pq), 0.03, 0.79 \right), \left((q, pq), 0.02, 0.76 \right), \left((p, pr), 0.03, 0.65 \right), \left((r, pr), 0.04, 0.67 \right), \left((q, qr), 0.1, 0.83 \right), \right. \\ \left. \left((r, qr), 0.1, 0.7 \right), \left((q, qs), 0.1, 0.4 \right), \left((s, qs), 0.07, 0.5 \right), \left((r, rs), 0.06, 0.87 \right), \left((s, rs), 0.04, 0.84 \right) \right\}.$$

In Figure 8, edges qr and qs are B_{VI} .

Theorem 3.34. An edge $e \in N^*$ of a VIG $G = (M, N, P)$ is a B_{VI} if and only if it is in each $\max(r, s)_{VIGP}$ and in each $\min(r, s)_{VILP}$ for some r and $s \in M^*$.

Proof . Let $G = (M, N, P)$ be a VIG. Suppose that $e = rs$ is a B_{VI} . So, \exists nodes q and z in G so that (i) : $G_{VI(G-e)}(q, z) < G_{VI_G}(q, z)$, and (ii) : $L_{VI(G-e)}(q, z) > L_{VI_G}(q, z)$. From (i) it can be observe that the deleting of e from G deletes each $\max(q, z)_{VILP}$ and from (ii) it can be seen that the deletion of e deletes each $\min(q, z)_{VILP}$. So, e is in each $\max(q, z)_{VIGP}$ and $\min(q, z)_{VILP}$.

Conversely, suppose that e is in each $\max(q, z)_{VIGP}$ and in each $\min(q, z)_{VILP}$. Then, the deletion of e from G results in the deletion of each $\max(q, z)_{VIGP_s}$ and $\min(q, z)_{VILP_s}$. It shows that G_{VI} will lessen and L_{VI} will enhance between q and z . Therefore, $G_{VI(G-e)}(q, z) < G_{VI_G}(q, z)$ and $L_{VI(G-e)}(q, z) > L_{VI_G}(q, z)$. This proves that $e = rs$ is a B_{VI} . \square




Figure 8: VIG having B_{VI}

4 Conclusion

Vague incidence graph, belonging to the FG family has good capabilities when facing with problems that cannot be expressed by FGs. It has various applications in modern science and technology, especially in the fields of computer science, decision making, and operation research. So, VIG and VISG are defined and their properties are investigated by several examples. Also, specific ideas analogous to vague cut vertices and vague bridges in VGs, vague incidence cut-vertices, and vague incidence bridges are explored. Finally, new concepts of VIGs such as VIGP, VILP, VIBP, VIGLM, and B_{VI} are introduced. One may extend this study to introduce more generalized forms of the connectivity index in VIGs such as Randic index, Zagreb index, Hyper Zagreb index, Wiener index etc. This research can be extended towards spherical and T-spherical fuzzy sets.

References

- [1] M. Akram, A. Farooq, A.B. Saeid, and K.P. Shum, *Certain types of vague cycles and vague trees*, J. Intell. Fuzzy Syst. **28** (2015), no. 2, 621–631.
- [2] M. Akram, F. Feng, S. Sarwar, and Y.B. Jun, *Certain types of vague graphs*, U.P.B. Sci. Bull. Series A. **76** (2014), no. 1, 141–154.
- [3] M. Akram, S. Samanta, and M. Pal, *Cayley Vague Graphs*, J. Fuzzy Math. **25** (2017), no. 2, 1–14.
- [4] R.A. Borzooei and H. Rashmanlou, *Cayley interval-valued fuzzy graphs*, U.P.B. Sci. Bull. Ser. A: Appl. Math. Phys. **78** (2016), no. 3, 83–94.
- [5] R.A. Borzooei and H. Rashmanlou, *Degree of vertices in vague graphs*, J. Appl. Math. Inf. **33** (2015), no. 5, 545–557.
- [6] R.A. Borzooei and H. Rashmanlou, *Domination in vague graphs and its applications*, J. Intell. Fuzzy Syst. **29** (2015), 1933–1940.
- [7] R.A. Borzooei, H. Rashmanlou, S. Samanta, and M. Pal, *Regularity of vague graphs*, J. Intell. Fuzzy Syst. **30** (2016), 3681–3689.
- [8] R.A. Brualdi and J.L.Q. Massey, *Incidence and strong edge colorings of graphs*, Discret. Math. **122** (1993), 51–58.
- [9] T. Dinesh, *A study on graph structures, incidence algebras and their fuzzy analogous*, Ph.D. Thesis, Kannur University, Kerala, India, 2012.

- [10] T. Dinesh, *Fuzzy incidence graph-an introduction* Adv. Fuzzy Sets Syst. **21** (2016), 33–48.
- [11] S. Das, S. Poulik, G. Ghorai, *Picture fuzzy phi-tolerance competition graphs with its application*, J. Ambient Intell. Human. Comput. **15** (2023), 547–559.
- [12] I. Fang, I. Nazeer, T. Rashid, and J.B. Lio, *Connectivity and Wiener index of fuzzy incidence graphs*, Math. Prob. Eng. **2021** (2021) 1–7.
- [13] W.-L. Gau and D.J. Buehrer, *Vague sets*, IEEE Transactions on Systems, Man, Cybernet. **23** (1993), no. 2, 610–614.
- [14] J.N. Mordeson and S. Mathew, *Fuzzy end nodes in fuzzy incidence graphs*, New Math. Nat. Comput. **13** (2017), 13–20.
- [15] J.N. Mordeson and S. Mathew, *Human trafficking: Source, transit, destination, designations*, New Math. Nat. Comput. **13** (2017), 209–218.
- [16] J.N. Mordeson and S. Mathew, *Local look at human trafficking*, New Math. Nat. Comput. **13** (2017), 327–340.
- [17] J.N. Mordeson, S. Mathew, and R.A. Borzooei, *Vulnerability and government response to human trafficking: Vague fuzzy incidence graphs*, New Math. Nat. Comput. **14** (2018), 203–219.
- [18] I. Nazeer, T. Rashid, and J.L.G. Guirao, *Domination of fuzzy incidence graphs with the algorithm and application for the selection of a medical lab*, Math. Prob. Eng. **2021** (2021), Article ID 6682502. 1–11.
- [19] I. Nazeer, T. Rashid, M.T. Hussain, and J.L.G. Guirao, *Domination in join of fuzzy incidence graphs using strong pairs with application in trading system of different countries*, Symmetry **13** (2021), no. 7, 1–15.
- [20] T. Nguyen Xuan and T. Phan Nhat, *On the existence of equilibrium points of vector functions*, Numer. Funct. Anal. Optim. **19** (1998), 141–156.
- [21] S. Poulik, S. Das, and G. Ghorai, *Randic index of bipolar fuzzy graphs and its application in network systems*, J. Appl. Math. Comput. **68** (2021), 2317–2341.
- [22] S. Poulik and G. Ghorai, *Connectivity concepts in bipolar fuzzy incidence graphs*, Thai J. Math. **20** (2022), no. 4, 1609–1619.
- [23] N. Ramakrishna, *Vague graphs*, Int. J. Cogn. Comput. **7** (2009), 51–58.
- [24] H. Rashmanlou and R.A. Borzooei, *New concepts of interval-valued intuitionistic (S, T) -fuzzy graphs*, J. Intell. Fuzzy Syst. **30** (2016), no. 4, 1893–1901.
- [25] H. Rashmanlou and R.A. Borzooei, *Product vague graphs and its applications*, J. Intell. Fuzzy Syst. **30** (2016), 371–382.
- [26] H. Rashmanlou and R.A. Borzooei, *Vague graphs with application*, J. Intell. Fuzzy Syst. **30** (2016), 3291–3299.
- [27] H. Rashmanlou, R.A. Borzooei, S. Samanta, and M. Pal, *Properties of interval valued intuitionistic (s, t) -fuzzy graphs*, Pac. Sci. Rev. A: Nat. Sci. Engin. **18** (2016), no. 1, 30–37.
- [28] H. Rashmanlou, Y.B. Jun, and R.A. Borzooei, *More results on highly irregular bipolar fuzzy graphs*, Ann. Fuzzy Math. Inf. **8** (2014), 149–168.
- [29] H. Rashmanlou, S. Samanta, M. Pal, R.A. Borzooei, *A study on bipolar fuzzy graphs*, J. Intell. Fuzzy Syst. **28** (2015), 571–580.
- [30] A. Rosenfeld, *Fuzzy graphs, fuzzy sets and their applications*, Zadeh, L.A., Fu, K.S., Shimura, M., Eds.; Academic Press: New York, NY, USA, 1975, pp. 77–95.
- [31] S. Samanta, M. Akram, and M. Pal, *m-step fuzzy competition graphs*, J. Appl. Math. Comput. **47** (2015), no. 1, 461–472.
- [32] S. Samanta and M. Pal, *Some more results on bipolar fuzzy sets and bipolar fuzzy intersection graphs*, J. Fuzzy Math. **22** (2014) 253–262.
- [33] A.A. Talebi, H. Rashmanlou, and N. Mehdipoor, *Isomorphism on vague graphs*, Ann. Fuzzy Math. Inf. **6** (2013), no. 3, 575–588.

-
- [34] A.A. Talebi, H. Rashmanlou, and S.H. Sadati, *Interval-valued Intuitionistic Fuzzy Competition Graph*, J. Multiple-Valued Logic Soft Comput. **34** (2020).
- [35] A.A. Talebi, H. Rashmanlou, and S.H. Sadati, *New concepts on m-polar interval-valued intuitionistic fuzzy graph*, TWMS J. Appl. Engin. Math. **10** (2020), no. 3, 806–818.
- [36] L.A. Zadeh, *Fuzzy sets*, Inf. Control **8** (1965), 338–353.