

On a class of multivalent meromorphic functions defined by combinational differential operator

Mohammadreza Foroutan^{a,*}, Mostafa Soltani^a, Ali Ebadian^b

^aDepartment of Mathematics, Payame Noor University, P.O.Box 19395-3697, Tehran, Iran

^bDepartment of Mathematics, Faculty of Science, Urmia University, Urmia, Iran

(Communicated by Mugur Alexandru Acu)

Abstract

In this paper, we define a class of meromorphically multivalent functions in $\mathbb{U}^* = \{z : z \in \mathbb{C} : 0 < |z| < 1\}$ by using a differential operator. Important properties of this class such as coefficient estimates, distortion theorem, radius of starlikeness and convexity, closure theorems, and convolution properties are obtained. We also study δ -neighborhoods and partial sums for this class.

Keywords: meromorphic functions, multivalent functions, differential operator, coefficient estimates, convolution, neighborhoods

2020 MSC: 30C45; 30C50

1 Introduction

Let Σ_p denote the class of functions of the form:

$$f(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k} z^{p+k} \quad (a_{p+k} \geq 0; p \in \mathbb{N} = \{1, 2, \dots\}), \quad (1.1)$$

which are analytic and p -valent in the punctured unit disk $\mathbb{U}^* = \{z : z \in \mathbb{C} : 0 < |z| < 1\} = \mathbb{U} \setminus \{0\}$. A function $f \in \Sigma_p$ is meromorphically starlike of order ρ ($0 \leq \rho < p$) if

$$-\Re \left\{ \frac{zf'(z)}{f(z)} \right\} > \rho.$$

A function $f \in \Sigma_p$ is meromorphically convex of order θ ($0 \leq \theta < p$) if

$$-\Re \left\{ 1 + \frac{zf''(z)}{f'(z)} \right\} > \theta.$$

*Corresponding author

Email addresses: foroutan_mohammadreza@yahoo.com, mr_forootan@pnu.ac.ir, (Mohammadreza Foroutan), soltanimath62@gmail.com (Mostafa Soltani), a.ebadian@urmia.ac.ir (Ali Ebadian)

If $f \in \Sigma_p$ is given by (1.1) and $g \in \Sigma_p$ is given by

$$g(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} b_{p+k} z^{p+k} \quad (b_{p+k} \geq 0; p \in \mathbb{N}). \quad (1.2)$$

then the Hadamard product (or convolution) of f and g is defined by

$$(f * g)(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k} b_{p+k} z^{p+k} = (g * f)(z) \quad (z \in \mathbb{U}^*; p \in \mathbb{N}). \quad (1.3)$$

For functions $f(z) \in \Sigma_p$, Aouf [1] defined the following differential operator:

$$\begin{aligned} S_{\lambda,p}^0 f(z) &= f(z) \\ S_{\lambda,p}^1 f(z) &= (1 - \lambda)f(z) + \frac{\lambda}{p} z f'(z) + \frac{2\lambda}{z^p} \\ &= \frac{1}{z^p} + \sum_{k=0}^{\infty} \left(\frac{p + \lambda k}{p} \right) a_{p+k} z^{p+k} \\ &= S_{\lambda,p} f(z) \quad (\lambda \geq 0; p \in \mathbb{N}) \\ S_{\lambda,p}^2 f(z) &= S_{\lambda,p} (S_{\lambda,p}^1 f(z)) \end{aligned}$$

and

$$\begin{aligned} S_{\lambda,p}^n f(z) &= S_{\lambda,p} (S_{\lambda,p}^{n-1} f(z)) \\ &= (1 - \lambda) S_{\lambda,p}^{n-1} f(z) + \frac{\lambda}{p} z (S_{\lambda,p}^{n-1} f(z))' + \frac{2\lambda}{z^p} \quad (\lambda \geq 0; n, p \in \mathbb{N}). \end{aligned}$$

It can be easily seen that

$$S_{\lambda,p}^n f(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} \left(\frac{p + \lambda k}{p} \right)^n a_{p+k} z^{p+k}, \quad (n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, p \in \mathbb{N}). \quad (1.4)$$

Also Orhan et al. [4] defined the differential operator $T_{\sigma\mu p}^n$ in the following way:

$$\begin{aligned} T_{\sigma\mu p}^0 f(z) &= f(z) \\ T_{\sigma\mu p}^1 f(z) &= T_{\sigma\mu p} f(z) = \sigma\mu \frac{[z^{p+1} f(z)]''}{z^{p-1}} + (\sigma - \mu) \frac{[z^{p+1} f(z)]'}{z^p} + (1 - \sigma + \mu) f(z) \end{aligned} \quad (1.5)$$

and, in general,

$$T_{\sigma\mu p}^n f(z) = T_{\sigma\mu p} (T_{\sigma\mu p}^{n-1} f(z)), \quad (1.6)$$

where $0 \leq \mu \leq \sigma$ and $n \in \mathbb{N}_0$. Recently, Sharma [6] used the generalized modified Srivastava-Gupta operators defined in [7] by using iterative combinations in ordinary and simultaneous approximation and obtained several interesting results. If the function $f(z) \in \Sigma_p$ is given by (1.1) then from (1.5) and (1.6), we obtain

$$T_{\sigma\mu p}^n f(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} \Psi_k(\sigma, \mu, n, p) a_{k+p} z^{k+p} \quad (1.7)$$

where

$$\Psi_k(\sigma, \mu, n, p) = [1 + (k + 2p)(\sigma - \mu + (k + 2p + 1)\sigma\mu)]^n. \quad (1.8)$$

Making use of the differential operators $S_{\lambda,p}^n f(z)$ and $T_{\sigma\mu p}^n f(z)$ defined as in (1.4) and (1.7), respectively, we define the following differential operator for the functions $f(z) \in \Sigma_p$ as follows:

$$D_{\lambda,\sigma,\mu,\omega,p}^n f(z) = (1 - \omega) S_{\lambda,p}^n f(z) + \omega T_{\sigma\mu p}^n f(z), \quad (1.9)$$

for $n \in \mathbb{N}, \lambda \geq 0, 0 \leq \mu \leq \sigma, 0 \leq \omega \leq 1$. Let $f(z)$ be given by (1.1), then using (1.4) and (1.7), the relation (1.9) can be easily written as

$$D_{\lambda, \sigma, \mu, \omega, p}^n f(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} z^{p+k} \tag{1.10}$$

where

$$\Phi_k(n, \lambda, \sigma, \mu, \omega, p) = (1 - \omega) \left(\frac{p + \lambda k}{p} \right) + \omega \Psi_k(\sigma, \mu, n, p) \tag{1.11}$$

and

$$\Psi_k(\sigma, \mu, n, p) = [1 + (k + 2p)(\sigma - \mu + (k + 2p + 1)\sigma\mu)]^n, \tag{1.12}$$

for $n \in \mathbb{N}, \lambda \geq 0, 0 \leq \mu \leq \sigma, 0 \leq \omega \leq 1$. Therefore in terms of convolution (1.9) is equivalent to

$$D_{\lambda, \sigma, \mu, \omega, p}^n f(z) = (f * h)(z),$$

where

$$h(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} \Phi_k(n, \lambda, \sigma, \mu, \omega, p) z^{p+k}.$$

Using the differential operator $D_{\lambda, \sigma, \mu, \omega, p}^n f(z)$, we define the following subclass of multivalent meromorphic functions

Definition 1.1. A function $f(z) \in \Sigma_p$ is said to be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$ if it satisfies the following inequality:

$$\left| \frac{z^{p+2} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))'' + z^{p+1} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' - p^2}{\gamma z^{p+1} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' + \alpha(1 + \gamma)p - p} \right| < \beta \tag{1.13}$$

where $0 \leq \alpha < 1, 0 < \beta \leq 1, 0 \leq \gamma \leq 1$.

Meromorphically multivalent functions have been extensively studied, for example, see Najafzadeh and Ebadian [3], Atshan and Kulkarni [2], Orhan et al. [4] and Auof [1].

2 Coefficients Estimates

Theorem 2.1. A function f defined by (1.1) is in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$ if and only if

$$\sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)a_{p+k} \leq \beta p(1-\alpha)(1+\gamma) \tag{2.1}$$

where $\Phi_k(n, \lambda, \sigma, \mu, \omega, p)$ is given by (1.11)

Proof . Assume (2.1) holds. it is enough to show that

$$M = \left| z^{p+2} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))'' + z^{p+1} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' - p^2 \right| - \beta \left| \gamma z^{p+1} (D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' + \alpha(1 + \gamma)p - p \right| < 0.$$

For $|z| = r < 1$, by (2.1), we have

$$\begin{aligned} M &= \left| \sum_{k=0}^{\infty} (p+k)^2 \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} r^{2p+k} \right| - \beta \left| p(1-\alpha)(1+\gamma) - \gamma \sum_{k=0}^{\infty} (p+k) \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} r^{2p+k} \right| \\ &\leq \sum_{k=0}^{\infty} (p+k)^2 \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} r^{2p+k} - \beta p(1-\alpha)(1+\gamma) + \beta \gamma \sum_{k=0}^{\infty} (p+k) \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} r^{2p+k} \\ &< \sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)a_{p+k} - \beta p(1-\alpha)(1+\gamma) \\ &< 0. \end{aligned}$$

Hence $f \in \mathcal{H}_p(\alpha, \beta, \gamma)$. Conversely, suppose that $f(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, then (1.13) holds true. So, we have

$$\left| \frac{z^{p+2}(D_{\lambda, \sigma, \mu, \omega, p}^n f(z))'' + z^{p+1}(D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' - p^2}{\gamma z^{p+1}(D_{\lambda, \sigma, \mu, \omega, p}^n f(z))' + \alpha(1 + \gamma)p - p} \right| = \left| \frac{\sum_{k=0}^{\infty} (p+k)^2 \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} z^{2p+k}}{p(1-\alpha)(1+\gamma) - \gamma \sum_{k=0}^{\infty} (p+k) \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} z^{2p+k}} \right| < \beta.$$

Since $\Re \epsilon z \leq |z|$, for all z , we have

$$\Re \left\{ \frac{\sum_{k=0}^{\infty} (p+k)^2 \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} z^{2p+k}}{p(1-\alpha)(1+\gamma) - \gamma \sum_{k=0}^{\infty} (p+k) \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} z^{2p+k}} \right\} < \beta.$$

Now by letting $z \rightarrow 1^-$ through real axis, we obtain

$$\sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma] \Phi_k(n, \lambda, \sigma, \mu, \omega, p) a_{p+k} \leq \beta p(1-\alpha)(1+\gamma).$$

Hence the result follows. \square

Corollary 2.2. If $f(z)$ defined by (1.1) is in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$, then

$$a_{p+k} \leq \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)[p+k+\beta\gamma] \Phi_k(n, \lambda, \sigma, \mu, \omega, p)}.$$

The result is sharp for the function $f(z)$ given by

$$f(z) = \frac{1}{z^p} + \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)[p+k+\beta\gamma] \Phi_k(n, \lambda, \sigma, \mu, \omega, p)} z^{p+k} \quad (2.2)$$

where $\Phi_k(n, \lambda, \sigma, \mu, \omega, p)$ is given by (1.11).

3 Distortion Theorem

Theorem 3.1. If $f(z)$ defined by (1.1) is in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$, then for $0 < |z| = r < 1$

$$\frac{1}{r^p} - \frac{\beta p(1-\alpha)(1+\gamma)}{p(p+\beta\gamma) \Phi_0(n, \lambda, \sigma, \mu, \omega, p)} r^p \leq |f(z)| \leq \frac{1}{r^p} + \frac{\beta p(1-\alpha)(1+\gamma)}{p(p+\beta\gamma) \Phi_0(n, \lambda, \sigma, \mu, \omega, p)} r^p \quad (3.1)$$

and

$$\frac{p}{r^{p+1}} - \frac{\beta p(1-\alpha)(1+\gamma)}{(p+\beta\gamma) \Phi_0(n, \lambda, \sigma, \mu, \omega, p)} r^{p-1} \leq |f'(z)| \leq \frac{p}{r^{p+1}} + \frac{\beta p(1-\alpha)(1+\gamma)}{(p+\beta\gamma) \Phi_0(n, \lambda, \sigma, \mu, \omega, p)} r^{p-1} \quad (3.2)$$

where

$$\Phi_0(n, \lambda, \sigma, \mu, \omega, p) = \left[(1-\omega) + \omega \left[1 + 2p(\sigma - \mu + (2p+1)\sigma\mu) \right]^n \right]. \quad (3.3)$$

The bounds are attained for the function $f(z)$ given by

$$f(z) = \frac{1}{z^p} + \frac{\beta p(1-\alpha)(1+\gamma)}{p(p+\beta\gamma) \Phi_0(n, \lambda, \sigma, \mu, \omega, p)} z^p. \quad (3.4)$$

Proof . In view of Theorem 2.1 we have

$$p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p) \sum_{k=0}^{\infty} a_{p+k} \leq \sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)a_{p+k} \leq \beta p(1-\alpha)(1+\gamma)$$

which is equivalent to

$$\sum_{k=0}^{\infty} a_{p+k} \leq \frac{\beta p(1-\alpha)(1+\gamma)}{p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}. \tag{3.5}$$

Thus for $0 < |z| = r < 1$, we get

$$\begin{aligned} |f(z)| &\leq \frac{1}{r^p} + \sum_{k=0}^{\infty} a_{p+k}r^{p+k} \\ &\leq \frac{1}{r^p} + r^p \sum_{k=0}^{\infty} a_{p+k} \\ &\leq \frac{1}{r^p} + \frac{\beta p(1-\alpha)(1+\gamma)}{p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}r^p \end{aligned} \tag{3.6}$$

and

$$\begin{aligned} |f(z)| &\geq \frac{1}{r^p} - \sum_{k=0}^{\infty} a_{p+k}r^{p+k} \\ &\geq \frac{1}{r^p} - r^p \sum_{k=0}^{\infty} a_{p+k} \\ &\geq \frac{1}{r^p} - \frac{\beta p(1-\alpha)(1+\gamma)}{p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}r^p \end{aligned} \tag{3.7}$$

which, together, yield (3.1). Furthermore, it follows from Theorem 2.1 that

$$\sum_{k=0}^{\infty} (p+k)a_{p+k} \leq \frac{\beta p(1-\alpha)(1+\gamma)}{(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}. \tag{3.8}$$

Hence

$$\begin{aligned} |f'(z)| &\leq \frac{p}{r^{p+1}} + \sum_{k=0}^{\infty} (p+k)a_{p+k}r^{p+k-1} \\ &\leq \frac{p}{r^{p+1}} + r^{p-1} \sum_{k=0}^{\infty} (p+k)a_{p+k} \\ &\leq \frac{p}{r^{p+1}} + \frac{\beta p(1-\alpha)(1+\gamma)}{(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}r^{p-1} \end{aligned} \tag{3.9}$$

and

$$\begin{aligned} |f'(z)| &\geq \frac{p}{r^{p+1}} - \sum_{k=0}^{\infty} (p+k)a_{p+k}r^{p+k-1} \\ &\geq \frac{p}{r^{p+1}} - r^{p-1} \sum_{k=0}^{\infty} (p+k)a_{p+k} \\ &\geq \frac{p}{r^{p+1}} - \frac{\beta p(1-\alpha)(1+\gamma)}{(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}r^{p-1}, \end{aligned} \tag{3.10}$$

which together yield (3.2). It can be easily seen that the function $f(z)$ defined by (3.4) is extremal for Theorem 3.1. \square

4 Radius of Starlikeness and Convexity

Theorem 4.1. Let the function $f(z)$ defined by (1.1) be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$ then $f(z)$ is meromorphically p -valent starlike of order ρ ($0 \leq \rho < p$) in $0 < |z| < r(n, \lambda, \sigma, \mu, \omega, p, \rho, \alpha, \beta, \gamma)$, where

$$r(n, \lambda, \sigma, \mu, \omega, p, \rho, \alpha, \beta, \gamma) = \inf_k \left[\frac{(p - \rho)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p + k)(p + k + \beta\gamma)}{\beta p(1 - \alpha)(1 + \gamma)(3p + k - \rho)} \right]^{\frac{1}{2p + k}}, \quad (4.1)$$

$k \geq 0, p \in \mathbb{N}, n \in \mathbb{N}_0$, the result is sharp.

Proof . It is sufficient to show that

$$\left| \frac{zf'(z) + pf(z)}{f(z)} \right| \leq p - \rho,$$

for $0 < |z| < r(n, \lambda, \sigma, \mu, \omega, p, \rho, \alpha, \beta, \gamma)$. Note that

$$\begin{aligned} \left| \frac{zf'(z) + pf(z)}{f(z)} \right| &= \left| \frac{\sum_{k=0}^{\infty} (2p + k)a_{p+k}z^{p+k}}{z^{-p} + \sum_{k=0}^{\infty} a_{p+k}z^{p+k}} \right| \\ &\leq \frac{\sum_{k=0}^{\infty} (2p + k)a_{p+k}r^{2p+k}}{1 - \sum_{k=0}^{\infty} a_{p+k}r^{2p+k}}. \end{aligned}$$

Thus, $\left| \frac{zf'(z) + pf(z)}{f(z)} \right| \leq p - \rho$, if

$$\sum_{k=0}^{\infty} \frac{(3p + k - \rho)}{(p - \rho)} a_{p+k} r^{2p+k} \leq 1. \quad (4.2)$$

Theorem 2.1 ensures that

$$\sum_{k=0}^{\infty} \frac{(p + k)[p + k + \beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{\beta p(1 - \alpha)(1 + \gamma)} a_{p+k} \leq 1, \quad (4.3)$$

in view of (4.3), it follows that (4.2) will be true if

$$\frac{(3p + k - \rho)}{(p - \rho)} r^{2p+k} \leq \frac{(p + k)[p + k + \beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{\beta p(1 - \alpha)(1 + \gamma)} \quad (4.4)$$

or if

$$r \leq \left[\frac{(p - \rho)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p + k)(p + k + \beta\gamma)}{\beta p(1 - \alpha)(1 + \gamma)(3p + k - \rho)} \right]^{\frac{1}{2p + k}}. \quad (4.5)$$

Setting $r(n, \lambda, \sigma, \mu, \omega, p, \rho, \alpha, \beta, \gamma)$ in (4.5), the result follows. The result is sharp with the extremal function $f(z)$ given by (2.2). \square

Theorem 4.2. Let the function $f(z)$ defined by (1.1) be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$. Then $f(z)$ is meromorphically p -valent convex of order θ ($0 \leq \theta < p$) in $0 < |z| < r(n, \lambda, \sigma, \mu, \omega, p, \theta, \alpha, \beta, \gamma)$, where

$$r(n, \lambda, \sigma, \mu, \omega, p, \theta, \alpha, \beta, \gamma) = \inf_k \left[\frac{p(p - \theta)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p + k + \beta\gamma)}{\beta p(1 - \alpha)(1 + \gamma)(3p + k - \theta)} \right]^{\frac{1}{2p + k}}, \quad (4.6)$$

$k \geq 0, p \in \mathbb{N}, n \in \mathbb{N}_0$, the result is sharp.

Proof . It suffices to show that

$$\left| \frac{(zf'(z))' + pf'(z)}{f'(z)} \right| \leq p - \theta,$$

for $0 < |z| < r(n, \lambda, \sigma, \mu, \omega, p, \theta, \alpha, \beta, \gamma)$. Note that

$$\begin{aligned} \left| \frac{(zf'(z))' + pf'(z)}{f'(z)} \right| &= \left| \frac{\sum_{k=0}^{\infty} (p+k)(2p+k)a_{p+k}z^{p+k-1}}{-pz^{-(p+1)} + \sum_{k=0}^{\infty} (p+k)a_{p+k}z^{p+k-1}} \right| \\ &\leq \frac{\sum_{k=0}^{\infty} (p+k)(2p+k)a_{p+k}r^{2p+k}}{p - \sum_{k=0}^{\infty} (p+k)a_{p+k}r^{2p+k}}. \end{aligned}$$

Thus, $\left| \frac{(zf'(z))' + pf'(z)}{f'(z)} \right| \leq p - \theta$, if

$$\sum_{k=0}^{\infty} \frac{(p+k)(3p+k-\theta)}{p(p-\theta)} a_{p+k} r^{2p+k} \leq 1. \tag{4.7}$$

Theorem 2.1 ensures that

$$\sum_{k=0}^{\infty} \frac{(p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{\beta p(1-\alpha)(1+\gamma)} a_{p+k} \leq 1, \tag{4.8}$$

so in view of (4.8) it follows that (4.7) will be true if

$$\frac{(p+k)(3p+k-\theta)}{p(p-\theta)} r^{2p+k} \leq \frac{(p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{\beta p(1-\alpha)(1+\gamma)}, \tag{4.9}$$

or, if

$$r \leq \left[\frac{p(p-\theta)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p+k+\beta\gamma)}{\beta p(1-\alpha)(1+\gamma)(3p+k-\theta)} \right]^{\frac{1}{2p+k}}. \tag{4.10}$$

Therefore, we obtain the $r(n, \lambda, \sigma, \mu, \omega, p, \theta, \alpha, \beta, \gamma)$ in (4.6). The result is sharp for the extremal function $f(z)$ given by (2.2). \square

5 Closure Theorems

Theorem 5.1. Let

$$f_{p-1} = \frac{1}{z^p} \tag{5.1}$$

and

$$f_{p+k} = \frac{1}{z^p} + \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p+k+\beta\gamma)} z^{p+k}, \tag{5.2}$$

where, $k \geq 0, p \in \mathbb{N}, n \in \mathbb{N}_0$, and $z \in U^*$. Then $f(z)$ is in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$ if and only if it can be expressed in the form

$$f(z) = \sum_{k=-1}^{\infty} c_{p+k} f_{p+k}(z) \tag{5.3}$$

where $c_{p+k} \geq 0$ and $\sum_{k=-1}^{\infty} c_{p+k} = 1$.

Proof . Let $f(z) = \sum_{k=-1}^{\infty} c_{p+k} f_{p+k}(z)$, where $c_{p+k} \geq 0$ and $\sum_{k=-1}^{\infty} c_{p+k} = 1$. Then

$$\begin{aligned} f(z) &= \sum_{k=-1}^{\infty} c_{p+k} f_{p+k}(z) \\ &= \frac{1}{z^p} + \sum_{k=0}^{\infty} c_{p+k} \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p+k+\beta\gamma)} z^{p+k}. \end{aligned} \quad (5.4)$$

We have

$$\sum_{k=0}^{\infty} c_{p+k} \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p+k+\beta\gamma)} \times \frac{(p+k)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)(p+k+\beta\gamma)}{\beta p(1-\alpha)(1+\gamma)} = \sum_{k=0}^{\infty} c_{p+k} = 1 - c_{p-1} \leq 1. \quad (5.5)$$

Then Theorem 2.1 shows $f(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$. Conversely, suppose $f(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$. Then by Corollary 2.2, we have

$$a_{p+k} \leq \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}.$$

Set

$$c_{p+k} = \frac{(p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{\beta p(1-\alpha)(1+\gamma)} a_{p+k} \quad \text{and} \quad c_{p-1} = 1 - \sum_{k=0}^{\infty} c_{p+k}.$$

Then

$$\begin{aligned} f(z) &= \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k} z^{p+k} \\ &= \frac{1}{z^p} + \sum_{k=0}^{\infty} \frac{\beta p(1-\alpha)(1+\gamma)}{(p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)} c_{p+k} z^{p+k} \\ &= \frac{1}{z^p} + \sum_{k=0}^{\infty} (f_{p+k}(z) - \frac{1}{z^p}) c_{p+k} \\ &= \frac{1}{z^p} \left(1 - \sum_{k=0}^{\infty} c_{p+k} \right) + \sum_{k=0}^{\infty} c_{p+k} f_{p+k}(z) \\ &= \frac{1}{z^p} c_{p-1} + \sum_{k=0}^{\infty} c_{p+k} f_{p+k}(z) \\ &= \sum_{k=-1}^{\infty} c_{p+k} f_{p+k}(z). \end{aligned}$$

This completes the proof of Theorem 5.1. \square

Theorem 5.2. The class $\mathcal{H}_p(\alpha, \beta, \gamma)$ is closed under convex linear combinations.

Proof . Let each of the functions

$$f_j(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k,j} z^{p+k} \quad (a_{p+k,j} \geq 0; j = 1, 2), \quad (5.6)$$

be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$. It is sufficient to show that the function $h(z)$ defined by

$$h(z) = (1-t)f_1(z) + tf_2(z) \quad (0 \leq t \leq 1), \quad (5.7)$$

is also in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$. Since

$$h(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} [(1-t)a_{p+k,1} + ta_{p+k,2}]z^{p+k} \quad (0 \leq t \leq 1), \tag{5.8}$$

by Theorem 2.1, we have

$$\begin{aligned} & \sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)[(1-t)a_{p+k,1} + ta_{p+k,2}] \\ &= (1-t) \sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)a_{p+k,1} + t \sum_{k=0}^{\infty} (p+k)[p+k+\beta\gamma]\Phi_k(n, \lambda, \sigma, \mu, \omega, p)a_{p+k,2} \\ &\leq (1-t)\beta p(1-\alpha)(1+\gamma) + t\beta p(1-\alpha)(1+\gamma) = \beta p(1-\alpha)(1+\gamma), \end{aligned}$$

which shows that $h(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, hence the result follows. \square

6 Convolution Properties

Theorem 6.1. Let the functions $f_j(z)$, for $j = 1, 2$, defined by (5.6) be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$. Then $(f_1 * f_2)(z) \in \mathcal{H}_p(\phi, \beta, \gamma)$, where

$$\phi = 1 - \frac{p\beta(1+\gamma)(1-\alpha)^2}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}, \tag{6.1}$$

such that $\Phi_0(n, \lambda, \sigma, \mu, \omega, p)$ is given by (3.3). The result is sharp for the functions $f_j(z)$, for $j = 1, 2$, given by

$$f_j(z) = \frac{1}{z^p} + \frac{p\beta(1+\gamma)(1-\alpha)}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}z^p. \tag{6.2}$$

Proof . Employing the technique used earlier by Schlid and Silverman[5], we will find the largest ϕ such that

$$\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\phi)} a_{p+k,1} a_{p+k,2} \leq 1, \tag{6.3}$$

for $f_j(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, for $j = 1, 2$. Since $f_j(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, for $j = 1, 2$, we readily see that

$$\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} a_{p+k,j} \leq 1, \tag{6.4}$$

by applying the Cauchy-Schwarz inequality, we obtain

$$\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \sqrt{a_{p+k,1} a_{p+k,2}} \leq 1. \tag{6.5}$$

This implies that we need only to show that

$$\frac{a_{p+k,1} a_{p+k,2}}{(1-\phi)} \leq \frac{\sqrt{a_{p+k,1} a_{p+k,2}}}{(1-\alpha)}, \tag{6.6}$$

or equivalently that

$$\sqrt{a_{p+k,1} a_{p+k,2}} \leq \frac{1-\phi}{(1-\alpha)}. \tag{6.7}$$

Hence by the inequality (6.5), it is sufficient to prove that

$$\frac{p\beta(1+\gamma)(1-\alpha)}{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)} \leq \frac{1-\phi}{1-\alpha}, \tag{6.8}$$

which implies that

$$\phi \leq 1 - \frac{p\beta(1+\gamma)(1-\alpha)^2}{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}. \quad (6.9)$$

Now, define the function $\Lambda(k)$ by

$$\Lambda(k) = 1 - \frac{p\beta(1+\gamma)(1-\alpha)^2}{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}. \quad (6.10)$$

We note that $\Lambda(k)$ is an increasing function of k , therefore we conclude that

$$\phi \leq \Lambda(0) = 1 - \frac{p\beta(1+\gamma)(1-\alpha)^2}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}, \quad (6.11)$$

therefore the proof is complete. \square

Theorem 6.2. Let the functions $f_1(z), f_2(z)$ defined by (5.6) be in the classes $\mathcal{H}_p(\alpha_1, \beta, \gamma), \mathcal{H}_p(\alpha_2, \beta, \gamma)$, respectively. Then $(f_1 * f_2)(z) \in \mathcal{H}_p(\eta, \beta, \gamma)$, where

$$\eta = 1 - \frac{p\beta(1+\gamma)(1-\alpha_1)(1-\alpha_2)}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}, \quad (6.12)$$

such that $\Phi_0(n, \lambda, \sigma, \mu, \omega, p)$ is given by (3.3), and the result is sharp for the functions $f_j(z) (j = 1, 2)$ given by

$$f_1(z) = \frac{1}{z^p} + \frac{p\beta(1+\gamma)(1-\alpha_1)}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)} z^p, \quad (6.13)$$

and

$$f_2(z) = \frac{1}{z^p} + \frac{p\beta(1+\gamma)(1-\alpha_2)}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)} z^p. \quad (6.14)$$

Proof . Use the arguments similar to those in the proof of Theorem 6.1. \square

Theorem 6.3. If $f_1(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k,1} z^{p+k}$ and $f_2(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} a_{p+k,2} z^{p+k}$ be in $\mathcal{H}_p(\alpha, \beta, \gamma)$, where $0 \leq a_{p+k,2} \leq 1$, $k = 0, 1, 2, \dots$, and $p \in \mathbb{N}$, then $(f_1 * f_2)(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$.

Proof . Since

$$\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} a_{p+k,1} a_{p+k,2} \leq \sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} a_{p+k,1} \leq 1,$$

then by Theorem 2.1, $(f_1 * f_2)(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$. \square

Corollary 6.4. If $f(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, then the integral operator

$$\mathcal{F}_{c,p}(z) = \frac{c}{z^{p+c}} \int_0^z t^{c+p-1} f(t) dt, \quad c > 0, \quad (6.15)$$

is also in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$.

Proof . It is easy to check that

$$\mathcal{F}_{c,p}(z) = f(z) * \left(\frac{1}{z^p} + \sum_{k=0}^{\infty} \frac{c}{c+2p+k} z^{p+k} \right). \quad (6.16)$$

Since $0 < \frac{c}{c+2p+k} \leq 1$, by Theorem 6.3, the proof is trivial. \square

Theorem 6.5. Let the functions $f_j(z)(j = 1, 2)$ defined by (5.6) be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$, and

$$p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p) - 2p\beta(1 + \gamma)(1 - \alpha) \geq 0, \tag{6.17}$$

then the function $h(z)$ defined by

$$h(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} (a_{p+k,1}^2 + a_{p+k,2}^2)z^{p+k}, \tag{6.18}$$

belongs to the class $\mathcal{H}_p(\alpha, \beta, \gamma)$, where $\Phi_0(n, \lambda, \sigma, \mu, \omega, p)$ is given by (3.3).

Proof . Since $f_1(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, we have

$$\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} a_{p+k,1} \leq 1, \tag{6.19}$$

and so

$$\sum_{k=0}^{\infty} \left[\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \right]^2 a_{p+k,1}^2 \leq 1. \tag{6.20}$$

Similarly, since $f_2(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, we have

$$\sum_{k=0}^{\infty} \left[\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \right]^2 a_{p+k,2}^2 \leq 1. \tag{6.21}$$

Hence

$$\sum_{k=0}^{\infty} \frac{1}{2} \left[\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \right]^2 (a_{p+k,1}^2 + a_{p+k,2}^2) \leq 1. \tag{6.22}$$

In view of Theorem 2.1, it is sufficient to show that

$$\sum_{k=0}^{\infty} \left[\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \right] (a_{p+k,1}^2 + a_{p+k,2}^2) \leq 1. \tag{6.23}$$

Thus the inequality (6.23) holds if, for $k = 0, 1, 2, \dots$

$$\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \leq \frac{1}{2} \left[\frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \right]^2, \tag{6.24}$$

or equivalently if

$$(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p) - 2p\beta(1+\gamma)(1-\alpha) \geq 0, \tag{6.25}$$

for $k = 0, 1, 2, \dots$. The left hand side of (6.25) is an increasing function of k , hence it holds for all k if

$$p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p) - 2p\beta(1 + \gamma)(1 - \alpha) \geq 0, \tag{6.26}$$

which is true by our assumption. Hence the proof is complete. \square

Theorem 6.6. Let the functions $f_j(z)(j = 1, 2)$ defined by (5.6), be in the class $\mathcal{H}_p(\alpha, \beta, \gamma)$. Then the function $h(z)$ defined by (6.18) belongs in the class $\mathcal{H}_p(\tau, \beta, \gamma)$, where

$$\tau = 1 - \frac{2p\beta(1 + \gamma)(1 - \alpha)^2}{p(p + \beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}, \tag{6.27}$$

such that $\Phi_0(n, \lambda, \sigma, \mu, p)$ is given by (3.3). The result is sharp for the functions $f_j(z)(j = 1, 2)$ defined by (6.2),

Proof . Since

$$\sum_{k=0}^{\infty} \frac{[(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)]^2}{[p\beta(1+\gamma)(1-\alpha)]^2} a_{p+k,j}^2 \leq \left[\sum_{k=0}^{\infty} \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} a_{p+k,j} \right]^2 \leq 1, \quad (6.28)$$

for $f_j(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, where $j = 1, 2$, so we get

$$\sum_{k=0}^{\infty} \frac{[(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)]^2}{2[p\beta(1+\gamma)(1-\alpha)]^2} (a_{p+k,1}^2 + a_{p+k,2}^2) \leq 1. \quad (6.29)$$

We have to find the largest τ such that

$$\frac{1}{1-\tau} \leq \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{2p\beta(1+\gamma)(1-\alpha)^2}, \quad (6.30)$$

that is

$$\tau \leq 1 - \frac{2p\beta(1+\gamma)(1-\alpha)^2}{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}. \quad (6.31)$$

If we define $L(k)$ by

$$L(k) = 1 - \frac{2p\beta(1+\gamma)(1-\alpha)^2}{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}, \quad (6.32)$$

that it is easy to see that $L(k)$ is an increasing function of k , thus we conclude that

$$\tau \leq L(0) = 1 - \frac{2p\beta(1+\gamma)(1-\alpha)^2}{p(p+\beta\gamma)\Phi_0(n, \lambda, \sigma, \mu, \omega, p)}, \quad (6.33)$$

which completes the proof. \square

7 Neighborhoods and Partial Sums

Definition 7.1. For every $\delta > 0$ and a non-negative sequence $\mathcal{S} = \{s_k\}_{k=0}^{\infty}$, where

$$s_k = \frac{(p+k)(p+k+\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1+\gamma)(1-\alpha)} \quad (7.1)$$

$$(k \geq 0, p \in \mathbb{N}, 0 \leq \alpha < 1, 0 < \beta \leq 1, 0 \leq \gamma \leq 1, \lambda \geq 0, 0 \leq \mu \leq \sigma),$$

the δ -neighborhood of a function $f \in \Sigma_p$ is defined by

$$\mathcal{N}_\delta(f) = \left\{ g \in \Sigma_p : g(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} b_{p+k} z^{p+k} \text{ and } \sum_{k=0}^{\infty} s_k |b_{p+k} - a_{p+k}| \leq \delta \right\}. \quad (7.2)$$

Theorem 7.2. Let $f \in \mathcal{H}_p(\alpha, \beta, \gamma)$ be given by (1.1). If f satisfies

$$\frac{f(z) + \epsilon z^{-p}}{1 + \epsilon} \in \mathcal{H}_p(\alpha, \beta, \gamma) \quad (\epsilon \in \mathbb{C}, |\epsilon| < \delta, \delta > 0), \quad (7.3)$$

then

$$\mathcal{N}_\delta(f) \subset \mathcal{H}_p(\alpha, \beta, \gamma). \quad (7.4)$$

Proof . It is not difficult to see that a function f belongs to $\mathcal{H}_p(\alpha, \beta, \gamma)$ if and only if

$$\frac{z^{p+2}(D_{\lambda, \sigma, \mu, \delta, p}^n f(z))'' + z^{p+1}(D_{\lambda, \sigma, \mu, \delta, p}^n f(z))' - p^2}{\beta\gamma z^{p+1}(D_{\lambda, \sigma, \mu, \delta, p}^n f(z))' + \beta\alpha(1 + \gamma)p - \beta p} \neq \nu \quad (\nu \in \mathbb{C}, |\nu| = 1), \quad (7.5)$$

which is equivalent to

$$\frac{(f * h)(z)}{z^{-p}} \neq 0, \quad (7.6)$$

where $h(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} c_{p+k} z^{p+k}$, such that

$$c_{p+k} = \frac{(p+k)(p+k - \nu\beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\nu\beta(1 + \gamma)(1 - \alpha)}, \quad \text{and} \quad |c_{p+k}| \leq \frac{(p+k)(p+k + \beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1 + \gamma)(1 - \alpha)}.$$

Furthermore, under the hypotheses (7.3), using (7.6) we obtain

$$\frac{1}{z^{-p}} \left(\frac{f(z) + \epsilon z^{-p}}{1 + \epsilon} * h(z) \right) \neq 0. \quad (7.7)$$

Now assume that $\left| \frac{(f * h)(z)}{z^{-p}} \right| < \delta$. Then by (7.7) we have

$$\left| \frac{1}{1 + \epsilon} \frac{(f * h)(z)}{z^{-p}} + \frac{\epsilon}{1 + \epsilon} \right| \geq \frac{|\epsilon|}{|1 + \epsilon|} - \frac{1}{|1 + \epsilon|} \left| \frac{(f * h)(z)}{z^{-p}} \right| > \frac{|\epsilon| - \delta}{|1 + \epsilon|} > 0.$$

This is a contradiction as $|\epsilon| < \delta$. Therefore $\left| \frac{(f * h)(z)}{z^{-p}} \right| \geq \delta$. If we now let

$$g(z) = \frac{1}{z^p} + \sum_{k=0}^{\infty} b_{p+k} z^{p+k} \in \mathcal{N}_\delta(f),$$

then we have

$$\begin{aligned} \delta - \left| \frac{(g * h)(z)}{z^{-p}} \right| &\leq \left| \frac{(f - g) * h(z)}{z^{-p}} \right| = \left| \sum_{k=0}^{\infty} (a_{p+k} - b_{p+k}) c_{p+k} z^{2p+k} \right| \\ &\leq \sum_{k=0}^{\infty} |a_{p+k} - b_{p+k}| |c_{p+k}| |z|^{2p+k} \\ &< \sum_{k=0}^{\infty} \frac{(p+k)(p+k + \beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1 + \gamma)(1 - \alpha)} |a_{p+k} - b_{p+k}| \\ &\leq \delta. \end{aligned}$$

Thus, $\frac{(g * h)(z)}{z^{-p}} \neq 0$ which implies that $g(z) \in \mathcal{H}_p(\alpha, \beta, \gamma)$, and the proof is complete. \square

Theorem 7.3. Let $f \in \Sigma_p$ be given by (1.1) and the partial sums $k_0(z)$ and $k_q(z)$ be defined by $k_0(z) = \frac{1}{z^p}$ and $k_q(z) = \frac{1}{z^p} + \sum_{k=0}^{q-1} a_{p+k} z^{p+k}$ ($q > 0$), also suppose that

$$\sum_{k=0}^{\infty} \theta_{p+k} a_{p+k} \leq 1 \quad (7.8)$$

where $\theta_{p+k} = \frac{(p+k)(p+k + \beta\gamma)\Phi_k(n, \lambda, \sigma, \mu, \omega, p)}{p\beta(1 + \gamma)(1 - \alpha)}$. Then, for $q > 0$, we have

$$\Re \left\{ \frac{f(z)}{k_q(z)} \right\} > 1 - \frac{1}{\theta_q} \quad (7.9)$$

and

$$\Re\left\{\frac{k_q(z)}{f(z)}\right\} > \frac{\theta_q}{1 + \theta_q}. \quad (7.10)$$

Proof . Under the hypotheses of the theorem we can see from (7.8) that

$$\theta_{p+k+1} > \theta_{p+k} > 1 \quad (k = 0, 1, 2, \dots).$$

Therefore making use of (7.8) again we have

$$\sum_{k=0}^{q-1} a_{p+k} + \theta_q \sum_{k=q}^{\infty} a_{p+k} \leq \sum_{k=0}^{\infty} \theta_{p+k} a_{p+k} \leq 1. \quad (7.11)$$

Let

$$w(z) = \theta_q \left[\frac{f(z)}{k_q(z)} - \left(1 - \frac{1}{\theta_q}\right) \right] = 1 + \frac{\theta_q \sum_{k=q}^{\infty} a_{p+k} z^{2p+k}}{1 + \sum_{k=0}^{q-1} a_{p+k} z^{2p+k}}. \quad (7.12)$$

Applying (7.11) and (7.12) we find

$$\left| \frac{w(z) - 1}{w(z) + 1} \right| = \left| \frac{\theta_q \sum_{k=q}^{\infty} a_{p+k} z^{2p+k}}{2 + 2 \sum_{k=0}^{q-1} a_{p+k} z^{2p+k} + \theta_q \sum_{k=q}^{\infty} a_{p+k} z^{2p+k}} \right| \leq \frac{\theta_q \sum_{k=q}^{\infty} a_{p+k}}{2 - 2 \sum_{k=0}^{q-1} a_{p+k} - \theta_q \sum_{k=q}^{\infty} a_{p+k}} \leq 1, \quad (7.13)$$

which shows that $\Re w(z) > 0$. From (7.12) we immediately obtain (7.9). Similarly by letting

$$\varphi(z) = (1 + \theta_q) \left[\frac{k_q(z)}{f(z)} - \frac{\theta_q}{1 + \theta_q} \right],$$

we can prove (7.10), therefore the proof is complete. \square

References

- [1] M.K. Aouf, *A class of meromorphic multivalent functions with positive coefficients*, Taiwan. J. Math. **12** (2008), 2517–2533.
- [2] W.G. Atshan and S.R. Kulkarni, *On application of differential subordination for certain subclass of meromorphically p -valent functions with positive coefficients defined by linear operator*, J. Ineq. Pure Appl. Math. **10** (2009), 1–11.
- [3] Sh. Najafzadeh and A. Ebadian, *Convex family of meromorphically multivalent functions on connected sets*, Math. Com. Mod. **57** (2013), 301–305.
- [4] H. Orhan, D. Raducanu, and E. Deniz, *Subclasses of meromorphically multivalent functions defined by a differential operator*, Comput. Math. Appl. **61** (2011), 966–979.
- [5] A. Schlid and H. Silverman, *Convolution of univalent functions with negative coefficients*, Ann. Univ. Curie-Sklodowska Sect. A **29** (1975), 99–107.
- [6] P.M. Sharma, *Iterative combinations for Srivastava-Gupta operators*, Asian-Eur. J. Math. **14** (2021), no. 7, 2150108.
- [7] H.M. Srivastava and V. Gupta, *A family of summation-integral type operators*, Math. Comput. Model. **37** (2003), no. 12-13, 1307–1315.