

An application to a system of fractional hybrid differential equations of best proximity point (pair) theorems via measure of noncompactness

Mallika Sarmah*, Anupam Das, Dipak Sarma

Department of Mathematics, Cotton University, Panbazar, Guwahati-781001, Assam, India

(Communicated by Mugur Alexandru Acu)

Abstract

This paper establishes the best proximity point (BPP) theorem by using a newly developed contraction operator. Using the results, the optimal solution to a system of fractional hybrid differential equations is then investigated. To further illustrate the findings, an additional example is given.

Keywords: Measure of noncompactness (MNC), Hybrid differential equation, Best proximity point (BPP)
2020 MSC: 47H09, 47H10, 45E99

1 Introduction

Finding the points that are closest to a given point or subgroup is one of the main issues in approximation theory. For \mathcal{W} , a normed linear space (NLS), if $\check{e} \cap T(\check{e}) \neq \phi$, where $\check{e} \in \mathcal{E}$ and \mathcal{E} is a non-empty subset of \mathcal{W} , then the map $T : \mathcal{E} \rightarrow \mathcal{W}$ has a fixed point. If there isn't a fixed point for T , then for any \check{e} in \mathcal{E} , the distance between \check{e} and $T(\check{e})$ is positive. If there is a minimum distance between \check{e} and $T(\check{e})$, then \check{e} is the best proximity point of T . As a consequence, Ky Fan develops his best approximation theory. On the other hand, when T maps \mathcal{E} into a distinct subset \mathcal{B} of \mathcal{W} , there is an issue. In this case, the extension of the problem is to find a point that approximates the distance between these two subsets. These points are known as best proximity points.

The structure of this paper will be as follows: We first review some basic terms and ideas related to best proximity theory. We then prove the best proximity point theorem for both cyclic and noncyclic contractive operators. We next go over their particular cases in the next section. Lastly, we use our findings to look at the best way to solve a system of fractional hybrid differential equations.

After Kuratowski and Hausdorff's generalization, numerous scholars investigated the concept of *MNC* in order to obtain significant extensions of the theory of compact operators. The main competence is the application of measures of noncompactness to ensure that the mappings satisfy the relevant inequalities. To help the reader understand our problem and objective, we thus provide a brief history. We review the standard fixed point problem in a Banach space \mathcal{W} by using certain regularity assumptions from Schauder [2].

*Corresponding author

Email addresses: mallikasarmah29@gmail.com (Mallika Sarmah), math.anupam@gmail.com (Anupam Das), dipak.sarma@cottonuniversity.ac.in (Dipak Sarma)

The authors of [10] first addressed the *BPP* results using *MNC*. After that, they used their findings to look into the existence of optimum solutions of a system of second-order differential equations. The application of *MNC* to find optimal solutions for ψ -Hilfer fractional differential equations with initial conditions was investigated by the authors of [11]. Using the Caputo fractional derivative of order $\check{r} \in (0, 1]$, the authors of [5] defined both strong and mild solutions to the fractional hybrid boundary value problems in two types and investigated the presence of at least one mild solution for each type. Finally, they proved how crucial it is to consider the initial conditions. In [13], the authors used Petryshyn’s fixed point theorem and *MNC* to investigate the solvability of functional-integral equations in Banach algebra. Additionally, they looked into the general class of functional equations, which includes a number of integral equations that arise in real-world and non-linear analysis situations. Using a Darbo-type fixed point theorem and *MNC*, the authors of [9] investigate the existence of solutions to infinite systems of second-order differential equations in the Banach sequence space l_p . Using a method related to *MNC*, the authors of [12] work on existence results for an infinite system of differential equations of order n with boundary conditions in the Banach spaces c_0 and l_1 . In [7], the authors used a method related to *MNC* and the generalized Meir–Keeler fixed point theorem to demonstrate the existence of a solution for an infinite system of nonlinear integral equations in the Banach spaces $l_p, p > 1$. The authors examined if there are any solutions for nonlinear integral equations in [8]. Additionally, they presented an iteration approach to provide very accurate solutions for the nonlinear integral equations. At last, they determined the convergence condition and provided an upper bound of error.

Motivated by these works, in this paper we established a best proximity point theorem with the help of a newly defined contraction operator by using *MNC*, and studied the existence of optimum solutions of a system of fractional hybrid differential equations.

Theorem 1.1. [6] A continuous operator $J : K \rightarrow K$ where K is a nonempty, convex and compact subset of a Banach space \check{W} , then J admits at least a fixed point.

Clearly it is the generalization of Brouwer fixed point theorem. Consider a Banach space \check{W} and a closed ball $\mathcal{C}(\bar{f}, \bar{g}) = \{\bar{m} \in \check{W} : \|\bar{m} - \bar{f}\| \leq \bar{g}\}$ in \check{W} . Suppose \bar{z} , for nonempty (z) denotes the closure of z and $\overline{conv}(z)$ denotes the closed and convex hull of the non empty set z which is the smallest convex and closed set containing z .

Also $M_{\check{W}}$ and $N_{\check{W}}$ represents the family of non empty bounded subsets of \check{W} and subfamily of \check{W} consisting all relatively compact sets respectively; $R = (-\infty, \infty)$, and $R_+ = [0, \infty)$. Measure of noncompactness (*MNC*) is defined axiomatically as follows:

Definition 1.2. [1] A map $\check{G} : M_{\check{W}} \rightarrow R_+$ is a *MNC* (measure of noncompactness) in a Banach space \check{W} , if the following conditions are holds for \check{G} :

1. $\ker \check{G} = \{\check{\mathfrak{X}} \in M_{\check{W}} : \check{G}(\check{\mathfrak{X}}) = 0\} \neq \phi$,
2. $\check{\mathfrak{X}} \in \ker \check{G}$ if and only if $\check{\mathfrak{X}}$ is relatively compact,
3. $\check{\mathfrak{X}}_1 \subseteq \check{\mathfrak{X}}_2 \Rightarrow \check{G}(\check{\mathfrak{X}}_1) \leq \check{G}(\check{\mathfrak{X}}_2)$,
4. $\check{G}(\overline{\check{\mathfrak{X}}}) = \check{G}(\check{\mathfrak{X}})$,
5. $\check{G}(\overline{conv}(\check{\mathfrak{X}})) = \check{G}(\check{\mathfrak{X}})$,
6. $\check{G}(\delta\check{\mathfrak{X}}_1 + (1 - \delta)\check{\mathfrak{X}}_2) \leq \delta\check{G}(\check{\mathfrak{X}}_1) + (1 - \delta)\check{G}(\check{\mathfrak{X}}_2)$, for $\delta \in [0, 1]$,
7. $\max\{\check{G}(\check{\mathfrak{X}}_1), \check{G}(\check{\mathfrak{X}}_2)\} = \check{G}(\check{\mathfrak{X}}_1 \cup \check{\mathfrak{X}}_2)$,
8. The set $\check{\mathfrak{X}}_\infty = \bigcap_{n=1}^\infty \check{\mathfrak{X}}_n$ is compact and non empty, if $(\check{\mathfrak{X}}_n)$ is a decreasing sequence of closed sets which are non empty in $M_{\check{W}}$ and $\lim_{n \rightarrow \infty} \check{G}(\check{\mathfrak{X}}_n) = 0$.

In particular, the space $\check{W} = \mathcal{C}(I)$, where I is the closed and bounded interval, is the set of real valued continuous functions on I . Then \check{W} is a Banach space with the norm $\|D\| = \sup\{|D(\check{s})| : \check{s} \in I\}$, $D \in \check{W}$. Assume that $U(\neq \phi) \subseteq \check{W}$ is bounded. For $D \in U$ and $r > 0$, the modulus of continuity of D , represented by $\beth(D, r)$ i.e.,

$$\beth(D, r) = \sup\{|D(\check{s}_1) - D(\check{s}_2)| : \check{s}_1, \check{s}_2 \in I, |\check{s}_1 - \check{s}_2| \leq r\}.$$

Furthermore, we define

$$\beth(U, r) = \sup\{\beth(D, r) : D \in U\}; \beth_0(U) = \lim_{r \rightarrow 0} \beth(U, r).$$

A Hausdorff *MNC* \beth is given by $\beth(U) = \frac{1}{2}\beth_0(U)$, see [2]. It is widely known that the map \beth_0 is a *MNC* in \check{W} .

2 Preliminaries

We collect some fundamental definitions and notations needed for the paper.

Definition 2.1. [6] Consider \check{W} be a Banach space. Then

1. \check{W} is uniformly convex Banach space if there exists a strictly increasing function $\bar{D} : (0, 2] \rightarrow [0, 1]$ such that,

$$\begin{cases} \|\acute{r}_0 - \bar{a}_0\| \leq \bar{\omega}, \\ \|\acute{k}_0 - \bar{a}_0\| \leq \bar{\omega}, \implies \left\| \frac{\acute{r}_0 + \acute{k}_0}{2} - \bar{a}_0 \right\| \leq \left(1 - \bar{D} \left(\frac{\bar{s}_1}{\bar{\omega}} \right) \right) \bar{\omega}; \\ \|\acute{r}_0 - \acute{k}_0\| \geq \bar{s}_1 \end{cases}$$

for all $\acute{r}_0, \acute{k}_0, \bar{a}_0 \in \check{W}$, $\bar{\omega} > 0$ and $\bar{s}_1 \in [0, 2\bar{\omega}]$.

2. \check{W} is strictly convex Banach space if for $\acute{r}_0, \acute{k}_0, \bar{a}_0 \in \check{W}$ and $\bar{\omega} > 0$, the following conditions are holds:

$$\begin{cases} \|\acute{r}_0 - \bar{a}_0\| \leq \bar{\omega}, \\ \|\acute{k}_0 - \bar{a}_0\| \leq \bar{\omega}, \implies \left\| \frac{\acute{r}_0 + \acute{k}_0}{2} - \bar{a}_0 \right\| < \bar{\omega}. \\ \acute{r}_0 \neq \acute{k}_0. \end{cases}$$

Consider a normed linear space (NLS) \check{W} . For any two non empty subset N_1, N_2 of \check{W} , the pair (N_1, N_2) is closed if and only if both N_1, N_2 are closed; $(N_1, N_2) \subseteq (H, C)$ if and only if $N_1 \subseteq H, N_2 \subseteq C$. In addition, we denote, $dist(H, C) = \inf \{ \|\check{\mu} - \check{\nu}\| : (\check{\mu}, \check{\nu}) \in H \times C \}$,

$$H_0 = \{ \check{\mu} \in H : \text{there exists } \bar{\nu}_0 \in C, \|\check{\mu} - \bar{\nu}_0\| = dist(H, C) \}$$

$$C_0 = \{ \check{\nu} \in C : \text{there exists } \check{\mu}_1 \in H, \|\check{\mu}_1 - \check{\nu}\| = dist(H, C) \}.$$

Definition 2.2. [6] Consider \mathfrak{Z} be a NLS. A non empty pair (H, C) of \mathfrak{Z} is proximal if $H = H_0$ and $C = C_0$. Also for a reflexive Banach space \mathfrak{D} , if the pair (H, C) be a closed, nonempty, convex and bounded in \mathfrak{D} , then (H_0, C_0) is also convex, nonempty and closed pair. Consider a function $T : H \cup C \rightarrow H \cup C$. We say that T is,

1. relatively nonexpansive, if $\|T(\check{\mu}) - T(\check{\nu})\| \leq \|\check{\mu} - \check{\nu}\|$ for any $(\check{\mu}, \check{\nu}) \in H \times C$,
2. cyclic, if $T(H) \subseteq C$ and $T(C) \subseteq H$,
3. non cyclic, if $T(H) \subseteq H$ and $T(C) \subseteq C$,
4. compact, if $(\overline{T(H)}, \overline{T(C)})$ is compact.

Definition 2.3. [6] Consider (H, C) be a nonempty pair in a Banach space \check{W} and $\bar{E} : H \cup C \rightarrow H \cup C$ be a cyclic function, then $\check{q} \in H \cup C$ is called a BPP of \bar{E} if $\|\check{q} - \bar{E}(\check{q})\| = dist(H, C)$. If \bar{E} is non cyclic, then the pair $(\check{q}, \check{w}) \in H \times C$ is best proximity pair if $\|\check{q} - \check{w}\| = dist(H, C)$, for $\check{q} = \bar{E}(\check{q}), \check{w} = \bar{E}(\check{w})$.

Corollary 2.4. [6] Suppose a Banach space \check{W} and a nonempty, convex and compact pair (N_1, N_2) in \check{W} . Let a cyclic and relatively nonexpansive mapping $T : N_1 \cup N_2 \rightarrow N_1 \cup N_2$. Then T have a BPP.

Corollary 2.5. [6] Suppose a strictly convex Banach space \check{W} and a compact, nonempty and convex pair (N_1, N_2) in \check{W} . Let a relatively non-expansive and non-cyclic mapping $T : N_1 \cup N_2 \rightarrow N_1 \cup N_2$. Then T have a best proximity pair.

Definition 2.6. [5] The left Caputo fractional derivative of order $\check{r} > 0$ is defined by

$$D^{\check{r}} \check{F}(\tau_1) = \frac{1}{\check{\gamma}(m - \check{r})} \int_a^{\tau_1} (\tau_1 - \check{\varrho})^{m - \check{r} - 1} D^m \check{F}(\check{\varrho}) d\check{\varrho}, \tag{2.1}$$

where $m - 1 < \check{r} < m$, $D = \frac{d}{d\check{\varrho}}$ and $\check{\gamma}(\cdot)$ denotes Euler's gamma function.

3 Main result

Definition 3.1. [3] Consider the function $\mathbf{C} : \mathfrak{U} \subseteq \check{\mathcal{W}} \rightarrow \check{\mathcal{W}}$ and $\check{\Phi} : M_{\check{\mathcal{W}}} \rightarrow R^+$. \mathbf{C} is said to be $\check{\Phi}$ -admissible if,

$$\check{\Phi}(\check{z}) \geq 1 \implies \check{\Phi}(\text{Conv}\mathbf{C}\check{z}) \geq 1,$$

where $\check{z} \subseteq \mathfrak{U}$ and $\check{z}, \mathbf{C}\check{z} \in M_{\check{\mathcal{W}}}$.

Definition 3.2. [3] Assume the function $\check{\omega} : R^+ \times R^+ \rightarrow R^+$ with the following conditions:

- (i) $\max\{\check{\mu}_0, \check{\nu}_0\} \leq \check{\omega}(\check{\mu}_0, \check{\nu}_0)$, for all $\check{\mu}_0, \check{\nu}_0 \geq 0$,
- (ii) $\check{\omega}$ is non-decreasing and continuous,
- (iii) $\check{\omega}(\check{u}_0 + \check{b}_0, \check{m}_0 + \check{t}_0) \leq \check{\omega}(\check{u}_0, \check{m}_0) + \check{\omega}(\check{b}_0, \check{t}_0)$.

We use \mathbb{Q} to represent the collection of this functions. For example, $\check{\omega}(\check{\mu}, \check{\nu}) = \check{\mu} + \check{\nu}$.

Definition 3.3. [3] Assume a map $\check{\varrho} : R^+ \rightarrow R^+$ with a non negative sequence $\{\check{a}_n\}$ such that

$$\lim_{n \rightarrow \infty} \check{\varrho}(\check{a}_n) = 0 \quad \text{if} \quad \lim_{n \rightarrow \infty} \check{a}_n = 0.$$

We use \mathbb{K} to represent the collection of this functions. For example, $\check{\varrho}(\check{\mu}) = \check{\mu}$. Assume that a map $\check{\Theta} : R^+ \rightarrow R^+$ and $\check{\varrho} \in \mathbb{K}$ with the following conditions:

- (i) $\check{\Theta}$ is continuous with $\check{\Theta}(\check{\mu}) = 0$ if and only if $\check{\mu} = 0$.
- (ii) $\lim_{n \rightarrow \infty} \check{\varrho}(\check{\mu}_n) < \check{\Theta}(\check{\mu})$ if $\lim_{n \rightarrow \infty} \check{\mu}_n = \check{\mu} > 0$.

We use $\mathbb{S}_{\check{\varrho}}$ to represent the collection of this class of functions. \mathbf{H} and \mathbf{C} will be nonempty convex subsets of a Banach space $\check{\mathcal{W}}$ in this section.

Definition 3.4. Consider (\mathbf{H}, \mathbf{C}) be convex and nonempty pair in a Banach space $\check{\mathcal{W}}$ with a *MNC* $\check{\mathcal{G}}$ on $\check{\mathcal{W}}$. A mapping $\mathbf{T} : \mathbf{H} \cup \mathbf{C} \rightarrow \mathbf{H} \cup \mathbf{C}$ which is cyclic (*noncyclic*), is said to be a $(\check{\Theta}, \check{\omega}, \check{F}, \check{\varrho})$ -contractive operator such that for any convex, nonempty, proximal, closed, bounded and \mathbf{T} -invariant pair (ι_1, ι_2) such that $\text{dist}(\iota_1, \iota_2) = \text{dist}(\mathbf{H}, \mathbf{C})$, we have,

$$\check{\Phi}(\check{z}) \check{\Theta} [\check{\omega}(\check{\mathcal{G}}(\mathbf{T}(\iota_1) \cup \mathbf{T}(\iota_2)), \check{F}(\check{\mathcal{G}}(\mathbf{T}(\iota_1) \cup \mathbf{T}(\iota_2))))] \leq \check{\varrho} [\check{\omega}(\check{\mathcal{G}}(\iota_1 \cup \iota_2), \check{F}(\check{\mathcal{G}}(\iota_1 \cup \iota_2)))] , \quad (3.1)$$

where \check{z} is a non empty subset of $\check{\mathbb{H}} = \mathbf{H} \cup \mathbf{C}$, $\check{\Theta} \in \mathbb{S}_{\check{\varrho}}$, $\check{\omega} \in \mathbb{Q}$, $\check{\varrho} \in \mathbb{K}$ and $\check{F} : R^+ \rightarrow R^+$ is a non decreasing and continuous function. Also the function \mathbf{T} is $\check{\Phi}$ -admissible with $\check{\Phi}(\check{\mathbb{H}}) \geq 1$.

Theorem 3.5. Consider a relatively non expansive, cyclic and $(\check{\Theta}, \check{\omega}, \check{F}, \check{\varrho})$ -contractive operator $\mathbf{T} : \mathbf{H} \cup \mathbf{C} \rightarrow \mathbf{H} \cup \mathbf{C}$, then \mathbf{T} has a *BPP*, if $\mathbf{H}_0 \neq \phi$.

Proof . Since $\mathbf{H}_0 \neq \phi$, $(\mathbf{H}_0, \mathbf{C}_0) \neq \phi$. By the given conditions on \mathbf{T} , clearly $(\mathbf{H}_0, \mathbf{C}_0)$ is a closed, convex, proximal and \mathbf{T} -invariant pair. For each $\check{p}_0 \in \mathbf{H}_0$, there is a $\check{q}_0 \in \mathbf{C}_0$ satisfying $\|\check{p}_0 - \check{q}_0\| = \text{dist}(\mathbf{H}, \mathbf{C})$. Since \mathbf{T} is relatively non expansive, we get, $\|\mathbf{T}\check{p}_0 - \mathbf{T}\check{q}_0\| \leq \|\check{p}_0 - \check{q}_0\| = \text{dist}(\mathbf{H}, \mathbf{C})$, which implies $\mathbf{T}\check{p}_0 \in \mathbf{C}_0$, that is $\mathbf{T}(\mathbf{H}_0) \subseteq \mathbf{C}_0$. Similarly, $\mathbf{T}(\mathbf{C}_0) \subseteq \mathbf{H}_0$, hence we get \mathbf{T} is cyclic on $\mathbf{H}_0 \cup \mathbf{C}_0$.

Let us assume that $\check{\mathfrak{X}}_0 = \mathbf{H}_0$, $\check{\mathfrak{W}}_0 = \mathbf{C}_0$ and $\left\{ \left(\check{\mathfrak{X}}_n, \check{\mathfrak{W}}_n \right) \right\}$ be a sequence of pairs with $\check{\mathfrak{X}}_n = \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{X}}_{n-1}))$ and $\check{\mathfrak{W}}_n = \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{W}}_{n-1}))$, for all $n \in \mathbb{N}$. Now our claim is, $\check{\mathfrak{X}}_{n+1} \subseteq \check{\mathfrak{W}}_n$ and $\check{\mathfrak{W}}_n \subseteq \check{\mathfrak{X}}_{n-1}$ for all $n \in \mathbb{N}$. In fact $\check{\mathfrak{W}}_1 = \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{W}}_0)) = \overline{\text{conv}}(\mathbf{T}(\mathbf{C}_0)) \subseteq \overline{\text{conv}}(\mathbf{H}_0) = \mathbf{H}_0 = \check{\mathfrak{X}}_0$. Hence we can write,

$$\mathbf{T}(\check{\mathfrak{W}}_1) \subseteq \mathbf{T}(\check{\mathfrak{X}}_0) \quad \text{and} \quad \check{\mathfrak{W}}_2 = \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{W}}_1)) \subseteq \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{X}}_0)) = \check{\mathfrak{X}}_1.$$

With the similar argument, we get by using induction, $\check{\mathfrak{W}}_n \subseteq \check{\mathfrak{X}}_{n-1}$. Similarly, we get $\check{\mathfrak{X}}_{n+1} \subseteq \check{\mathfrak{W}}_n$, for all $n \in \mathbb{N}$. Hence we can write $\check{\mathfrak{X}}_{n+2} \subseteq \check{\mathfrak{W}}_{n+1} \subseteq \check{\mathfrak{X}}_n \subseteq \check{\mathfrak{W}}_{n-1}$, for all $n \in \mathbb{N}$. So in $\mathbf{H}_0 \times \mathbf{C}_0$, the decreasing sequence of *NBCC* pairs is $\left\{ \left(\check{\mathfrak{X}}_{2n}, \check{\mathfrak{W}}_{2n} \right) \right\}$. Moreover,

$$\mathbf{T}(\check{\mathfrak{W}}_{2n}) \subseteq \mathbf{T}(\check{\mathfrak{X}}_{2n-1}) \subseteq \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{X}}_{2n-1})) = \check{\mathfrak{X}}_{2n}, \quad (3.2)$$

$$\mathbf{T}(\check{\mathfrak{X}}_{2n}) \subseteq \mathbf{T}(\check{\mathfrak{W}}_{2n-1}) \subseteq \overline{\text{conv}}(\mathbf{T}(\check{\mathfrak{W}}_{2n-1})) = \check{\mathfrak{W}}_{2n}. \quad (3.3)$$

Hence we get $(\tilde{\mathfrak{X}}_{2n}, \mathfrak{W}_{2n})$ is T -invariant pair, for all $n \in \mathbb{N}$. Now, if the pair $(\check{x}, \check{y}) \in H_0 \times C_0$ is proximal, we have,

$$dist(\tilde{\mathfrak{X}}_{2n}, \mathfrak{W}_{2n}) \leq \|T^{2n}\check{x} - T^{2n}\check{y}\| \leq \|\check{x} - \check{y}\| = dist(H, C).$$

Now we are to show that, for all $n \in \mathbb{N}$, the pair $(\tilde{\mathfrak{X}}_n, \mathfrak{W}_n)$ is proximal. For $n=0$ we have $(\tilde{\mathfrak{X}}_0, \mathfrak{W}_0)$ is proximal pair. Let us assume that $(\tilde{\mathfrak{X}}_k, \mathfrak{W}_k)$ is proximal and an arbitrary $\tilde{\pi}$ such that $\tilde{\pi} \in \tilde{\mathfrak{X}}_{k+1} = \overline{conv}(T(\tilde{\mathfrak{X}}_k))$. So $\tilde{\pi} = \sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} T(J_{\acute{k}})$ with $J_{\acute{k}} \in \tilde{\mathfrak{X}}_k, \acute{n} \in [1, \infty), \Psi_{\acute{k}} \geq 0$ and $\sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} = 1$. By assumption, we have $(\tilde{\mathfrak{X}}_k, \mathfrak{W}_k)$ is proximity pair, there exists $\tilde{\rho}_{\acute{k}} \in \mathfrak{W}_k$ for $1 \leq \acute{k} \leq \acute{n}$ such that $\|J_{\acute{k}} - \tilde{\rho}_{\acute{k}}\| = dist(\tilde{\mathfrak{X}}_k, \mathfrak{W}_k) = dist(H, C)$. Consider $\tilde{\rho} = \sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} T(\tilde{\rho}_{\acute{k}})$. Then $\tilde{\rho} \in \overline{conv}(T(\mathfrak{W}_k)) = \mathfrak{W}_{k+1}$, and

$$\|\tilde{\pi} - \tilde{\rho}\| = \left\| \sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} T(J_{\acute{k}}) - \sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} T(\tilde{\rho}_{\acute{k}}) \right\| \leq \sum_{\acute{k}=1}^{\acute{n}} \Psi_{\acute{k}} \|J_{\acute{k}} - \tilde{\rho}_{\acute{k}}\| = dist(H, C). \tag{3.4}$$

Hence $(\tilde{\mathfrak{X}}_{k+1}, \mathfrak{W}_{k+1})$ is the proximal pair and by induction hypothesis our claim is proved.

If $\max\{\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n_0}), \check{\mathcal{G}}(\mathfrak{W}_{2n_0})\} = 0$, for some $n_0 \in [1, \infty) \cup \{0\}$, then we have $T : \tilde{\mathfrak{X}}_{2n_0} \cup \mathfrak{W}_{2n_0} \rightarrow \tilde{\mathfrak{X}}_{2n_0} \cup \mathfrak{W}_{2n_0}$ is compact. By Corollary 2.4, T have a *BPP*. Hence we consider that $\max\{\check{\mathcal{G}}(\tilde{\mathfrak{X}}_n), \check{\mathcal{G}}(\mathfrak{W}_n)\} > 0$, for all $n \in [1, \infty)$. Since $\tilde{\mathfrak{X}}_{2n+1} \subseteq T(\tilde{\mathfrak{X}}_{2n})$ and $\mathfrak{W}_{2n+1} \subseteq T(\mathfrak{W}_{2n})$, we have,

$$\begin{aligned} & \acute{\theta} \left[\check{\omega} \left(\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n+1} \cup \mathfrak{W}_{2n+1}), \check{F} \left(\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n+1} \cup \mathfrak{W}_{2n+1}) \right) \right) \right] \\ &= \acute{\theta} \left[\check{\omega} \left(\max\{\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n+1}), \check{\mathcal{G}}(\mathfrak{W}_{2n+1})\}, \check{F} \left(\max\{\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n+1}), \check{\mathcal{G}}(\mathfrak{W}_{2n+1})\} \right) \right) \right] \\ &= \acute{\theta} \left[\check{\omega} \left(\max\left\{ \check{\mathcal{G}}\left(\overline{conv}\left(T(\tilde{\mathfrak{X}}_{2n})\right)\right), \check{\mathcal{G}}\left(\overline{conv}\left(T(\mathfrak{W}_{2n})\right)\right) \right\}, \check{F} \left(\max\left\{ \check{\mathcal{G}}\left(\overline{conv}\left(T(\tilde{\mathfrak{X}}_{2n})\right)\right), \check{\mathcal{G}}\left(\overline{conv}\left(T(\mathfrak{W}_{2n})\right)\right) \right\} \right) \right) \right] \\ &= \acute{\theta} \left[\check{\omega} \left(\max\{\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n})), \check{\mathcal{G}}(T(\mathfrak{W}_{2n}))\}, \check{F} \left(\max\{\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n})), \check{\mathcal{G}}(T(\mathfrak{W}_{2n}))\} \right) \right) \right] \\ &= \acute{\theta} \left[\check{\omega} \left(\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n}) \cup T(\mathfrak{W}_{2n})), \check{F} \left(\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n}) \cup T(\mathfrak{W}_{2n})) \right) \right) \right] \\ &\leq \acute{\phi}(\check{\chi}) \acute{\theta} \left[\check{\omega} \left(\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n}) \cup T(\mathfrak{W}_{2n})), \check{F} \left(\check{\mathcal{G}}(T(\tilde{\mathfrak{X}}_{2n}) \cup T(\mathfrak{W}_{2n})) \right) \right) \right] \\ &\leq \acute{\varrho} \left[\check{\omega} \left(\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}), \check{F} \left(\check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}) \right) \right) \right]. \end{aligned}$$

Clearly, $\left\{ \check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}) \right\}_{n=1}^{\infty}$ is a decreasing and non-negative sequence. Thus, there exists $\mathcal{V} \geq 0$ such that $\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}) = \mathcal{V}$. If possible, consider $\mathcal{V} > 0$. Then we have as $n \rightarrow \infty$

$$\begin{aligned} \acute{\theta}[\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))] &\leq \acute{\varrho}[\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))] \\ &< \acute{\theta}[\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))], \end{aligned}$$

a contradiction. Thus $\mathcal{V}=0$, that is $\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}) = 0$. Thus

$$\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n} \cup \mathfrak{W}_{2n}) = \max\left\{ \lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathfrak{X}}_{2n}), \lim_{n \rightarrow \infty} \check{\mathcal{G}}(\mathfrak{W}_{2n}) \right\} = 0.$$

If $\tilde{\mathfrak{X}}_{\infty} = \cap_{n=0}^{\infty} \tilde{\mathfrak{X}}_{2n}$ and $\mathfrak{W}_{\infty} = \cap_{n=0}^{\infty} \mathfrak{W}_{2n}$, We get the non empty, compact, convex pair $(\tilde{\mathfrak{X}}_{\infty}, \mathfrak{W}_{\infty})$ which is T -invariant with $dist(\tilde{\mathfrak{X}}_{\infty}, \mathfrak{W}_{\infty}) = dist(H, C)$. Thus T has a *BPP*. \square

Theorem 3.6. Consider a relatively nonexpansive, noncyclic and $(\acute{\Theta}, \check{\omega}, \check{F}, \check{\varrho})$ -contractive operator $T : H \cup C \rightarrow H \cup C$ on a strictly convex Banach space \mathcal{W} , then T has a best proximity pair, if $H_0 \neq \phi$.

Proof . Following the proof of the Theorem 3.5, define a pair $(\tilde{\mathcal{X}}_n, \mathcal{W}_n)$ as $\tilde{\mathcal{X}}_n = \overline{conv} (T(\tilde{\mathcal{X}}_{n-1}))$ and $\mathcal{W}_n = \overline{conv} (T(\mathcal{W}_{n-1}))$, $n \in \mathbb{N}$ with $\tilde{\mathcal{X}}_0 = H_0$ and $\mathcal{W}_0 = C_0$. We get a NBCC and decreasing sequence of pair $\{(\tilde{\mathcal{X}}_n, \mathcal{W}_n)\}$ in $H_0 \times C_0$. Also,

$$T(\tilde{\mathcal{X}}_n) \subseteq T(\tilde{\mathcal{X}}_{n-1}) \subseteq \overline{conv} (T(\tilde{\mathcal{X}}_{n-1})) = \tilde{\mathcal{X}}_n, \tag{3.5}$$

$$T(\mathcal{W}_n) \subseteq T(\mathcal{W}_{n-1}) \subseteq \overline{conv} (T(\mathcal{W}_{n-1})) = \mathcal{W}_n. \tag{3.6}$$

Thus, the pair $(\tilde{\mathcal{X}}_n, \mathcal{W}_n)$ is T -invariant for all $n \geq 1$. Following the proof of the Theorem 3.5, we obtain a proximal pair $(\tilde{\mathcal{X}}_n, \mathcal{W}_n)$, for all non negative integer n such that $dist(\tilde{\mathcal{X}}_n, \mathcal{W}_n) = dist(H, C)$. Now if $\max\{\check{\mathcal{G}}(\tilde{\mathcal{X}}_{n_0}), \check{\mathcal{G}}(\mathcal{W}_{n_0})\} = 0$, for some positive integer n_0 , then $T : \tilde{\mathcal{X}}_{n_0} \cup \mathcal{W}_{n_0} \rightarrow \tilde{\mathcal{X}}_{n_0} \cup \mathcal{W}_{n_0}$ is compact. Hence from Corollary 2.5, we get the desired result. Thus we consider that, $max\{\check{\mathcal{G}}(\tilde{\mathcal{X}}_n), \check{\mathcal{G}}(\mathcal{W}_n)\} > 0$. Since $\tilde{\mathcal{X}}_{n+1} \subseteq T(\tilde{\mathcal{X}}_n)$ and $\mathcal{W}_{n+1} \subseteq T(\mathcal{W}_n)$, we have,

$$\begin{aligned} & \acute{\Theta} [\check{\omega} (\check{\mathcal{G}}(\tilde{\mathcal{X}}_{n+1} \cup \mathcal{W}_{n+1}), \check{F}(\check{\mathcal{G}}(\tilde{\mathcal{X}}_{n+1} \cup \mathcal{W}_{n+1})))] \\ &= \acute{\Theta} [\check{\omega} (\max\{\check{\mathcal{G}}(\tilde{\mathcal{X}}_{n+1}), \check{\mathcal{G}}(\mathcal{W}_{n+1})\}, \check{F}(\max\{\check{\mathcal{G}}(\tilde{\mathcal{X}}_{n+1}), \check{\mathcal{G}}(\mathcal{W}_{n+1})\}))] \\ &= \acute{\Theta} [\check{\omega} (\max\{\check{\mathcal{G}}(\overline{conv}(T(\tilde{\mathcal{X}}_n))), \check{\mathcal{G}}(\overline{conv}(T(\mathcal{W}_n)))\}, \check{F}(\max\{\check{\mathcal{G}}(\overline{conv}(T(\tilde{\mathcal{X}}_n))), \check{\mathcal{G}}(\overline{conv}(T(\mathcal{W}_n)))\})))] \\ &= \acute{\Theta} [\check{\omega} (\max\{\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n)), \check{\mathcal{G}}(T(\mathcal{W}_n))\}, \check{F}(\max\{\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n)), \check{\mathcal{G}}(T(\mathcal{W}_n))\})))] \\ &= \acute{\Theta} [\check{\omega} (\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n) \cup T(\mathcal{W}_n)), \check{F}(\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n) \cup T(\mathcal{W}_n))))] \\ &\leq \acute{\phi}(\check{\omega}) \acute{\Theta} [\check{\omega} (\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n) \cup T(\mathcal{W}_n)), \check{F}(\check{\mathcal{G}}(T(\tilde{\mathcal{X}}_n) \cup T(\mathcal{W}_n))))] \\ &\leq \check{\varrho} [\check{\omega} (\check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n), \check{F}(\check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n)))] . \end{aligned}$$

Clearly, $\{\check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n)\}_{n=1}^\infty$ is a decreasing and non-negative sequence. Thus, there exists $\mathcal{V} \geq 0$ such that $\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n) = \mathcal{V}$. If possible, consider $\mathcal{V} > 0$. Then we have

$$\begin{aligned} \acute{\Theta} [\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))] &\leq \check{\varrho} [\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))]; \text{ as } n \rightarrow \infty \\ &< \acute{\Theta} [\check{\omega}(\mathcal{V}, \check{F}(\mathcal{V}))], \end{aligned}$$

a contradiction. Thus $\mathcal{V} = 0$, that is $\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n) = 0$. Thus

$$\lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathcal{X}}_n \cup \mathcal{W}_n) = \max\left\{ \lim_{n \rightarrow \infty} \check{\mathcal{G}}(\tilde{\mathcal{X}}_n), \lim_{n \rightarrow \infty} \check{\mathcal{G}}(\mathcal{W}_n) \right\} = 0.$$

If $\tilde{\mathcal{X}}_\infty = \bigcap_{n=0}^\infty \tilde{\mathcal{X}}_n$ and $\mathcal{W}_\infty = \bigcap_{n=0}^\infty \mathcal{W}_n$, We get the non empty, compact, convex pair $(\tilde{\mathcal{X}}_\infty, \mathcal{W}_\infty)$ which is T -invariant with $dist(\tilde{\mathcal{X}}_\infty, \mathcal{W}_\infty) = dist(H, C)$. Thus T has a best proximity pair. \square

The next results are special cases of Theorem 3.5. The noncyclic version of the following corollaries are satisfied in strictly convex Banach spaces.

Corollary 3.7. Consider a cyclic relatively nonexpansive mapping $T : H \cup C \rightarrow H \cup C$ such that for any nonempty, bounded, closed, convex, proximal and T -invariant pair (ι_1, ι_2) with $dist(\iota_1, \iota_2) = dist(H, C)$,

$$2^{\check{\omega}} \acute{\Theta} [\check{\omega} (\check{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)), \check{F}(\check{\mathcal{G}}(T(\iota_1) \cup T(\iota_2))))] \leq \check{\varrho} [\check{\omega} (\check{\mathcal{G}}(\iota_1 \cup \iota_2), \check{F}(\check{\mathcal{G}}(\iota_1 \cup \iota_2)))] , \tag{3.7}$$

for $\check{\omega} \geq 0$. Then T has a BPP if $H_0 \neq \emptyset$.

Proof . Putting $\acute{\Phi}(\acute{w}) = 2^{\acute{w}}$, $\acute{w} \geq 0$ in equation (3.1) of Definition 3.4 and using Theorem 3.5, the result follows. \square

Corollary 3.8. Consider a cyclic relatively nonexpansive mapping $T : H \cup C \rightarrow H \cup C$ such that for any nonempty, bounded, closed, convex, proximal and T -invariant pair (ι_1, ι_2) with $\text{dist}(\iota_1, \iota_2) = \text{dist}(H, C)$,

$$2^{\acute{\Theta}} [\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)) + \acute{F}(\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)))] \leq \acute{\varrho} [\acute{\mathcal{G}}(\iota_1 \cup \iota_2) + \acute{F}(\acute{\mathcal{G}}(\iota_1 \cup \iota_2))]. \tag{3.8}$$

Then T has a *BPP* if $H_0 \neq \emptyset$.

Proof . Putting $\acute{\omega}(\acute{\mu}, \acute{\nu}) = \acute{\mu} + \acute{\nu}$ in equation (3.7) of Corollary 3.7 and using Theorem 3.5, the result follows. \square

Corollary 3.9. Consider a cyclic relatively nonexpansive mapping $T : H \cup C \rightarrow H \cup C$ such that for any nonempty, bounded, closed, convex, proximal and T -invariant pair (ι_1, ι_2) with $\text{dist}(\iota_1, \iota_2) = \text{dist}(H, C)$,

$$2^{\acute{\Theta}} [\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)) + \acute{F}(\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)))] \leq \acute{\mathcal{G}}(\iota_1 \cup \iota_2) + \acute{F}(\acute{\mathcal{G}}(\iota_1 \cup \iota_2)). \tag{3.9}$$

Then T has a *BPP* if $H_0 \neq \emptyset$.

Proof . Putting $\acute{\varrho}(\acute{\mu}) = \acute{\mu}$ in equation (3.8) of Corollary 3.8 and using Theorem 3.5, the result follows. \square

Corollary 3.10. Consider a cyclic relatively nonexpansive mapping $T : H \cup C \rightarrow H \cup C$ such that for any nonempty, bounded, closed, convex, proximal and T -invariant pair (ι_1, ι_2) with $\text{dist}(\iota_1, \iota_2) = \text{dist}(H, C)$,

$$2^{\acute{\Theta}} [\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)) + \acute{F}(\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)))] \leq \acute{\mathcal{G}}(\iota_1 \cup \iota_2) + \acute{F}(\acute{\mathcal{G}}(\iota_1 \cup \iota_2)). \tag{3.10}$$

Then T has a *BPP* if $H_0 \neq \emptyset$.

Proof . Putting $\acute{\Theta}(\acute{\mu}) = \acute{\mu}$ in equation (3.9) of Corollary 3.9 and using Theorem 3.5, the result follows. \square

Corollary 3.11. Consider a cyclic relatively nonexpansive mapping $T : H \cup C \rightarrow H \cup C$ such that for any nonempty, bounded, closed, convex, proximal and T -invariant pair (ι_1, ι_2) with $\text{dist}(\iota_1, \iota_2) = \text{dist}(H, C)$,

$$\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)) \leq \acute{\mathcal{G}}(\iota_1 \cup \iota_2). \tag{3.11}$$

Then T has a *BPP* if $H_0 \neq \emptyset$.

Proof . Putting $\acute{F}(\acute{\mu}) = 0$ in equation (3.10) of Corollary 3.10, we get

$$\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2)) \leq 2^{\acute{w}} [\acute{\mathcal{G}}(T(\iota_1) \cup T(\iota_2))] \leq \acute{\mathcal{G}}(\iota_1 \cup \iota_2),$$

and using Theorem 3.5, the result follows. \square

4 Applications

We use our conclusions to look into the optimum solution to a system of fractional hybrid differential equation. Consider the following system of fractional hybrid differential equations:

$$D^{\acute{\nu}} [\acute{\tilde{\epsilon}}(\tau_1) - \acute{F}(\tau_1, \acute{\tilde{\epsilon}}(\tau_1))] = \mathcal{H}(\acute{\varrho}, \acute{\tilde{\epsilon}}(\acute{\varrho})), \tag{4.1}$$

$$D^{\acute{\nu}} [\acute{\tilde{\varkappa}}(\tau_1) - \acute{F}(\tau_1, \acute{\tilde{\varkappa}}(\tau_1))] = \mathcal{Z}(\acute{\varrho}, \acute{\tilde{\varkappa}}(\acute{\varrho})). \tag{4.2}$$

with $\acute{\tilde{\epsilon}}(0) = \acute{\tilde{\pi}}_0$, $\acute{\tilde{\varkappa}}(0) = \acute{\tilde{\beta}}_0$, $\acute{\tilde{\pi}}_0, \acute{\tilde{\beta}}_0 \in \mathbf{R}$, $\acute{\nu} \in (0, 1]$, $\tau_1 \in [0, 1] = I$ and $\mathcal{H}, \mathcal{Z}, \acute{F}$ are continuous functions with $\|\mathcal{H}(\acute{\varrho}, \acute{\tilde{\epsilon}}(\acute{\varrho}))\| \leq l_1$, $\|\mathcal{Z}(\acute{\varrho}, \acute{\tilde{\varkappa}}(\acute{\varrho}))\| \leq l_2$. Now, we consider the following system of integral equations equivalent to equations (4.1) and (4.2):

$$\begin{cases} \tilde{\varepsilon}(\tau_1) = \tilde{\pi}_0 - \tilde{F}(0, \tilde{\pi}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{H}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho}, \\ \tilde{\varkappa}(\tau_1) = \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho})) d\tilde{\varrho}, \end{cases} \tag{4.3}$$

for $\tau_1 \in [0, 1] = I$. Also, assume that $(R, \|\cdot\|)$ is a Banach space and two closed ball $\tilde{P}_1 = E_1(\tilde{\pi}_0, \tilde{b})$, $\tilde{P}_2 = E_2(\tilde{\beta}_0, \tilde{b})$ in R , with $\tilde{b} \in [0, \infty)$. Consider a standard Banach space $N = C(\mathfrak{F}, R)$ of continuous function with supremum norm for $\mathfrak{F} \subseteq I$. Let

$$N_1 = C(\mathfrak{F}, \tilde{P}_1) = \{\tilde{\varepsilon} : \mathfrak{F} \rightarrow \tilde{P}_1 : \tilde{\varepsilon} \in N\}, \quad \text{and} \quad N_2 = C(\mathfrak{F}, \tilde{P}_2) = \{\tilde{\varkappa} : \mathfrak{F} \rightarrow \tilde{P}_2 : \tilde{\varkappa} \in N\}.$$

Then (N_1, N_2) is NBCC pair in N . Now for every $\tilde{\varepsilon} \in N_1, \tilde{\varkappa} \in N_2$,

$$\|\tilde{\varepsilon} - \tilde{\varkappa}\| = \sup_{\tau_1 \in I} \|\tilde{\varepsilon}(\tau_1) - \tilde{\varkappa}(\tau_1)\| \geq \|\tilde{\pi}_0 - \tilde{\beta}_0\|.$$

Thus $\text{dist}(N_1, N_2) = \|\tilde{\pi}_0 - \tilde{\beta}_0\|$. Now, we define $T : N_1 \cup N_2 \rightarrow N$ such that,

$$T(\tilde{\varepsilon}(\tau_1)) = \begin{cases} \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho}, & \tilde{\varepsilon} \in N_1, \\ \tilde{\pi}_0 - \tilde{F}(0, \tilde{\pi}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{H}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho}, & \tilde{\varepsilon} \in N_2. \end{cases}$$

Clearly, T is cyclic, and if $\|\tilde{r}_0 - T(\tilde{r}_0)\| = \text{dist}(N_1, N_2)$, for $\tilde{r}_0 \in N_1 \cup N_2$, then \tilde{r}_0 is the optimum solution of the system (4.3). Thus \tilde{r}_0 is the BPP for the operator T .

Theorem 4.1. [4] Consider $\acute{U} \in C[\vartheta_1, \vartheta_2]$ with $\vartheta_1 < \vartheta_2$. Let \acute{V} is Lebesgue integrable on $[\vartheta_1, \vartheta_2]$ and \acute{V} does not change its sign in $[\vartheta_1, \vartheta_2]$ with $\check{a} \in (\vartheta_1, \vartheta_2)$. Then the generalized mean value theorem of integral calculus gives,

$$\int_{\vartheta_1}^{\vartheta_2} \acute{U}(\bar{t}) \acute{V}(\bar{t}) d\bar{t} = \acute{U}(\check{a}) \int_{\vartheta_1}^{\vartheta_2} \acute{V}(\bar{t}) d\bar{t}.$$

Theorem 4.2. Consider $\acute{U} \in C[\vartheta_1, \vartheta_2]$ with $\vartheta_1 < \vartheta_2$ and $\check{\gamma} > 0$. Let \acute{V} is Lebesgue integrable on $[\vartheta_1, \vartheta_2]$ and \acute{V} does not change its sign in $[\vartheta_1, \vartheta_2]$. Then there exists $\check{a} \in (\vartheta_1, \check{\varepsilon}) \subset (\vartheta_1, \vartheta_2)$ such that $\mathcal{I}_{\vartheta_1}^{\check{\gamma}}(\acute{U}\acute{V})(\check{\varepsilon}) = \acute{U}(\check{a}) \mathcal{I}_{\vartheta_1}^{\check{\gamma}} \acute{V}(\check{\varepsilon})$, for almost every $\check{\varepsilon} \in (\vartheta_1, \vartheta_2]$. For $\check{\gamma} \geq 1$ or $\acute{V} \in C[\vartheta_1, \vartheta_2]$, the above result holds for every $\check{\varepsilon} \in (\vartheta_1, \vartheta_2]$.

Proof . Here

$$\mathcal{I}_{\vartheta_1}^{\check{\gamma}}(\acute{U}\acute{V})(\check{\varepsilon}) = \frac{1}{\check{\gamma}(\check{\gamma})} \int_{\vartheta_1}^{\tau_1} (\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} \acute{U}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) \acute{V}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho}.$$

Consider $\tilde{\acute{V}}(\check{\varepsilon}) = \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \acute{V}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho}))$. Now we have,

$$\begin{aligned} \mathcal{I}_{\vartheta_1}^{\check{\gamma}}(\acute{U}\acute{V})(\check{\varepsilon}) &= \int_{\vartheta_1}^{\tau_1} \acute{U}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) \tilde{\acute{V}}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \\ &= \acute{U}(\check{a}) \int_{\vartheta_1}^{\tau_1} \tilde{\acute{V}}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \\ &= \acute{U}(\check{a}) \int_{\vartheta_1}^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \acute{V}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \\ &= \acute{U}(\check{a}) \mathcal{I}_{\vartheta_1}^{\check{\gamma}} \acute{V}(\check{\varepsilon}). \end{aligned}$$

□

Theorem 4.3. Assume that \mathfrak{I}_0 is a *MNC* on \mathbb{N} with $0 < \check{r} \leq 1$, $\tilde{F}(\check{\mu}, \check{\nu}) = \check{\mu}\check{\nu}$, $\check{\mu}, \check{\nu} \in R$, $|\tau_1\tilde{\varepsilon}(\tau_1) - \tau_2\tilde{\varepsilon}(\tau_2)| \leq |\tau_2 - \tau_1|$, $\left\{ |\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r} + 1)} \mathfrak{l}_2 \tau_1^{\check{r}} \right\} \leq L$, for $L > 0$. Then an optimal solution exists for the system of equations (4.3) if:

For all $\tilde{\varepsilon} \in \mathbb{N}_1$, $\tilde{\varkappa} \in \mathbb{N}_2$, $\tau_1 \in \mathfrak{F}$ there exists $l \geq 0$ and $\check{s} \geq 0$ such that,

- (i) $|\tilde{\varepsilon}(\tau_1) - \tilde{\varkappa}(\tau_1)| \leq l$,
- (ii) $|\tilde{\beta}_0 - \tilde{\pi}_0| \leq \check{s}$,
- (iii) $|\mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) - \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho}))| \leq \check{\gamma}(\check{r} + 1) \left\{ |\tilde{\varepsilon} - \tilde{\varkappa}| - \check{s} - l \right\}$.

Proof . First, we show that the operator \mathbf{T} is cyclic. For $\tilde{\varepsilon} \in \mathbb{N}_1$,

$$\begin{aligned} \|\mathbf{T}(\tilde{\varepsilon}(\tau_1)) - \tilde{\beta}_0\| &= \left\| \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \tilde{\beta}_0 \right\| \\ &= \left\| \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) - \tilde{F}(0, \tilde{\beta}_0) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right\| \\ &\leq \left| \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) \right| + \left| \tilde{F}(0, \tilde{\beta}_0) \right| + \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\ &\leq |\tau_1\tilde{\varepsilon}(\tau_1)| + \mathfrak{l}_2 \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} d\tilde{\varrho} \right| \\ &\leq \tau_1|\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r})} \mathfrak{l}_2 \left| \int_0^{\tau_1} (\tau_1 - \tilde{\varrho})^{\check{r}-1} d\tilde{\varrho} \right| \\ &\leq |\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r})} \mathfrak{l}_2 \left| \left[\frac{-(\tau_1 - \tilde{\varrho})^{\check{r}}}{\check{r}} \right]_0^{\tau_1} \right| \\ &\leq |\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r} + 1)} \mathfrak{l}_2 \tau_1^{\check{r}} \\ &\leq L. \end{aligned}$$

Hence, $\mathbf{T}(\tilde{\varepsilon}(\tau_1)) \in \mathbb{N}_2$. By using a similar method, we can show that $\mathbf{T}(\tilde{\varepsilon}(\tau_1)) \in \mathbb{N}_1$ for $\tilde{\varepsilon} \in \mathbb{N}_2$. Thus, \mathbf{T} is cyclic. We now show that $\mathbf{T}(\mathbb{N}_1)$ is an equicontinuous and bounded subset of \mathbb{N}_2 . Consider $\tilde{\varepsilon} \in \mathbb{N}_1$ and $\tau_1 \in \mathfrak{F}$ with $0 < \check{r} \leq 1$. We have,

$$\begin{aligned} \|\mathbf{T}(\tilde{\varepsilon}(\tau_1))\| &= \left\| \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right\| \\ &\leq \left| \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) \right| + \left| \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) \right| + \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\ &\leq |\tilde{\beta}_0| + |\tau_1\tilde{\varepsilon}(\tau_1)| + \mathfrak{l}_2 \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{r}-1}}{\check{\gamma}(\check{r})} d\tilde{\varrho} \right| \\ &\leq |\tilde{\beta}_0| + \tau_1|\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r})} \mathfrak{l}_2 \left| \int_0^{\tau_1} (\tau_1 - \tilde{\varrho})^{\check{r}-1} d\tilde{\varrho} \right| \\ &\leq |\tilde{\beta}_0| + |\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r})} \mathfrak{l}_2 \left| \left[\frac{-(\tau_1 - \tilde{\varrho})^{\check{r}}}{\check{r}} \right]_0^{\tau_1} \right| \\ &\leq |\tilde{\beta}_0| + |\tilde{\varepsilon}(\tau_1)| + \frac{1}{\check{\gamma}(\check{r} + 1)} \mathfrak{l}_2 |\tau_1^{\check{r}}| \\ &\leq |\tilde{\beta}_0| + L. \end{aligned}$$

Therefore $\mathbf{T}(\mathbf{N}_1)$ is bounded. Suppose $\tau_1, \tau_2 \in \mathfrak{F}$, $\tau_1 > \tau_2$ and $\tilde{\varepsilon} \in \mathbf{N}_1$. Then

$$\begin{aligned}
& \|\mathbf{T}(\tilde{\varepsilon}(\tau_1)) - \mathbf{T}(\tilde{\varepsilon}(\tau_2))\| \\
&= \left\| \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \tilde{\beta}_0 + \tilde{F}(0, \tilde{\beta}_0) - \tilde{F}(\tau_2, \tilde{\varepsilon}(\tau_2)) \right. \\
&\quad \left. - \int_0^{\tau_2} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right\| \\
&\leq \left| \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) - \tilde{F}(\tau_2, \tilde{\varepsilon}(\tau_2)) \right| + \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \int_0^{\tau_2} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\
&\leq |\tau_1 \tilde{\varepsilon}(\tau_1) - \tau_2 \tilde{\varepsilon}(\tau_2)| + \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \int_0^{\tau_1} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} + \right. \\
&\quad \left. + \int_0^{\tau_1} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \int_0^{\tau_2} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\
&\leq |\tau_1 \tilde{\varepsilon}(\tau_1) - \tau_2 \tilde{\varepsilon}(\tau_2)| + \left| \int_0^{\tau_1} \frac{\{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} - (\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}\}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} + \int_{\tau_2}^{\tau_1} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\
&\leq |\tau_1 \tilde{\varepsilon}(\tau_1) - \tau_2 \tilde{\varepsilon}(\tau_2)| + \left| \int_0^{\tau_1} \frac{\{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} - (\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}\}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| + \left| \int_{\tau_2}^{\tau_1} \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right| \\
&\leq |\tau_1 \tilde{\varepsilon}(\tau_1) - \tau_2 \tilde{\varepsilon}(\tau_2)| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \mathbf{l}_2 \left| \int_0^{\tau_1} \{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} - (\tau_2 - \tilde{\varrho})^{\check{\gamma}-1}\} d\tilde{\varrho} \right| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \mathbf{l}_2 \left| \int_{\tau_2}^{\tau_1} (\tau_2 - \tilde{\varrho})^{\check{\gamma}-1} d\tilde{\varrho} \right| \\
&\leq |\tau_2 - \tau_1| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \mathbf{l}_2 \left| \left\{ \frac{-(\tau_1 - \tilde{\varrho})^{\check{\gamma}}}{\check{\gamma}} + \frac{(\tau_2 - \tilde{\varrho})^{\check{\gamma}}}{\check{\gamma}} \right\}_0^{\tau_1} \right| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \mathbf{l}_2 \left| \left\{ \frac{-(\tau_2 - \tilde{\varrho})^{\check{\gamma}}}{\check{\gamma}} \right\}_{\tau_2}^{\tau_1} \right| \\
&\leq |\tau_2 - \tau_1| + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \mathbf{l}_2 \left| \left\{ -(\tau_1 - \tilde{\varrho})^{\check{\gamma}} + (\tau_2 - \tilde{\varrho})^{\check{\gamma}} \right\}_0^{\tau_1} \right| + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \mathbf{l}_2 \left| \left\{ -(\tau_2 - \tilde{\varrho})^{\check{\gamma}} \right\}_{\tau_2}^{\tau_1} \right| \\
&= |\tau_2 - \tau_1| + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \mathbf{l}_2 \left| \left\{ (\tau_2 - \tau_1)^{\check{\gamma}} + (\tau_1)^{\check{\gamma}} - (\tau_2)^{\check{\gamma}} \right\} - (\tau_2 - \tau_1)^{\check{\gamma}} \right| \\
&= |\tau_2 - \tau_1| + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \mathbf{l}_2 \left| \left\{ (\tau_1)^{\check{\gamma}} - (\tau_2)^{\check{\gamma}} \right\} \right|.
\end{aligned}$$

As $\tau_2 \rightarrow \tau_1$, $\|\mathbf{T}(\tilde{\varepsilon}(\tau_1)) - \mathbf{T}(\tilde{\varepsilon}(\tau_2))\| \rightarrow 0$. That is $\mathbf{T}(\mathbf{N}_1)$ is equicontinuous. We can show that $\mathbf{T}(\mathbf{N}_2)$ is equicontinuous and bounded in \mathbf{N}_1 with the similar manner. Thus by Arzela-Ascoli theorem, we conclude that $(\mathbf{N}_1, \mathbf{N}_2)$ is a relatively compact pair. Now for each $(\tilde{\varepsilon}, \tilde{\varkappa}) \in \mathbf{N}_1 \times \mathbf{N}_2$,

$$\begin{aligned}
& \|\mathbf{T}(\tilde{\varepsilon}(\tau_1)) - \mathbf{T}(\tilde{\varkappa}(\tau_1))\| \\
&= \left\| \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \tilde{\pi}_0 + \tilde{F}(0, \tilde{\pi}_0) - \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) \right. \\
&\quad \left. - \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho})) d\tilde{\varrho} \right\| \\
&\leq \left| \tilde{\beta}_0 - \tilde{\pi}_0 \right| + \left| \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) - \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) \right| + \left| \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} - \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\tilde{\gamma}(\check{\gamma})} \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho})) d\tilde{\varrho} \right| \\
&\leq \check{s} + \left| \tau_1 \tilde{\varepsilon}(\tau_1) - \tau_1 \tilde{\varkappa}(\tau_1) \right| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \left| \int_0^{\tau_1} (\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} \left\{ \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) - \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho})) \right\} d\tilde{\varrho} \right| \\
&\leq \check{s} + \left| \tilde{\varepsilon}(\tau_1) - \tilde{\varkappa}(\tau_1) \right| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \tilde{\gamma}(\check{\gamma} + 1) \left| \int_0^{\tau_1} (\tau_1 - \tilde{\varrho})^{\check{\gamma}-1} \left\{ |\tilde{\varepsilon} - \tilde{\varkappa}| - \check{s} - l \right\} d\tilde{\varrho} \right| \\
&\leq \check{s} + \left| \tilde{\varepsilon}(\tau_1) - \tilde{\varkappa}(\tau_1) \right| + \frac{1}{\tilde{\gamma}(\check{\gamma})} \tilde{\gamma}(\check{\gamma} + 1) \left\{ |\tilde{\varepsilon} - \tilde{\varkappa}| - \check{s} - l \right\} \left| \left\{ \frac{-(\tau_1 - \tilde{\varrho})^{\check{\gamma}}}{\check{\gamma}} \right\}_0^{\tau_1} \right| \\
&\leq \check{s} + l + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \tilde{\gamma}(\check{\gamma} + 1) \left\{ |\tilde{\varepsilon} - \tilde{\varkappa}| - \check{s} - l \right\} |(\tau_1)^{\check{\gamma}}| \\
&\leq \check{s} + l + \frac{1}{\tilde{\gamma}(\check{\gamma} + 1)} \tilde{\gamma}(\check{\gamma} + 1) \left\{ |\tilde{\varepsilon} - \tilde{\varkappa}| - \check{s} - l \right\} \\
&= |\tilde{\varepsilon} - \tilde{\varkappa}| \\
&\leq \|\tilde{\varepsilon} - \tilde{\varkappa}\|.
\end{aligned}$$

Thus \mathbf{T} is relatively nonexpansive. Now, we assume that the pair $(\iota_1, \iota_2) \subseteq (\mathbf{N}_1, \mathbf{N}_2)$ is a *NBCC*, \mathbf{T} -invariant, proximal pair

and $dist(\iota_1, \iota_2) = dist(N_1, N_2)$. Using theorem 4.2, we get,

$$\begin{aligned} \beth_0(\mathbf{T}(\iota_1) \cup \mathbf{T}(\iota_2)) &= \max \left\{ \beth_0(\mathbf{T}(\iota_1)), \beth_0(\mathbf{T}(\iota_2)) \right\} \\ &\leq \max \left\{ \sup_{\tau_1 \in I} \left\{ \beth_0(\{\mathbf{T}\tilde{\varepsilon}(\tau_1) : \tilde{\varepsilon} \in \iota_1\}) \right\}, \sup_{\tau_1 \in I} \left\{ \beth_0(\{\mathbf{T}\tilde{\varkappa}(\tau_1) : \tilde{\varkappa} \in \iota_2\}) \right\} \right\} \\ &= \max \left\{ \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) d\tilde{\varrho} \right\} \right) \right\}, \right. \\ &\quad \left. \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tilde{\pi}_0 - \tilde{F}(0, \tilde{\pi}_0) + \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) + \int_0^{\tau_1} \frac{(\tau_1 - \tilde{\varrho})^{\check{\gamma}-1}}{\check{\gamma}(\check{\gamma})} \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho})) d\tilde{\varrho} \right\} \right) \right\} \right\} \\ &= \max \left\{ \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tilde{\beta}_0 - \tilde{F}(0, \tilde{\beta}_0) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \frac{1}{\check{\gamma}(\check{\gamma}+1)} \mathcal{Z}(\tilde{i}, \tilde{\varepsilon}(\tilde{i})) (\tau_1)^{\check{\gamma}} : \tilde{i} \in (0, \tau_1) \right\} \right) \right\}, \right. \\ &\quad \left. \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tilde{\pi}_0 - \tilde{F}(0, \tilde{\pi}_0) + \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) + \frac{1}{\check{\gamma}(\check{\gamma}+1)} \mathcal{H}(\tilde{i}, \tilde{\varkappa}(\tilde{i})) (\tau_1)^{\check{\gamma}} : \tilde{i} \in (0, \tau_1) \right\} \right) \right\} \right\} \\ &\leq \max \left\{ \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tau_1 \tilde{\varepsilon}(\tau_1) + \frac{1}{\check{\gamma}(\check{\gamma}+1)} \mathcal{Z}(\tilde{i}, \tilde{\varepsilon}(\tilde{i})) (\tau_1)^{\check{\gamma}} : \tilde{i} \in (0, \tau_1) \right\} \right) \right\}, \right. \\ &\quad \left. \sup_{\tau_1 \in I} \left\{ \beth_0 \left(\left\{ \tau_1 \tilde{\varkappa}(\tau_1) + \frac{1}{\check{\gamma}(\check{\gamma}+1)} \mathcal{H}(\tilde{i}, \tilde{\varkappa}(\tilde{i})) (\tau_1)^{\check{\gamma}} : \tilde{i} \in (0, \tau_1) \right\} \right) \right\} \right\} \\ &\leq \max \left\{ \beth_0 \left(\left\{ \iota_1 + \mathcal{Z}(\tilde{i}, \tilde{\varepsilon}(\tilde{i})) \right\} \right), \beth_0 \left(\left\{ \iota_2 + \mathcal{H}(\tilde{i}, \tilde{\varkappa}(\tilde{i})) \right\} \right) \right\} \leq \beth_0(\iota_1 \cup \iota_2). \end{aligned}$$

Thus by using Corollary 3.11, \mathbf{T} has a *BPP*. Hence, the system (4.3) has $s \in N_1 \cup N_2$ as an optimal solution. \square

Example 4.4. Assume the following system of equations with $\|\mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho}))\| \leq 1$, $\|\mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho}))\| \leq 1$, $1 = 1$, $\check{s} = 1$, $\check{r} = 1$, $\frac{1}{2}\|\tilde{\varepsilon} - \tilde{\varkappa}\| \leq \|\tilde{\varepsilon} - \tilde{\varkappa}\| - 2$ and $\tilde{\tau}_1 \in \mathfrak{F} = [0, 1]$ as

$$\begin{aligned} \tilde{\varepsilon}(\tau_1) &= \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) - \tilde{F}(0, 0) + \int_0^{\tau_1} \frac{1}{\check{\gamma}(1)} \sin\left(\frac{\tilde{\varepsilon}}{2}\right) d\tilde{\varrho}, \\ \tilde{\varkappa}(\tau_1) &= 1 - \tilde{F}(0, 1) + \tilde{F}(\tau_1, \tilde{\varkappa}(\tau_1)) + \int_0^{\tau_1} \frac{1}{\check{\gamma}(1)} \sin\left(\frac{\tilde{\varkappa}}{2}\right) d\tilde{\varrho}. \end{aligned}$$

Consider $N_1 = \{\tau_1\}$ and $N_2 = \{\tau_1 + 1\}$ on $\mathfrak{F} = [0, 1]$. Define an operator $\mathbf{T} : N_1 \cup N_2 \rightarrow N$ such that,

$$\mathbf{T}(\tilde{\varepsilon}(\tau_1)) = \begin{cases} 1 - \tilde{F}(0, 1) + \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) + \int_0^{\tau_1} \frac{1}{\check{\gamma}(1)} \sin\left(\frac{\tilde{\varepsilon}}{2}\right) d\tilde{\varrho}, & \tilde{\varepsilon} \in N_1, \\ \tilde{F}(\tau_1, \tilde{\varepsilon}(\tau_1)) - \tilde{F}(0, 0) + \int_0^{\tau_1} \frac{1}{\check{\gamma}(1)} \sin\left(\frac{\tilde{\varepsilon}}{2}\right) d\tilde{\varrho}, & \tilde{\varepsilon} \in N_2. \end{cases}$$

Here $\mathcal{H} : \mathfrak{F} \times \iota_2 \rightarrow \iota_2$, and $\mathcal{Z} : \mathfrak{F} \times \iota_1 \rightarrow \iota_1$ with $\iota_1 = \{\tau_1\}$, $\iota_2 = \{\tau_1 + 1\}$ on $\mathfrak{F} = [0, 1]$. Now for $\tilde{\varepsilon} \in N_1$, $\tilde{\varkappa} \in N_2$, we have,

$$\begin{aligned} \|\mathcal{Z}(\tilde{\varrho}, \tilde{\varepsilon}(\tilde{\varrho})) - \mathcal{H}(\tilde{\varrho}, \tilde{\varkappa}(\tilde{\varrho}))\| &= \left\| \sin\left(\frac{\tilde{\varepsilon}}{2}\right) - \sin\left(\frac{\tilde{\varkappa}}{2}\right) \right\| \\ &\leq \left\| 2 \cos\left(\frac{\tilde{\varepsilon} + \tilde{\varkappa}}{2}\right) \sin\left(\frac{\tilde{\varepsilon} - \tilde{\varkappa}}{2}\right) \right\| \\ &\leq 2 \left\| \cos\left(\frac{\tilde{\varepsilon} + \tilde{\varkappa}}{2}\right) \right\| \left\| \sin\left(\frac{\tilde{\varepsilon} - \tilde{\varkappa}}{2}\right) \right\| \\ &\leq 2 \left\| \frac{\tilde{\varepsilon} - \tilde{\varkappa}}{2} \right\| \leq \frac{1}{2} \|\tilde{\varepsilon} - \tilde{\varkappa}\| \\ &\leq \|\tilde{\varepsilon} - \tilde{\varkappa}\| - 2. \end{aligned}$$

Since the above system of equations satisfies all the conditions of theorem (4.3). Therefore $r \in \{0, 1\} = \mathbb{N}_1 \cup \mathbb{N}_2$ is the optimal solution for the above system of equations at $\tau_1 = 0$ as $\|r - T(r)\| = 1 = \text{dist}(\mathbb{N}_1, \mathbb{N}_2)$.

5 Conclusion

This paper established a best proximity point theorem with the help of newly defined contraction operator by using *MNC*, and we used it to determine the existence of optimum solutions of the system of fractional hybrid differential equations. The approach used in this article can therefore be used to obtain the existence of optimum solution of other fractional differential equations.

References

- [1] R.R. Akhmerov, M.I. Kamenskii, A.S. Potapov, A.E. Rodkina, and B.N. Sadovskii, *Measures of Noncompactness and Condensing Operators*, Vol. 55, Birkhäuser, Basel, 1992.
- [2] J. Banaś and K. Goebel, *Measure of Noncompactness in Banach Spaces*, Lecture Notes in Pure and Applied Mathematics, Vol. 60, Marcel Dekker, New York, 1980.
- [3] A. Das, M. Paunović, V. Parvaneh, M. Mursaleen, and Z. Bagheri, *Existence of a solution to an infinite system of weighted fractional integral equations of a function with respect to another function via a measure of noncompactness*, *Demonst. Math.* **56** (2023), no. 1, 20220192.
- [4] K. Diethelm, *The mean value theorem and a Nagumo-type uniqueness theorem for Caputo's fractional calculus*, *Fractional Cal. Appl. Anal.* **15** (2012), 304–313.
- [5] M.A.E. Herzallah and D. Baleanu, *On fractional order hybrid differential equations*, *Abstr. Appl. Anal.* **2014** (2014), Article ID 389386, 7 pages.
- [6] M. Gabeleh and J. Markin, *Global optimal solutions of a system of differential equations via measure of noncompactness*, *Filomat* **35** (2021), no. 15, 5059–5071.
- [7] B. Hazarika, H.M. Srivastava, R. Arab, and M. Rabbani, *Existence of solution for an infinite system of nonlinear integral equations via measure of noncompactness and homotopy perturbation method to solve it*, *J. Comput. Appl. Math.* **343** (2018), no. 1, 341–352.
- [8] B. Hazarika, H. M. Srivastava, R. Arab, and M. Rabbani, *Application of simulation function and measure of noncompactness for solvability of nonlinear functional integral equations and introduction to an iteration algorithm to find solution*, *Appl. Math. Comput.* **360** (2019), no. 1, 131–146.
- [9] S.A. Mohiuddine, H.M. Srivastava, and A. Alotaibi, *Application of measures of noncompactness to the infinite system of second-order differential equations in ℓ_p space*, *Adv. Difference Equ.* **2016** (2016), no. 1, 317.
- [10] H. K. Nashine, R. Arab, P.R. Patle, and D.K. Patel, *Best proximity point results via measure of noncompactness and application*, *Numerical Funct. Anal. Optim.* **42** (2021), no. 4, 430–442.
- [11] P.R. Patle, M. Gabeleh, V. Rakočević, and M.E. Samei, *New best proximity point(pair) theorems via MNC and application to the existence of optimum solutions for a system of ψ -Hilfer fractional differential equations*, *Rev. Real Acad. Cienc. Exactas Fis. Nat. Ser. A-Mat.* **117** (2023), 124.
- [12] H.M. Srivastava, A. Das, B. Hazarika, and S. A. Mohiuddine, *Existence of solutions of infinite systems of differential equations of general order with boundary conditions in the spaces c_0 and l_1 via the measure of noncompactness*, *Math. Meth. Appl. Sci.* **41** (2018), no. 10, 3558–3569.
- [13] H.M. Srivastava, A. Deep, S. Abbas, and B. Hazarika, *Solvability for a class of generalized functional-integral equations by means of Petryshyn's fixed point theorem*, *J. Nonlinear Convex Anal.* **22** (2021), no. 12, 2715–2737.